

RESEARCH ARTICLE

Open Access



Integrative analysis of transcriptomics and proteomics of skeletal muscles of the Chinese indigenous Shaziling pig compared with the Yorkshire breed

Hu Yang^{1,2}, Xing-li Xu², Hai-ming Ma^{1*} and Jun Jiang¹

Abstract

Background: The Shaziling pig (*Sus scrofa*) is a well-known indigenous breed in China. One of its main advantages over European breeds is its high meat quality. However, little genetic information is available for the Shaziling pig. To screen for differentially expressed genes and proteins that might be responsible for the meat quality, the *longissimus dorsi* muscles from Shaziling and Yorkshire pig breeds were investigated using an integrative analysis of transcriptomics and proteomics, involving high-throughput sequencing, the two-dimensional gel electrophoresis, and mass spectrometry.

Results: Sequencing produced 79,320 unigenes by *de novo* assembly, and 488 differentially expressed genes in the *longissimus dorsi* muscle of Shaziling pig compared with the Yorkshire breed were identified. Gene Ontology term enrichment of biological functions and Kyoto Encyclopedia of Genes and Genomes analysis showed that the gene products were mainly involved in metabolism, protein binding, and regulation of skeletal muscle development. At the protein level, 23 differentially expressed proteins were identified, which were potentially associated with fatty acid metabolism, the glycolytic pathway, and skeletal muscle growth. Eight differentially expressed genes were confirmed by real-time PCR. These results give an insight into the mechanisms underlying the formation of skeletal muscle in the Shaziling pig.

Conclusions: Certain differentially expressed genes and proteins are involved in fatty acid metabolism, intramuscular fat deposition, and skeletal muscle growth in the Shaziling pig. These results provide candidate genes for improving meat quality and will promote further transcriptomic research in Shaziling pigs.

Keywords: Shaziling Pig, RNA-seq, Proteomics, Meat quality

Background

Pigs are important domestic animals used for meat production worldwide. Pork quality is influenced by many factors, including breed, nutrition, and post-slaughter handling [1]. Among these factors, the breed is the most important. There are more pig breeds in China than in any other country in the world [2]. In China, 118 indigenous pig breeds are listed as the Domestic Animal Diversity in the World index [3]. Over a long period,

breeders have attempted to increase muscle yield and decrease carcass fatness, and great progress for these traits has been made in swine breeding. For example, Landrace pigs and Yorkshire pigs grow more quickly and have higher lean meat than other pig breeds. However, some studies suggest that such intensive selection for increased muscle growth and decreased carcass fatness has led to a deterioration in meat quality [4–6]. Compared with European breeds, Chinese native breeds have higher intramuscular fat (IMF), and increased tenderness and meat quality [7–10]. The Shaziling pig is a well-known breed, which has evolved for centuries in Hunan Province, China, where it feeds on grains, tubers,

* Correspondence: mahaiming2000@163.com

¹College of Animal Science and Technology, Hunan Agricultural University, Changsha 410128, People's Republic of China

Full list of author information is available at the end of the article



and wild herbs. It is a Chinese fat-type line with high IMF, superior meat quality, and strong resistance to general diseases. Compared with the Shaziling breed, the Yorkshire has a faster growth rate and a leaner meat ratio.

Skeletal muscle contains several fiber types [11]. Muscle fiber types and the proportion of fiber types affect meat quality directly [12]. Therefore, studies on development and growth are beneficial to improve meat quality. Skeletal muscle development is very complicated, and comprises several stages: determination of myoblasts, proliferation of myoblasts, differentiation and fusion of myoblasts into myotubes and myofibers, and growth and maturation [13, 14]. Analyzing these stages would provide a good basis for understanding muscle fiber development. Previous reports showed that muscle fibers could be classified into red and white fibers [15]. These compositional differences between fibers determine their distinct metabolic type and physiological functions and affect meat quality [16, 17]. Skeletal muscles have been explored extensively using molecular biology [18, 19], and proteomics and transcriptomics techniques have been applied to study the porcine skeletal muscle of different breeds [20, 21].

Over the past decade, proteomic technologies have been used successfully to study skeletal muscle [11, 22]. Proteomic analysis based on two-dimensional gel electrophoresis (2-DE) and mass spectrometry (MS) is a classical method in quantitative proteomics to separate mixtures of proteins into two dimensions and has been a powerful tool in meat science [23, 24]. Many reports concerning differential proteomics among different pig breeds have been published [25–27]. Another technology for characterizing molecular changes in skeletal muscle is analysis of the transcriptome. Recently, transcriptome studies have been applied to analyze differentially expressed genes (DEGs), identify novel genes, describe metabolic pathways, and forecast the relationship between genotypes and phenotypes [28–30]. Next-generation sequencing technology has provided a new tool to quantify transcriptomes and analyze gene expression on a global scale. Transcriptional and proteomic methods could be used to analyze the changes from the mRNA expression to the protein abundance. In addition, post-transcriptional regulation is very important for mRNA stability, translation initiation, and protein stability [31]. Thus, it is necessary to combine proteomic and transcriptional methods simultaneously to analyze skeletal muscle growth and development. Currently, several reports have presented results of both proteomic and transcriptional analyses. For instance, *longissimus dorsi* muscle (LM) proteome and transcriptome profiles of Yorkshire pig and Casertana pig breeds were compared using 2-DE and a microarray. As a result, a large number of genes were identified that are involved in glycolytic metabolism and skeletal muscle growth [32]. In addition, Kim et al. [33] also compared the LM

proteome and transcriptome profiles of different pig breeds using 2-DE and a microarray.

In the present study, we performed transcriptomic and proteomic analyses, along with functional enrichment of Gene Ontology (GO) terms and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis, to characterize the expression profiles in the LM of Shaziling and Yorkshire pigs. The aim of this study was to reveal the differences in breed-related protein and transcript expression patterns between the two breeds. These results could provide an insight into the mechanisms of growth and development of porcine skeletal muscle.

Methods

Sample preparation

Three 25-day-old, male full-sibs for each of Shaziling pigs and Yorkshire pigs were slaughtered following the Animal Care Guidelines of the Ethics committee of Hunan Agricultural University. Samples of LM were collected from the same area of the ribs and placed in liquid nitrogen immediately. All samples were kept at -80°C after collection until used.

Protein extraction

The frozen LM tissues (100 mg) from Shaziling pigs and Yorkshire pigs were crushed in a mortar containing liquid nitrogen and were then sonicated for 10 s using a Sonoplus (Bandelin Electronic, Berlin Germany). The crushed tissue was homogenized in 1 mL of cold dissolution buffer containing 7 M urea, 2 M thiourea, 1 % dithiothreitol (DTT), 4 % (w/v) CHAPS, 20 μL protease inhibitor cocktail (BBI, Canada), and 2 % (v/v) pharmalyte (pH 3–10; BioRad, Hercules, CA, USA). The homogenate was centrifuged at $15,000 \times g$, for 20 min at 4°C . The supernatant fraction was filtered and kept at -80°C for subsequent analysis. The total protein content was determined using a Bradford assay kit (Bio-Rad).

2-DE and images analysis of gel

Approximately 100 μg of extracted protein was diluted with rehydration buffer [8 M urea, 2 M thiourea, 50 mM DTT, 4 % (w/v) CHAPS, and 0.5 % carrier ampholytes (pH 3–10, Bio-Rad)]. The mix was loaded onto 13-cm, nonlinear, immobilized pH gradient strips (IPG, pH 3–10; BioRad), which were rehydrated overnight. After rehydration, first-dimension isoelectric focusing was carried out at 20°C at 30 V for 12 h, 500 V for 1 h, 1000 V for 1 h, and then gradually increased to 8,000 V. Isoelectric focusing was performed using an Ettan IPGphor III system (GE Healthcare, USA) gel apparatus. The focused IPG strips were incubated for 15 min in equilibration buffer containing 6 M urea, 50 mM Tris-HCl, pH 8.8, 30 % glycerol, 2 % sodium dodecyl sulfate (SDS), and 1 % DTT. Then, strips were then incubated again for

15 min in a buffer containing 6 M urea, 50 mM Tris-HCl, pH 8.8, 2 % SDS, 30 % glycerol, and 4 % iodoacetamide. In the second dimension, the equilibrated IPG strips were placed onto SDS polyacrylamide gels (12.5 % T, 2.6 % C) for protein separation using the Ettan DALT six electrophoresis system (GE Healthcare). When the bromophenol blue dye front reached the bottom of the gel, electrophoresis was stopped and analytical gels were subjected to silver staining and Coomassie blue was used to stain preparative gels before identification by MS. Each sample was analyzed in triplicate.

Spot choosing and tryptic digestion

Gel images were scanned using an Image Scanner UMax Powerlook 2110XL (GE Amersham) and analyzed using Image Master 2D Platinum software Hofer SE 600 (Version 5.0; GE Amersham). The protein spots were compared automatically using the software, including matching and quality. Spots whose average density was different by more than 1.5-fold between the two pig breeds were analyzed and marked. The differentially abundant spots were cut out from the preparative gel carefully and washed twice with 200 μ L of 50 mM ammonium bicarbonate with 50 % acetonitrile and incubated for 15 min at room temperature. The gel pieces were swollen in a digestion solution containing 5 μ L of 25 mM ammonium bicarbonate and 10 ng of trypsin at 4 °C for 30 min; in-gel tryptic digestion was run overnight at 37 °C. Subsequently, the supernatant was extracted twice with solvent A (80 % acetonitrile, 0.1 % trifluoroacetic acid) for 15 min at 3 °C. Finally, the digested tryptic peptides were passed through a Zip-Tip to remove salts, according to the manufacturer's protocol.

Mass spectrometry

The samples were stored at -70 °C before analysis by MALDI-TOF/TOF. Protein identification was performed using a 5800 MALDI-TOF/TOF mass spectrometer (AB SCIEX) according to the manufacturer's instructions. Mass spectra were acquired in reflector mode, and recorded in the range of 800–4,500 Da. Eight of the most intense ion signals were selected as precursors for the acquisition of MS/MS. The resulting peptide masses were submitted into the database of the National Center for Biotechnology Information non-redundant (NCBI nr) and the Swiss-Prot database using the Mascot server (Matrix Science, London, UK) to identify proteins. The search parameters were set at $\pm$ 100 ppm for peptide-mass mapping (PMF), peptide tolerance and $\pm$ 0.4 Da for the MS/MS tolerance.

RNA isolation and transcriptome analysis

Total RNA was extracted from the LM of the two breeds of the pig using the total RNA extraction kit

(Qiagen, Valencia, CA, USA), in accordance with the manufacturer's instructions. The RNA quantity and integrity were checked using a NanoDrop 2000 spectrophotometer and bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA). A TruSeq RNA Sample Preparation kit v2 (Illumina, San Diego, CA, USA) was used to construct cDNA libraries. Subsequently, the libraries were sequenced using an Illumina HiSeq 2500 instrument (Illumina, San Diego, CA, USA) that generated paired-end reads of 100 bp.

De novo assembling and functional annotation of reads

Raw sequencing reads of each sample were trimmed and assembled *de novo* using CLC Genomics Workbench (CLC Bio, Aarhus, Denmark). After the adapter sequences, ambiguous bases and sequences less than 20 nucleotides were removed, credible contigs generated by *de novo* sequence assembly. The assembled contigs were annotated against the UniProt and NCBI non-redundant (nr) database using BlastX alignment with an E-value cut off of $<1e-5$. Based on the annotation results, GO terms were extracted using Blast2GO (<http://www.blast2go.com>) and the results were classified as biological processes, cellular components, and molecular functions. The EuKaryotic Orthologous Groups (KOG) and KEGG databases were used to predict the functions and define the main metabolic pathways, separately.

Gene expression quantification and differential expression analysis

The expression level of each gene was calculated using the reads per kb per million reads values by the Qualimap v0.5 software [34]. The DEGseq program [35] and R packages were used to identify significantly DEGs between Shaziling pigs and Yorkshire pigs. In this study, the criteria were a fold change (FC) greater than two and cut-off of the false discovery rate of 5 %. For the unigenes that were considered as having differential expression, GO functional enrichment was carried out when the P value was less than 0.05. DEGseq provides statistical routines for determining DEGs.

Validation of DEGs by quantitative real-time PCR (qPCR)

To validate the expression level of the DEGs, eight DEGs from 2-DE results were analyzed by using qPCR. These genes included Enolase 3 (*ENO3*); ATP synthase subunit beta, mitochondrial (*ATP5B*); Enolase 1 (*ENO1*); Triose-phosphate Isomerase (*TPI*); alpha-actin (*ACTA1*); actin, alpha, cardiac muscle 1 (*ACTC1*); myosin light chain, phosphorylatable, fast skeletal muscle (*MYLPF*); and heat shock protein 27 (*HSPB1*), which mainly belonged to metabolic pathways and structural proteins. Total RNA was extracted from the LM of Shaziling and Yorkshire pigs using the Trizol reagent (Invitrogen, CA, USA),

according to the manufacturer's instructions, and the RNA was treated with RNase-free DNase (Qiagen, Valencia, CA, USA). The quality and concentration of the total RNA was evaluated by gel electrophoresis. The cDNA synthesis of samples was performed using the Quantitect Reverse Transcription kit (Qiagen). The primers (Table 1) were designed by the Roche Universal Probe Library Assay Design Center (<https://www.roche-applied-science.com>) and synthesized by Sangon Biotechnology Corporation (Shanghai, China). qPCR was performed on an Applied Biosystems 7300 Real time PCR system (Applied Biosystems, USA) with the SYBR Premix Ex Taq Kit (Takara Biotechnology, Japan) in a 20- μ L PCR mix. *GAPDH* was used as the reference gene to measure the expression levels of mRNA between the samples and data were calculated by the $2^{-\Delta\Delta C_t}$ comparative CT method [36].

Results and discussion

Identification of differentially abundant proteins by 2-DE and MS

The *Longissimus dorsi* proteomes from the two pig breeds were analyzed by the 2-DE technique. Thirty-eight protein spots (Fig. 1) were significantly different ($FC > 1.5$) between the two breeds: 27 protein spots were upregulated in Shaziling pigs and 11 in Yorkshire. The differentially abundant protein spots were identified by MALDI-TOF/TOF-MS: two spots could not be identified successfully; however, the 36 remaining spots were identified by matching peptide data to the UniProt database. The characteristics of the identified proteins and the identification parameters are listed in Table 2. Some of the identified proteins were resolved as multiple spots. For example, spots 2480, 2654, and 2659 represented the

same protein and showed similar quantitative trends. Ultimately, 23 different proteins were identified.

Function analysis of differentially expressed proteins


To further determine the functions of the differentially abundant proteins, functional categorization of all the identified proteins was performed using GO annotation. The annotated results of biological process are shown in Fig. 2. The 23 differential proteins were grouped into 11 categories. We focused on the categories of fatty acid metabolism, glycolytic pathway, and skeletal muscle growth and development. The result of enrichment analysis revealed that some of the identified proteins had an important impact on meat quality. For example, TPI plays an important role in ensuring immediate equilibration of the triosephosphates produced by aldolase in glycolysis, which is interconnected with lipid metabolism and to the pentose phosphate pathway [37]. In addition, a previous report showed that a quantitative increase in TPI was positively correlated with more tender meat [38].

Enolase 3 was another metabolic enzyme that was differentially abundant between the two breeds. Enolase 3 is a crucial enzyme in glycolysis that catalyzes the interconversion of diphosphoglycerate and phosphoenolpyruvate. Pig muscle Enolase 3 was investigated some time ago [39]. Many isoforms of Enolase 3 have been confirmed to have an influence on IMF in pigs [20, 40].

Muscles are composed primarily of different muscle fibers. Muscle fiber type is an important factor influencing meat quality [41]. For example, increasing the percentage of type IIb fibers could lead to the reduced meat quality because of altered metabolic rates and biochemical processes [42]. ACTA1 is a member of the actin family and is a major constituent of the contractile apparatus in skeletal

Table 1 Primer sequences for the quantitative real-time PCR amplification of the differential expressed genes in Shaziling and Yorkshire pigs

Genes	Primer sequences (5'-3')	Product size	GenBank sequence no.
HSPB1	F: CGGCAGGATGAGCACGGCTTCA R: GCGCCTCGAAAGTGACAGGGATGG	184 bp	gi 55926209
ENO1	F: GGGGCTCAACTGGGATCTACGA R: TCCGTGCCGTCCATCTCAATCA	191 bp	gi 753703906
ATP5B	F: CCCTTCTGCGGTGGGTAT R: CACGGGACAGCACAGTGGTAG	188 bp	gi 89574051
TPI1	F: CAGAGCACCCGCATCATTTACG R: AAGCGCCACCCACAAGGAAC	100 bp	gi 262263205
ACTC1	F: GGGGATGGCGTAACCCACA R: GGCAAGGCATAGCCCTCGTAA	50 bp	gi 545801458
MYLPF	F: GGCGGCAACGTGGACTACAA R: GGCCATCAAAGACCGAAGAGG	94 bp	gi 117660856
ENO3	F: CGGGAAGGACGCCACCAAT R: CCGTTGCGGTAGAACTCAGATGC	165 bp	gi 113205498
ACTA1	F: TCAGGAAGGACCTGTATGCCAACAA R: TGGACAGCGAGGCCAGGATG	186 bp	gi 268607671



muscle. Previous reports showed that alpha-actin levels positively correlate with the synthesis of muscle fiber proteins and muscle growth [43]. In this study, ACTA1 was differentially abundant between the two pig breeds.

Heat shock proteins (HSPs) including HSPA1, HSPA8, HSPB1, and other chaperone proteins have been associated with pig meat quality [44, 45]. In the present study, HSPB1 was more abundant in the Shaziling pig compared with the Yorkshire pig. This observation suggested that HSPB1 is correlated with meat quality. In addition to tenderness, further research is needed to confirm whether the IMF content of meat is affected by the expression of HSPB1.

Illumina sequencing and *de novo* assembly

Sequencing generated 30,751,992 and 35,933,274 100-bp paired-end reads from cDNA libraries of Shaziling and Yorkshire pigs, respectively. After trimming of low-quality reads and the removal of adapters, 29,522,984 and 34,937,710 reads were used for *de novo* assembly. The reads were assembled into 86,759 contigs (N50 = 939 bp) ranging from 100 to 49,881 bp, with an average length of 672 bp. Thereafter, the contigs were assembled into scaffolds (N50 = 1028 bp) with a mean length of 713 bp. After the final assembly of scaffolds using CAP3 [46], we obtained 79,320 unigenes (N50 = 1112 bp) with an average length of 733 bp and a maximum of 66,767 bp. The length distribution of the assembled unigenes is shown in Fig. 3. The result of the assembly is shown in Table 3. The transcriptome data have been submitted to the NCBI GEO database under accession number GSE70673.

Functional annotation of *longissimus dorsi* transcripts

To validate and annotate the protein functions of the 79,320 assembled unigenes, they were searched against the NCBI nr and UniProt protein database using the BLAST program (E-value cutoff $<1e^{-5}$). Of the 79,320 assembled unigenes, 23,055 sequences (29.07 %) were assigned to the nr database, and 25,784 sequences (32.50 %) were found to have significant similarity with proteins in the UniProt database (Table 4). Approximately 30 % of all assembled unigenes was perfectly matched to the Uniprot database and the nr database, respectively. Thus, most of the assembled unigenes were unknown, indicating that many new genes and non-coding RNA sequences were obtained. Generally, direct sequencing of transcriptomes is considered an effective tool to discover new genes [47, 48] and has been applied to various organisms [49, 50]. These unigenes provided much valuable information for further identification of genes related to development of the *longissimus dorsi*.

Functional classification of unigenes

GO is an extremely useful tool to classify the functions of a large quantity of genes, and is an international standardized gene functional classification system. GO analysis has been widely applied to predict the functions of genes in many organisms. Three ontologies, cellular component, molecular function, and biological process, are shared in the GO database. In this study, of the annotated unigenes, 23,702 could be assigned to the biological process (BP, 111,204 sequences), cellular components (CC, 68,072), and molecular functions (MF, 30,087) (Fig. 4) using the Blast2GO program [51]. In the biological process, the majority of unigenes were involved in catalytic activity and transcription factor activity, indicating that they might play a key role in the regulation of skeletal muscle development. In cellular components, organelle,

Table 2 Protein differentially expressed for Shaziling pig breeds and Yorkshire were identified by 2-DE and MALDI-TOF-MS/MS

Sport no	NCBI accession number	Protein name	Mr, kDa theor	pI theor	No. of peptides identified	Mascot score	Sequence coverage	Expect ^e
Up-regulated in Yorkshire pigs								
1375	gi 89574051	mitochondrial ATP synthase, H ⁺ transporting F1 complex beta subunit, partial, ATP5B	47059.6	4.99	16	1180	46 %	5.8e-11
1929	gi 283993079	L-gulonate3-dehydrogenase	35433.2	5.79	14	417	51 %	1.2e-03
2268	gi 545835136	PREDICTED: NADH dehydrogenase ubiquinone flavoprotein 2 isoform X1	25812.2	6.96	14	435	60 %	1.8e-039
2247	gi 55926209	heat shock protein beta-1	22984.7	6.23	9	423	41 %	2.9e-03
2393	gi 809283	Chain B, Structure Determination Of Aquomet Porcine Hemoglobin At 2.8 Angstrom Resolution	16082.4	6.76	12	511	86 %	4.6e-04
2377	gi 545848507	PREDICTED: alpha-crystallin B chain isoform X2	20116.4	6.76	9	212	51 %	3.7e-01
2497	gi 494389	Chain A, High Resolution X-Ray Structures Of Pig Metmyoglobin	16901.8	6.5	9	529	59 %	7.3e-04
2500	gi 494389	Chain A, High Resolution X-Ray Structures Of Pig Metmyoglobin	16901.8	6.5	12	596	82 %	1.5e-05
2575	gi 55926217	cytochrome c oxidase subunit 5B, mitochondrial precursor	14002	8.8	8	334	46 %	2.3e-02
2594	gi 809283	Chain B, Structure Determination Of Aquomet Porcine Hemoglobin At 2.8 Angstrom Resolution	16082.4	6.76	9	400	67 %	5.8e-03
Up-regulated in Shaziling pigs								
2480	gi 117660856	MYLPF	19.0	4.8	6	110	36 %	5.8e-00
2395	gi 117660874	MLC1f	21018.6	4.9	11	444	60 %	2.3e-40
2654	gi 117660856	MYLPF	19066.3	4.8	9	347	49 %	1.2e-03
2422	gi 117660890	MLC3f	16761.2	4.6	4	250	29 %	2.3e-02
2659	gi 117660856	MYLPF	19066.3	4.8	6	144	27 %	2.3e-01
2400	gi 545858131	keratin, type I cytoskeletal 10 isoform X2	58318.6	4.9	16	190	21 %	5.8e-01
2482	gi 117660856	MYLPF	19066.3	4.8	16	600	86 %	5.8e-05
2522	gi 117660874	MLC1f	21018.6	4.9	8	190	40 %	5.8e-01
1154	gi 545845559	PREDICTED: fibrinogen beta chain isoform X2	50399.8	7.94	18	gi 545845559	43 %	4.6e-03
2225	gi 262263205	triosephosphate isomerase 1	26878.9	6.54	14	463	74 %	2.9e-04
2203	gi 262263205	Triosephosphate somerese 1	26878.9	6.54	17	774	77 %	2.3e-07
1326	gi 311247991	PREDICTED: pyruvate dehydrogenase protein Xcomponent-like isoform	54036.4	8.3	12	317	28 %	1.2e-02
1416	gi 545801458	PREDICTED: actin, alpha cardiac muscle 1 isoform X1	42334	5.2	10	313	31 %	2.9e-02
2250	gi 262263205	Triosephosphate isomerase 1, TPI1	26878.9	6.5	17	795	82 %	1.8e-07
2584	gi 314907119	A-FABP adipocyte fatty acid-binding protein	14780.6	6.2	8	371	56 %	4.6e-03
1433	gi 545833443	PREDICTED: alpha-enolase isoform X1	38172.7	8.93	12	377	41 %	1.2e-03
2221	gi 55926209	HSPB1 heat shock protein beta-1	22984.7	6.23	8	236	31 %	1.5e-01

Table 2 Protein differentially expressed for Shaziling pig breeds and Yorkshire were identified by 2-DE and MALDI-TOF-MS/MS (Continued)


1805	gi 545859898	PREDICTED: beta-enolase isoform X1	48154.9	8.52	17	414	40 %	2.3e-03
2550	gi 545805333	PREDICTED: 14 kDa phosphohistidine phosphatase isoform X2	14036.8	5.91	14	454	74 %	2.3e-04
2217	gi 262263205	triosephosphate isomerase 1	26878.9	6.54	16	701	82 %	4.6e-06
1711	gi 46389777	troponin T fast skeletal muscle type	30720.1	8.68	14	423	41 %	2.9e-03
1137	gi 545845559	PREDICTED: fibrinogen beta chain isoform X2	50399.8	7.94	15	289	34 %	7.3e-02
2215	gi 262263205	triosephosphate isomerase 1	26878.9	6.54	18	684	76 %	2.3e-06
2223	gi 55926209	HSPB1 heat shock protein beta-1	22984.7	6.23	10	371	42 %	4.6e-03
1833	gi 545832797	PREDICTED: troponin T, slow skeletal muscle isoformX1	32449.6	5.54	9	418	32 %	9.2e-03
1374	gi 268607671	actin, alpha skeletal muscle	42366	5.23	11	348	35 %	9.2e-03


^aNumber of times we would expect to obtain an equal or higher score by chance

membrane, cell, and cell part were prominently represented. For molecular functions, “catalytic activity” and “binding” represented the majority of the category.

To further evaluate the function of the unigenes, KOG classification of all the unigenes was performed, and

16,171 unigenes could be assigned to 25 categories (Fig. 5). Among the 25 KOG categories, the highest proportion of unigenes matched to “signal transduction mechanisms” (13.8 %), followed by cluster for “general function prediction only” (12.6 %), “post-translational

**Fig. 2** GO annotation of different proteinic spots (**a**: molecular function distribution, **b**: Biological process distribution)



modification, protein turnover, chaperones” (9.8 %), and “carbohydrate transport and metabolism” (5.8 %). The KEGG pathway database is a powerful tool for the analysis of gene function in terms of gene networks [52]. To identify the biological pathways of transcriptome from the LM, all unigenes were submitted to KEGG pathway analysis. A total of 16,755 unigenes were assigned to 278 KEGG pathways (Additional file 1: Table S1), which fell into five main categories, including metabolism, environmental information processing, genetic information processing, cellular processes, and organism systems. Among the five main categories, the metabolism pathway represented the largest number of the unigenes (1013). These metabolic pathways were mainly carbohydrate metabolism, energy metabolism, amino acid metabolism, and lipid metabolism, which provided a valuable resource to investigate muscle growth and lipid metabolism processes.

Identification and analysis of DEGs

To identify the DEGs between the Shaziling and Yorkshire pigs, the relative expression of the genes was estimated using the RPKM value, which is a common method for estimating transcript levels [53]. According to the RNA-seq analysis, DEGs were selected using the criteria of FC ≥2 and false discovery rate ≤0.05; the FC distribution of DEGs is shown in Fig. 6. A total of 488 unigenes were identified

as DEGs between the two breeds, of which 297 were up-regulated in the Shaziling pig and the 192 genes were up-regulated in the Yorkshire pig. To better explore the functions of the DEGs, GO functional analysis was carried out. The results indicated that 488 DEGs were significantly enriched in 208 GO categories (Additional file 2: Table S2). Notably, some of the categories are involved in the metabolic process and the regulation of skeletal muscle development.

qPCR analysis

To evaluate the proteomics and transcriptomics results, the expressions of eight genes were determined in the two pig breeds by qPCR on the basis of their different influence on the formation of meat flavor and skeletal muscle development. Four of the genes encoded proteins involved in glycolysis and fat deposition (*ENO1*, *ENO3*, *ATP5B*, and *TPI1*), three encoded proteins that correlated with synthesis of muscle fiber proteins (*MYLPE*, *ACTA1*, and *ACTC1*) and the *HSPB1* gene, whose encoded protein plays a crucial role in maintaining cellular homeostasis [54] and protects against stress [23]. The fold-changes in expression of the eight genes were compared with the transcriptomics and proteomics analysis results. As shown in Fig. 7, the qPCR results revealed that five genes were in agreement with the RNA-Seq results and six genes

Table 3 Statistical summary of the *longissimus dorsi* muscle transcriptome for assembling

Statistics	Counts	Average length (bp)	N50 (bp)	Longest length (bp)	Total length (bp)
Contigs	86,759	672	939	49,881	58,319,316
Scaffold	81,915	713	1028	66,767	58,377,616
Unigenes	79,320	733	1112	66,767	58,152,234

Table 4 Functional annotations using transcript BLAST analyses


Public database	Hit unigenes number	Percentage (hit/total) %
Annotated in nr	23,055	29.07
Annotated in UniProt	25,784	32.50
Annotated in GO	23,702	29.8
Annotated in KOG	16,171	20.3
Annotated in KEGG	16,755	21.1



showed a similar trend to proteomics results. In the present research, we also observed different expression trends between the proteomics and transcriptome analyses.


Correlation analysis of mRNA and protein expression

RNA-seq analyses identified 488 DEGs (Additional file 3: Table S3), of which 297 were upregulated in the Shaziling pig and 192 in the Yorkshire pig. Proteomics revealed identified 38 differentially abundant proteins, of which 27 protein spots were upregulated in the Shaziling pig and 11 in the Yorkshire pig. Similar to previous reports, the transcriptomics and proteomics data were divergent. In the present study, *ENO1* and *ACTC1* were overexpressed in Shaziling pigs, and *ATP5B* was overexpressed in Yorkshire pigs: only for these three genes were the proteomics results consistent with the transcriptomic results. In 2009, Timperio and colleagues performed a comparative analysis of proteomics and transcriptomics from the livers of Chianina and Holstein Friesian cattle. The results indicated that only three of 39 differentially abundant proteins were validated by microarray analyses [55]. Other research also confirmed that proteomics and transcriptomics data seldom overlap [56]. These differences were probably caused by alternative splicing, differential regulation of translation, and annotation errors of databases [57]. Another aspect concerning little overlap between transcriptomics and proteomics data is biological factors [58]. Although proteomics and transcriptomics data have almost no overlap, interaction pathway analyses could indicate shared biological significance [33]. Taking this into

consideration, the differentially abundant proteins and DEGs that converged in the same metabolic pathways, especially regulation of skeletal muscle development, were meaningful. Some of the proteins and upregulated gene transcripts in Shaziling pigs were found to be involved in the same metabolic pathways, particularly the glycolytic pathway (*ENO1*, *TPI1*, and *HSPB1*). *ENO1*, a glycolytic enzyme, is positively correlated with meat tenderness [59]. *TPI* is also a glycolytic enzyme, and has been shown to correlate with meat tenderness in porcine muscles [60]. Notably, although the *TPI1* result of proteomics and transcriptomics data did not match, pathway analyses of either DEG transcripts or proteins for the Shaziling samples were involved in a metabolic network. HSP proteins are related to protein folding and the oxidative stress response. In our research, *HSPB1* was overexpressed in Shaziling pigs, and might be positively correlated with meat quality, which agreed with previous studies [61, 62]. Shaziling pigs have excellent meat quality like other Chinese indigenous pig breeds. The IMF content in Shaziling pigs is 3.5 %, in Jinhua pigs it is 3.38 %, and in Lantang pigs it is 2.46 %. By contrast, Yorkshire pigs and Landrace pigs have IMFs of 1.79 % [7, 63]. Increased IMF content can improve meat quality significantly, especially in terms of tenderness [64]. According to our results, genes related to tenderness have a higher level of expression in Shaziling pigs than in Yorkshire pigs, for example *TPI1*, *HSPB1*, and *ENO1*. Further analysis of the DEGs identified a number of novel genes and pathways (Additional file 1: Tables S1, Additional file 2: Table S2







and Additional file 3: Table S3), which have not been reported to affect meat quality previously. Further characterization of these novel genes might reveal the regulatory mechanism underlying meat quality.

Conclusion

The object of this study was to investigate differences in the growth and development of skeletal muscle between Shaziling and Yorkshire pigs. The combined use of proteomic and transcriptomic analyses was effective in detecting DEGs and proteins. As a result, 38 differentially abundant proteins and 488 DEGs were identified

by mass spectrometry and RNA-seq analysis, respectively. Some of the proteins and unigenes are associated with lipid metabolism or glycolytic metabolism, according to previously published results. Based on the putative results of GO term enrichment and KEGG pathway analyses, we determine that many of the differential abundant proteins and DEGs are related to lipid mobilization, energy metabolism, the cytoskeleton, and signal transduction. Our study provided valuable information that could contribute to a deeper understanding of the molecular mechanisms regulating the development and formation of skeletal muscle.

Additional files

Additional file 1: Table S1. KEGG Pathway information of unigenes. (XLS 705 kb)

Additional file 2: Table S2. GO enrichment of DEG. (XLS 77 kb)

Additional file 3: Table S3. Summary of the differently expressed genes. (XLS 58 kb)

Abbreviations

2-DE, two-dimensional fluorescence difference gel electrophoresis; DEGs, differentially expressed genes; GO, gene ontology; IMF, intramuscular fat; KEGG, Kyoto Encyclopedia of Genes and Genomes; LM, *longissimus dorsi* muscle; MS, mass spectrometry; NGS, next-generation sequencing; qPCR, quantitative real-time PCR

Acknowledgements

We would like to thank LY for exceptional assistance in the qPCR and DX in collection of samples of pig tissue.

Funding

This study was supported by the National High Technology Research and Development Program of China (2011AA100304), the Provincial Natural Science Foundation of Hunan (13JJ1021) and Science and technology project of hunan province (2014NK4135).

Availability of data and materials

The sequencing data from this study has been archived at the NCBI Sequence Read Archive under GEO database [GSE:70673].

Authors' contributions

HY performed analysis of the transcriptomics and proteomics and drafted the manuscript. XX and JJ participated in the biological experiments and data analysis. HM managed the whole project. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The experiment and all its procedures were reviewed and approved by the Institutional Animal Care and Use Committee (IACUC) at Hunan Agricultural University.

Author details

¹College of Animal Science and Technology, Hunan Agricultural University, Changsha 410128, People's Republic of China. ²College of Life Science and Resource Environment, Yichun University, Yichun 336000, People's Republic of China.

Received: 2 November 2015 Accepted: 2 June 2016

Published online: 13 June 2016

References

- Rosenvold K, Andersen HJ. Factors of significance, for pork quality – a review. *Meat Sci.* 2003;64:219–37.
- Megens HJ, Crooijmans RP, San Cristobal M, Hui X, Li N, Groenen MA. Biodiversity of pig breeds from China and Europe estimated from pooled DNA samples: differences in microsatellite variation between two areas of domestication. *Gen Sel Evol.* 2008;40(1):103–28.
- Scherf BD. World watch list for animal diversity. 3rd ed. Rome: FAO; 2000.
- Lefaucheur L. A second look into fibre typing – relation to meat quality. *Meat Sci.* 2010;84(2):257–70.
- Xu YJ, Jin ML, Wang LJ, Zhang AD, Zuo B, Xu DQ, et al. Differential proteome analysis of porcine skeletal muscles between Meishan and Large White. *J Anim Sci.* 2009;87(8):2519–27.
- Zhao SM, Ren LJ, Chen L, Zhang X, Cheng ML, Li WZ, et al. Differential expression of lipid metabolism related genes in porcine muscle tissue leading to different intramuscular fat deposition. *Lipids.* 2009;44(11):1029–37.
- Wu T, Zhang Z, Yuan Z, Lo LJ, Chen J, Wang Y, et al. Distinctive genes determine different intramuscular fat and muscle fiber ratios of the longissimus dorsi muscles in Jinhua and landrace pigs. *PLoS One.* 2013;8(1):e53181.
- Lu P, Li DF, Yin JD, Zhang LY, Wang ZY. Flavour differences of cooked longissimus muscle from Chinese indigenous pig breeds and hybrid pig breed (Duroc × Landrace × Large White). *Food Chem.* 2008;107(4):1529–37.
- Liu X, Xiong X, Yang J, Zhou L, Yang B, Ai H, et al. Genome-wide association analyses for meat quality traits in Chinese Erhualian pigs and a Western Duroc (Landrace × Yorkshire) commercial population. *Genet Sel Evol.* 2015; 47(1):44.
- Guo J, Shan T, Wu T, Zhu LN, Ren Y, An S, et al. Comparisons of different muscle metabolic enzymes and muscle fiber types in Jinhua and Landrace pigs. *J Anim Sci.* 2011;89(1):185–91.
- Kim NK, Joh JH, Park HR, Kim OH, Park BY, Lee CS. Differential expression profiling of the proteomes and their mRNAs in porcine white and red skeletal muscles. *Proteomics.* 2004;4(11):3422–8.
- Guo Y, Li J, Shang J, Jin Y. Development of muscle-related genes and their effects on meat quality. *Energy Procedia.* 2012;16:229–33.
- Picard B, Lefaucheur L, Berri C, Duclos MJ. Muscle fibre ontogenesis in farm animal species. *Reprod Nutr Dev.* 2002;42(5):415–31.
- Zhao X, Mo D, Li A, Gong W, Xiao S, Zhang Y, et al. Comparative analyses by sequencing of transcriptomes during skeletal muscle development between pig breeds differing in muscle growth rate and fatness. *PLoS One.* 2011;6(5):e19774.
- Karlsson AH, Klont RE, Fernandez X. Skeletal muscle fibres as factors for pork quality. *Livest Prod Sci.* 1999;60(2):255–69.
- Wimmers K, Murani E, Ngu NT, Schellander K, Ponsuksili S. Structural and functional genomics to elucidate the genetic background of microstructural and biophysical muscle properties in the pig. *J Anim Breed Genet.* 2007; 124(1):27–34.
- Quiroz-Rothe E, Rivero JL. Coordinated expression of myosin heavy chains, metabolic enzymes, and morphological features of porcine skeletal muscle fiber types. *Micros Res Techniq.* 2004;65(1–2):43–61.
- da Costa N, McGillivray C, Chang KC. Postnatal myosin heavy chain isoforms in prenatal porcine skeletal muscles: insights into temporal regulation. *Anat Rec Part A.* 2003;273(2):731–40.
- Lefaucheur L, Ecolan P, Plantard L, Gueguen N. New insights into muscle fiber types in the pig. *J Histochem Cytochem.* 2002;50(5):719–30.
- Liu J, Damon M, Guitton N, Guisile I, Ecolan P, Vincent A, et al. Differentially-expressed genes in pig Longissimus muscles with contrasting levels of fat, as identified by combined transcriptomics, reverse transcription PCR, and proteomic analyses. *J Agri Food Chem.* 2009;57(9):3808–17.
- Xu YJ, Qian H, Feng XT, Xiong YZ, Lei MG, Ren ZQ, et al. Differential proteome and transcriptome analysis of porcine skeletal muscle during debelopment. *J Proteomics.* 2012;75(7):2093–108.
- Li A, Mo D, Zhao X, Jiang W, Cong P, He Z. Comparison of the longissimus muscle proteome between obese and lean pigs at 180 days. *Mamm Genome.* 2013;24(1–2):72–9.
- Görg A, Weiss W, Dunn MJ. Current two dimensional electrophoresis technology for proteomics. *Proteomics.* 2004;4(12):3665–85.
- Timms JF, Cramer R. Difference gel electrophoresis. *Proteomics.* 2008; 8(23–24):4886–97.
- Lin CS, Hsu CW. Differentially transcribed genes in skeletal muscle of Duroc and Taoyuan pigs. *J Anim Sci.* 2005;83(9):2075–86.
- Hollung K, Grove H, Færgestad EM, Sidhu MS, Berg P. Comparison of muscle proteome profiles in pure breeds of Norwegian Landrace and Duroc at three different ages. *Meat Sci.* 2009;81(3):487–92.
- Kim NK, Lim JH, Song MJ, Kim QH, Park BY, Kim MJ, et al. Developmental proteomic profiling of porcine skeletal muscle during postnatal development. *Asian Austral J Anim.* 2007;20(10):1612–7.
- Marguerat S, Bähler J. RNA-seq: from technology to biology. *Cell Mol Life Sci.* 2010;67(4):569–79.
- Ramayo-Caldas Y, Mach N, Esteve-Codina A, Corominas J, Castello A, Ballester M, et al. Liver transcriptome profile in pigs with extreme phenotypes of intramuscular fatty acid composition. *BMC Genomics.* 2012; 13(1):547.
- Chen C, Ai H, Ren J, Li W, Li P, Qiao R, et al. A global view of porcine transcriptome in three tissues from a full-sib pair with extreme phenotypes

- in growth and fat deposition by paired-end RNA sequencing. *BMC Genomics*. 2011;12(1):448.
31. Tian Q, Stepaniants SB, Mao M, Weng L, Feetham MC, Doyle MJ. Integrated genomic and proteomic analyses of gene expression in mammalian cells. *Mol Cell Proteomics*. 2004;3(10):960–9.
 32. Murgiano L, D'Alessandro A, Egidi MG, Crisa A, Prosperini G, Timpero AM, et al. Proteomics and transcriptomics investigation on longissimus muscles in Large White and Casertana pig breeds. *J Proteome Res*. 2010;9(12):6450–66.
 33. Kim NK, Park HR, Lee HC, Yoon D, Son ES, Kim YS, et al. Comparative studies of skeletal muscle proteome and transcriptome profilings between pig breeds. *Mamm Genome*. 2010;21(5–6):307–19.
 34. García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, et al. Qualimap: evaluating next-generation sequencing alignment data. *Bioinformatics*. 2012;28(20):2678–9.
 35. Wang L, Feng Z, Wang X, Wang X, Zhang X. DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics*. 2010;26(1):136–8.
 36. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. *Nat Protoc*. 2008;3(6):1101–8.
 37. Orosz F, Oláh J, Ovádi J. Triosephosphate isomerase deficiency: new insights into an enigmatic disease. *Biochim Biophys Acta*. 2009;1792(12):1168–74.
 38. Laville E, Sayd T, Terlouw C, Chambon C, Damon M, Larzul C, et al. Comparison of sarcoplasmic proteomes between two groups of pig muscles selected for shear force of cooked meat. *J Agric Food Chem*. 2007; 55(14):5834–41.
 39. Farrar WW, Deal Jr WC. Purification and properties of pig liver and muscle enolases. *J Protein Chem*. 1995;14(6):487–97.
 40. Kwasiborski A, Sayd T, Chambon C, Sante'-Lhoutellier V, Rocha D, Terlouw C. Pig Longissimus lumborum proteome: part II: relationships between protein content and meat quality. *Meat Sci*. 2008;80(4):982–96.
 41. Klont RE, Brocks I, Eikelenboom G. Muscle fibre type and meat quality. *Meat Sci*. 1998;49s:219–29.
 42. Ryu YC, Kim BC. Comparison of histochemical characteristics in various pork groups categorized by postmortem metabolic rate and pork quality. *J Anim Sci*. 2006;84(4):894–901.
 43. Helfferich W, Jump DB, Anderson DB, Skjaerlund DM, Merkel RA, Bergen WG. Skeletal muscle alpha-actin synthesis is increased pretranslationally in pigs fed the phenethanolamine ractopamine. *Endocrinology*. 1990;126(6):3096–100.
 44. Guillemain N, Bonnet M, Jurie C, Picard B. Functional analysis of beef tenderness. *J Proteomics*. 2011;75(2):352–65.
 45. D'Alessandro A, Marrocco C, Zolla V, D'Andrea M, Zolla L. Meat quality of the longissimus lumborum muscle of Casertana and Large White pigs: metabolomics and proteomics intertwined. *J Proteomics*. 2011;75(2):610–27.
 46. Huang X, Madan A. CAP3: a DNA sequence assembly program. *Genome Res*. 1999;9(9):868–77.
 47. Wickramasinghe S, Cánovas A, Rincón G, Medrano JF. RNA-sequencing: a tool to explore new frontiers in animal genetics. *Livest Sci*. 2014;166:206–16.
 48. Qian X, Ba Y, Zhuang Q, Zhong G. RNA-Seq technology and its application in fish transcriptomics. *OMICS*. 2014;18(2):98–110.
 49. Gui D, Jia K, Xia J, Yang L, Chen J, Wu Y, et al. De novo assembly of the Indo-Pacific humpback dolphin leucocyte transcriptome to identify putative genes involved in the aquatic adaptation and immune response. *PLoS One*. 2013;8(8):e72417.
 50. Ruan R, Guo AH, Hao YJ, Zheng JS, De WD. Novo assembly and characterization of narrow-ridged finless porpoise renal transcriptome and identification of candidate genes involved in osmoregulation. *Int J Mol Sci*. 2015;16(1):2220–38.
 51. Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics*. 2005;21(18):3674–6.
 52. Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Res*. 2011;40(Database issue):D109–14.
 53. Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods*. 2008;5:621–8.
 54. Liu Y, Steinacker JM. Changes in skeletal muscle heat shock proteins: pathological significance. *Front Biosci*. 2001;6:D12–25.
 55. Timperio AM, D'Alessandro A, Pariset L, D'Amici GM, Valentini A, Zolla L. Comparative proteomics and transcriptomics analyses of livers from two different Bos taurus breeds: "Chianina and Holstein Friesian". *J Proteomics*. 2009;73(7):309–22.
 56. Nie L, Wu G, Culley DE, Schotten JCM, Zhang W. Integrative analysis of transcriptomic and proteomic data: challenges, solutions and applications. *Crit Rev Biotechnol*. 2007;27(2):63–75.
 57. Hornshøj H, Bendixen E, Conley LN, Andersen PK, Hedegaard J, Panitz F. Transcriptomic and proteomic profiling of two porcine tissues using high-throughput technologies. *BMC Genomics*. 2009;10(1):30.
 58. Samulin J, Berget I, Lien S, Sundvold H. Differential gene expression of fatty acid binding proteins during porcine adipogenesis. *Comp Biochem Physiol B: Biochem Mol Biol*. 2008;151(2):147–52.
 59. D'Alessandro A, Rinalducci S, Marrocco C, Zolla V, Napolitano F, Zolla L. Love me tender: an Omics window on the bovine meat tenderness network. *J Proteomics*. 2012;75(2):4360–80.
 60. Hwang IH, Park BY, Kim JH, Cho SH, Lee JM. Assessment of postmortem proteolysis by gel-based proteome analysis and its relationship to meat quality traits in pig longissimus. *Meat Sci*. 2005;69(1):79–91.
 61. Bouley J, Meunier B, Chambon C, De Smet S, Hocquette JF, Picard B. Proteomic analysis of bovine skeletal muscle hypertrophy. *Proteomics*. 2005; 2:490–500.
 62. Morzel M, Terlouw C, Chambon C, Micol D, Picard B. Muscle proteome and meat eating qualities of Longissimus thoracis of "Blonde d' Aquitaine" young bulls: a central role of HSP27 isoforms. *Meat Sci*. 2008;78:297–304.
 63. Yu K, Shu G, Yuan F, Zhu X, Gao P, Wang S, et al. Fatty acid and transcriptome profiling of longissimus dorsi muscles between pig breeds differing in meat quality. *Int J Biol Sci*. 2013;9(1):108–18.
 64. Van Laack R, Stevens S, Stalder K. The influence of ultimate pH and intramuscular fat content on pork tenderness and tenderization. *J Anim Sci*. 2001;79:392–7.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

