

RESEARCH NOTE

Open Access



Social media usage among health care providers

Zoya Surani¹, Rahim Hirani², Anita Elias³, Lauren Quisenberry⁴, Joseph Varon^{5,6}, Sara Surani⁷ and Salim Surani^{8,9*} 

Abstract

Objective: The objective of this study was to evaluate the use of social media among healthcare workers in an attempt to identify how it affects the quality of patient care.

Results: An anonymous survey of 35 questions was conducted in South Texas, on 366 healthcare workers. Of the 97% of people who reported owning electronic devices, 87.9% indicated that they used social media. These healthcare workers indicated that they spent approximately 1 h on social media every day. The healthcare workers below the age of 40 were more involved in social media compared to those above 40 ($p < 0.05$). The use of social media among physicians and nurses was noted to be identical (88% for each group), and both groups encouraged their patients to research their clinical conditions on social media ($p < 0.05$). A higher number of physicians reported awareness of a social media policy in their hospital compared to nurses ($p < 0.05$). However, a large proportion of healthcare workers (40%) were unaware of their workplace policy, which could potentially cause a privacy breach of confidential medical information. Further studies are required to evaluate specific effects of these findings on the quality of patient care.

Keywords: Social media, Healthcare workers, Physicians, Nurses, Depression, Policy

Introduction

Seventy percent of Americans use social media and connect with each other, currently, which can be compared to only 5% of Americans, which used social media for this purpose in 2005 [1]. Social media can be considered one of the most innovative and one of the most destructive market forces which has emerged. Health care is not immune to it either. Currently, more than 40% of the health care consumers utilize social media for their healthcare information needs. This being even more in the consumers who are 18–24 years age group, compared to the 45–54 year age group. 90% of the health care consumers in the 18–24 year age group utilizes and believe health care information presented in the social media [2].

Whether through advertising or interactions with other colleagues, hospitals and healthcare workers engage with

one another via social media in their work fields [3]. 41% of the health care consumers chose social media to decide on their health care providers. In addition 26% of the hospitals in USA is currently participating in the social media in some form [2]. Social media has provided to be a communication system for healthcare workers through which they can find patient health information and past medical history or they can review relevant information about illnesses [4]. In terms of benefits, social media has provided healthcare providers a way to overcome barriers in delivering healthcare to patients [5], has enhanced self-management skills among patients who can familiarize themselves with specific illnesses and also provides numerous opportunities for providers to conduct research in their relevant fields [6]. Some clinicians have suggested that social media is among the best ways to design, collect, and analyze data into scientific papers for academic journals [7].

However, potential harmful outcomes can result from social media and should be considered, especially as they

*Correspondence: srsurani@hotmail.com

⁹ University of North Texas, 1177 West Wheeler Ave, Aransas Pass, TX 78366, USA

Full list of author information is available at the end of the article

relate to patient confidentiality [8]. Many hospitals and practices have strict social media policies, but not all employees are aware of them.

Additionally, age is thought to be directly correlated with the time spent on social media [9]. One study concluded that medical students, who are younger, spend more time on social media as compared to the residents, which suggests this correlation [10]. Another study showed that the use of social media among physicians was 41% in 2010 and 90% in 2011, as compared to medical students who usage was more than 90% [3]. Thus taking into account the factor of age could help us narrow down the general age group to work with in order to reduce the harmful effects of using social media during working hours, as it could be the source of distraction. Moreover, only 31% of healthcare institutions have social media policy, per Institute of Health [2].

The purpose of our study was to understand healthcare workers knowledge regarding the social media policies in their institution; their utilization of social media and also the utilization of healthcare providers above and below the age of 40 years of age, as the data has shown the utilization of social media is significantly higher in populations below the age of 45 years.

Main text

Study design

This study was an anonymous exploratory research using sampling of convenience. It analyze the most commonly used social media, estimate daily time spent on social media between health care providers, analyze subject awareness regarding their hospital's social media policy, and to see whether healthcare workers encouraged their patients to use it as a tool to educate themselves regarding disease process. An Ethics and Institutional Review Board (IRB) of Corpus Christi Medical Center, Corpus Christi Texas USA approved the survey and protocols. Consent waiver was granted by the IRB. The survey was carried out in a local community hospital and specialty clinic of mid-size suburban town.

All health care providers in the coastal bend area of Texas can be included. The healthcare providers who do not desire to fill the complete questionnaire or unable to read the questionnaires were excluded.

Methods

A sample size of 360 was needed based on the population of the health care providers in the coastal bend region of Texas to be 6000. A total of 366/370 healthcare provides participated (Table 1). Since there was not a validated questionnaire available, and being a pilot study, the questionnaire, comprised of 35 questions was developed, after deliberating the questionnaire among the

Table 1 Demographics of participants

Category	Subcategory	N (%)
Gender	Male	108 (29.5)
	Female	257 (70.2)
	No response	1 (0.3)
Race	African American/Black/Caribbean	12 (3.3)
	Asian/Pacific Islander	40 (11.0)
	Caucasian	148 (40.8)
	Hispanic/Latino	156 (43.0)
	Native American	2 (0.6)
	Other	5 (1.4)
Age group	< 20	0 (0)
	20–29	92 (25.3)
	30–39	123 (33.8)
	40–49	64 (17.6)
	50 +	85 (23.4)
Job	Physician	68 (19.0)
	RN	151 (42.3)
	LVN	10 (2.8)
	CMA	6 (1.7)
	PA	2 (0.6)
	Dietitian	1 (0.3)
	Speech pathologist	1 (0.3)
	Pharmacist	5 (1.4)
	Pharmacy technician	5 (1.4)
	Physical therapist	7 (2.0)
	Other	101 (28.3)

RN registered nurses, LVN license vocational nurses, CMA certified medical assistants, PA physician assistants

investigators. These questions were pretested among five physicians and five nurses. These were modified again based on the feedback from this pilot group. These questionnaires were anonymously distributed and collected in the healthcare setting. The responses from the paper questionnaire were manually entered and analyzed electronically, mainly using Microsoft Excel, Microsoft Corporation, California USA. p values were calculated using two-tail Student *t* test. p value of < 0.05 was considered significant.

Results

366 surveys were analyzed. 108 participants were male (29.5%), 257 were female (70.2%), and 1 preferred not to respond (0.3%) (Table 1). Hispanics (43.0%) and Caucasians (40.8%) dominated the sample size (Table 1). 97% of the sample size indicated that they own a type of electronic device. Of those 97%, 87.9% of indicated that they used some kind of social media on a regular.

Out of 366 participants, 217 were under the age of 40, and 151 participants were over 40 (Table 1). In the

and Accountability Act (HIPAA) (Centers for Disease Control and Prevention (CDC) [16]). The consequences in the breach of privacy can be severe and may lead to civil or criminal penalties [17].

Social media can also be a large distraction in the workplace. With all the perks of accessing medical records easily through different websites or applications, it is also highly tempting to use that resource for recreational purposes during work [18]. One study showed that when Facebook™ use was allowed in the workplace, productivity decreased by 1.47% [18]. In another study, institutions that allowed their participants to use Facebook™ every day for 15 min demonstrated a decrease in work efficiency by 1.5% [19].

Our study suggests that social media is an important tool for healthcare providers to help familiarize patients with their clinical conditions. This is especially beneficial for those who are unable to access healthcare information easily, including ethnic minorities and lower socioeconomic groups [20]. In one study, 53.5% of the patients who participated, indicated the use of social media for medical and health information [21]. However, social media may be deleterious for patient care.

To our knowledge, there are no previous studies quantifying the number of healthcare providers recommending the use of social media to their patients. We found that nearly half of participants (42.8%) encouraged their patients to read about disease processes on social media.

Our study suggested that a large number of healthcare workers are unaware of the social media policy at their workplace. This is worrisome because understanding institutional policies regarding social media is crucial in protecting confidential medical information and avoiding HIPAA compliance issues. Moreover, if healthcare providers are unaware of their social media policy, this can affect medical professionalism as a whole. It is important to note that though unintentional, this example is a violation of HIPAA, which restricts an individual from posting medical images to protect patient privacy [22]. Even after organizations such as the AMA provided these guidelines, not many healthcare providers are aware of or in compliance with their social media policies [23]. In one study, only 54% of healthcare institutions surveyed had a social networking policy [24]. Our result is consistent with these data (59.2%).

Some studies have shown that the use of social media decreases with age [25]. In one survey, 83% of the participants who were active users on social media were under the age of 30 years [25]. In our study, healthcare workers below the age of 40 used more social media and spent considerably longer amounts of time on it compared to healthcare workers above the age of 40 ($p < 0.05$). These differences are multifactorial. In our study, more

healthcare workers who were above the age of 40 indicated that social media takes away from their family time.

Conclusion

The use of social media in our cohort was in line with the general public. However, our study suggests that physicians contribute to medical forums online more than nurses. More than half of physicians and nurses encouraged their patients to learn about their disease processes online, however, only a small percentage believe that the information regarding the illness is reliable, which in contrast to what the consumers believe. Further studies are needed to determine the impact of social media on patterns of healthcare quality and disparities of care.

Limitations

Small sample size, encompassing single community hospital and clinic limits it results to be applied to the larger population. Moreover, cultural and geographical differences among population across the world may limit the generalization of this paper and hence similar studies in multiple geographical locations are required. Study design being sampling of convenience, also places some limitations. Not having a validated questionnaire also is a limitation of this study. We though suggest development of validated questionnaire and conducting a randomized trial to address this important issue which health care industry is facing.

Abbreviations

USA: United States of America; IRB: Institutional Review Board; CDC: Center for Disease Control and Prevention.

Authors' contributions

ZS, LQ, SS1 and SS2 were involved in design, data collection, data entry and preparation of manuscript. JV was involved in study design and preparation and review of manuscript. RH and AE were involved in preparation of manuscript. All authors read and approved the final manuscript.

Author details

¹ Veterans Memorial High School, Corpus Christi, TX, USA. ² School of Natural Science, Hampshire College, Amherst, MA, USA. ³ Sanamed Education, Miami, FL, USA. ⁴ Texas Tech University, El Paso, TX, USA. ⁵ The University of Texas Health Science Center at Houston, Houston, TX, USA. ⁶ The University of Texas, Medical Branch at Galveston, Houston, TX, USA. ⁷ Harvard University, Cambridge, MA, USA. ⁸ Texas A&M University, Corpus Christi, TX, USA. ⁹ University of North Texas, 1177 West Wheeler Ave, Aransas Pass, TX 78366, USA.

Acknowledgements

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The raw data can be obtained from the correspondence author.

Consent to publish

Not applicable.

Ethics approval and consent to participate

The Ethics & Institutional Review Board of Corpus Christi Medical Center, Corpus Christi, Texas USA approved this study. The same board granted waiver of consent.

Funding

None.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 30 June 2017 Accepted: 25 November 2017

Published online: 29 November 2017

References

- Center PR. Social media fact sheet 2017. <http://www.pewinternet.org/fact-sheet/social-media/>. Accessed 26 Sep 2017.
- 24 Outstanding statistics and figures on how social media has impacted the health care industry. Referral MD; 2017. <https://getreferrallmd.com/2013/09/healthcare-social-media-statistics/>. Accessed 26 Sep 2017.
- George DR, Rovniak LS, Kraschnewski JL. Dangers and opportunities for social media in medicine. *Clin Obstet Gynecol*. 2013;56(3):453–62.
- Chretien KC, Kind T. Social media and clinical care: ethical, professional, and social implications. *Circulation*. 2013;127(13):1413–21.
- Kaplan AMHM. Users of the world, unite! The challenges and opportunities of social media. *Bus Horiz*. 2010;53:59–68.
- Mattingly TJ II. Innovative patient care practices using social media. *J Am Pharm Assoc JAPhA*. 2015;55(3):288–93.
- Campbell L, Evans Y, Pumper M, Moreno MA. Social media use by physicians: a qualitative study of the new frontier of medicine. *BMC Med Inform Decis Mak*. 2016;16(1):1–11.
- Campbell L, Evans Y, Pumper M, Moreno MA. Social media use by physicians: a qualitative study of the new frontier of medicine. *BMC Med Inform Decis Mak*. 2016;16:91.
- Stelzner MA. 2011 Social media marketing industry report 2011. <http://www.socialmediaexaminer.com/socialmediamarketingreport2011.pdf>. Accessed 3 Jan 2017.
- Thompson LA, Dawson K, Ferdig R, Black EW, Boyer J, Coutts J, et al. The intersection of online social networking with medical professionalism. *J Gen Intern Med*. 2008;23(7):954–7.
- Xiang ZGU. Role of social media in online travel information search. *Tour Manag*. 2010;3(2):179–88.
- Fromm RE Jr, Varon J, Sternbach GL. Emergency medicine cyberspace. *J Emerg Med*. 1996;14(1):91–2.
- Brown J, Ryan C, Harris A. How doctors view and use social media: a national survey. *J Med Internet Res*. 2014;16(12):e267.
- Ash JS, Berg M, Coiera E. Some unintended consequences of information technology in health care: the nature of patient care information system-related errors. *J Am Med Inform Assoc JAMIA*. 2004;11(2):104–12.
- Courtney KSO, Kuo A. Enabling health and healthcare through ICT. Amsterdam: IOS Press; 2013.
- (CDC) CfDcAP. HIPAA privacy rule and public health guidance from CDC and the US department of Health and Human Services. CDC; 2003. <http://www.cdc.gov/privacyrule/Guidance/Content.htm>. Accessed 23 Dec 2016.
- Spector N, Kappel DM. Guidelines for using electronic and social media: the regulatory perspective. *Online J Issues Nurs*. 2012;17(3):1.
- Cain J. Social media in health care: the case for organizational policy and employee education. *Am J Health Syst Pharm AJHP*. 2011;68(11):1036–40.
- Moqbel M. The effect of the use of social networking sites in the workplace on job performance. Texas A&M University; 2012. <http://www.citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.306.5775&rep=1&type=pdf>. Accessed 15 Feb 2017.
- Moorhead SA, Hazlett DE, Harrison L, Carroll JK, Irwin A, Hoving C. A new dimension of health care: systematic review of the uses, benefits, and limitations of social media for health communication. *J Med Internet Res*. 2013;15(4):e85.
- Diaz JA, Griffith RA, Ng JJ, Reinert SE, Friedmann PD, Moulton AW. Patients' use of the Internet for medical information. *J Gen Intern Med*. 2002;17(3):180–5.
- Cao F, Huang HK, Zhou XQ. Medical image security in a HIPAA mandated PACS environment. *Comput Med Imaging Graph*. 2003;27(2–3):185–96.
- Shore R, Halsey J, Shah K, Crigger BJ, Douglas SP, Ethical AMACo, et al. Report of the ama council on ethical and judicial affairs: professionalism in the use of social media. *J Clin Ethics*. 2011;22(2):165–72.
- Backman C, Dolack S, Dunyak D, Lutz L, Tegen A, Warner D, et al. Social media and healthcare. *J AHIMA*. 2011;82(3):20–5 (quiz 6).
- Duggan M, Brenner J. The demographics of social media users-2012. Washington: Pew Research Center; 2013. <http://www.pewinternet.org/2013/02/14/the-demographics-of-social-media-users-2012/>. Accessed 2 Mar 2017.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

