

RESEARCH

Open Access



Apigetrin inhibits adipogenesis in 3T3-L1 cells by downregulating PPAR γ and CEBP- α

Fatma Hadrich*  and Sami Sayadi*

Abstract

Background: Apigetrin, a flavonoid found in many plant leaves and seeds, has been known to possess antimutagenic, anti-cancer, antioxidant and anti-inflammatory properties. Here, we are investigating the effect of the apigetrin on adipocytes differentiation in 3T3-L1 adipocytes, and elucidating the mechanism of its action.

Methods: Lipids accumulation was measured by Oil Red O staining and cell cycle was analyzed by flow cytometry. The antioxidant effect of apigetrin was evaluated against hydrogen peroxide. The expression of various genes, involved in adipogenesis and inflammation, was studied by real-time PCR.

Results: Our results showed that apigetrin treatment inhibited significantly lipid accumulation without effect on cell viability at 100 μ M, and it exerted the anti-adipogenic effect during the early stages of differentiation. Flow cytometry analysis showed that apigenin-7-O-glucoside (Ap7G) inhibited cell proliferation during mitotic clonal expansion and caused cell cycle delay. Quantitative PCR analysis revealed that the mRNA levels of C/EBP- α , PPAR- γ , SREBP-1c and FAS were suppressed after apigetrin treatment at 100 μ M. Moreover, the mRNA level of pro-inflammatory genes (TNF- α and IL-6) were suppressed after apigetrin treatment, at high concentration preadipocyte cells.

Conclusion: Taken together, these results indicated that apigenin-7-O-glucoside inhibits adipogenesis of 3T3-L1 preadipocytes at early stage of adipogenesis.

Keywords: Apigetrin, Adipogenesis, Cell cycle, ROS, TNF- α , IL-6

Background

The prevalence of obesity has rapidly increased in the past several decades and became a global health problem [1]. Obesity is often characterized by physical inactivity and consumption of a high-fat/cal diet and associated with lifestyle-related diseases, such as hyperlipidaemia, hypertension [2], arteriosclerosis, type 2 diabetes mellitus, cancer, respiratory complications, and osteoarthritis [3].

At the cellular level, obesity is defined by an increase in adipose tissue mass, which is the result of an enlargement in fat cells and/or an increase in their number [4]. Adipogenesis is the process of preadipocytes differentiation into adipocytes. Several transcription factors are regulated including the CCAAT/enhancer-binding protein (C/EBP) gene family and peroxisome proliferator activated receptor- γ (PPAR γ) [5]. These transcription factors are also involved in the sequential expression of

adipocyte specific proteins [6, 7]. Therefore, inhibition of adipogenesis is critical for achieving an anti-obesity effect and the search for agents that could control this process has been extensively undertaken [8].

Natural products derived from plants are receiving a lot of attention as treatment options and as chemopreventive agents. Flavonoids, are naturally occurring polyphenolic compounds abundant in food plants, have attracted significant public attention due to their various biological activities [9]. These products possess many biological and pharmacological activities owing to their potential anticancer, anti-inflammatory, antioxidant, and antimicrobial properties, as well as important nutritional supplements to the human diet [10, 11]. Their anti-obesity effect can act through various mechanisms by preventing weight gain and this may be an excellent alternative strategy for developing safe anti-obesity drugs. Among flavonoids compounds that have been tested for potential anti-obesity, we cited the quercetin, rutin, naringenin, luteolin and resveratrol [12, 13].

* Correspondence: fati_hadrich@yahoo.fr; sami.sayadi@cbs.mrt.tn
Environmental Bioprocesses Laboratory, AUF Regional Excellence Pole (AUF-PER-LBP), Sfax Biotechnology Center, P.O. Box 1177, 3038 Sfax, Tunisia



Apigenin is found abundantly in vegetables and fruits and it has a variety of physiological properties such as antioxidant, anti-inflammatory [14]. Added to its low toxicity, this flavonoid has been shown to have selective effects in inhibiting cell growth and inducing apoptosis in cancer cells [15]. Apigenin has been used as a dietary supplement and was reported to exert anti-obesity effect in 3T3-L1 cells through activation of AMPK [16].

The pure form of apigenin is unstable but in natural form in food, it is present mostly as glucoside conjugate, which is called apigenin. The conjugated form could be an important determinant for its bioavailability and absorption. Previous reports have investigated the pharmacological effects of the apigenin including anti-inflammatory, antioxidant and anti-diabetic effects. However, its anti-adipogenic effect remains unclear. Therefore, the objective of this study was to investigate the effect of the apigenin on mouse preadipocyte differentiation and to elucidate the mechanism underlying its effect, with a particular focus on the expression of molecules involved in adipogenesis.

Materials and methods

Reagents

3T3-L1 cells were provided from the Health Science Research Resources Bank (HSRRB, Osaka, Japan). Apigenin was purchased from Extrasynthese Company (Genay, France). Dulbecco's modified Eagle's medium (DMEM high-glucose), Dexamethasone, 3-iso-butyl-1-methylxanthine, and Insulin were purchased from Sigma-Aldrich (Missouri, USA). MTT was provided from Gibco, DCFH-DA, was purchased from Sigma Aldrich.

Primers against PPAR γ , C/EBP α , FAS, SREBP-1c, TNF- α and IL-6 were purchased from Santa cruz Biotechnology (Beverly, MA, USA).

Cell culture

3T3-L1 cells were cultured in DMEM medium containing 10% FBS at 37 °C and 5% CO₂. Cells were plated at a density of 2×10^5 cells. After reaching the confluence, adipocyte differentiation was initiated using the same medium containing 10 mg/L insulin, 0.5 mmol/L isobutylmethylxanthine, and 1 μ mol/L dexamethasone for 2 days. The medium was then replaced with DMEM, containing 5 mg/L insulin for more 2 days, and then changed to fresh medium every 2 days. Apigenin was diluted in DMSO, the final quantity of the solvent was 0.1% for control and treated cells for all experiment.

Oil red O staining and quantification

After differentiation, 3T3-L1 cells were washed twice with phosphate buffered saline (PBS, pH 7.4), fixed with lipid droplet assay fixative solution (Cayman kit) for 1 h, and then stained with Oil red O solution (in 60%

isopropanol) at room temperature for 10 min. After washing, cells were checked and pictures were taken using a microscope (BioZero BZ-8000; Keyence, Osaka, Japan). Moreover, the dye was extracted with dye extraction solution and the absorbance (OD. 420 nm) was measured by a Spectra Max microplate reader (Spectra Max 190; Molecular Devices Corporation, CA, USA).

MTT assay

3T3-L1 cells were harvested in 96-well plate. After reaching the confluence, the culture medium was replaced by 100 μ L containing Apigenin (0–200 μ mol/L) and the cells were incubated for further 48 h. The culture medium was removed and replaced by 100 μ L of fresh culture medium containing 10% of sterile filtered MTT (Sigma-Aldrich). After 6 h, the insoluble formazan crystals were dissolved in 100 μ L/well SDS and the absorbance was measured at 570 nm. The inhibition (%) was expressed as the percentage of viable cell compared to control.

Trypan blue assay

3T3-L1 cells were seeded for 2 days after confluence in the presence of apigenin. Then, cells viability was quantified by Trypan Blue assay. After washing twice with PBS, cells were trypsinized and immediately stained with 0.5% trypan blue dye (Trypan Blue, Sigma-Aldrich) for 3 min. Cells were observed under an optical microscope, and the viability was calculated as the percentage ratio of the number of unstained cells relative to the total cells counted.

Measurement of intracellular ROS level

Intracellular reactive oxygen species (ROS) level was measured using a fluorescent dye, DCFH-DA according to [17]. Differentiated cells (Day 7) were cultured in DMEM containing, 400 μ M H₂O₂ for 4 h and then with 30 μ M DCFH-DA for 30 min. Then, fluorescence intensity (excitation/emission 485/528 nm) was measured using a Varioscan HT plate reader. The values were calculated as a percentage (%) of control.

Flow cytometry analysis

Post-confluent 3T3-L1 cells were treated with differentiation media in the absence or presence of several doses of apigenin for 24 h. Both detached and adherent cells were collected by trypsinization and washed with phosphate buffered saline (PBS). Cells were fixed with 70% ethanol at 4 °C for overnight. After removing of ethanol, cells were stained with propidium iodide (Sigma-Aldrich) for 30 min in the obscurity. Fluorescent cells analysis was carried out by using Guava EasyCyte (Guava Technologies, Hayward, CA, USA).

Gene expression analysis

Total RNA was extracted from 3T3-L1 cells by acid-GTC-phenol method [18]. After DNase I (Takara Bio, Otsu, Shiga, Japan) treatment and RNA repurification, the cDNA was synthesized using M-MLV Reverse Transcriptase (Takara), and subjected for PCR. Quantitative PCR analysis was carried out using SYBR Premix Ex Taq (Takara). Each cDNA was amplified (95 °C for 5 s, 60 °C for 30 s, 72 °C for 30 s, for 40 cycles) using specific primers (Table 1).

Statistical analysis

The results of three experiments were pooled and expressed as mean $\pm$ standard deviation (SD). Data were subjected to a one-way analysis of variance (ANOVA) using Graph Pad Prism 6.0 followed by Tukey's multiple comparison test at $p < 0.05$.

Results

Apigetrin inhibits MDI-induced adipogenesis of 3T3-L1 preadipocytes

To investigate the anti-adipogenic effect of apigetrin, 3T3-L1 preadipocytes were induced to differentiate with MDI in the presence or absence of apigetrin, and cells were stained with oil red O solution. Our results showed that apigetrin dose-dependently inhibits adipogenesis of 3T3-L1 preadipocytes (Fig. 1a and b). Intracellular lipid accumulation was reduced after apigetrin treatment by 5%, 37% and 62% at 50, 100 and 200 μ M, respectively compared to untreated group (Fig. 1b). To check whether this anti-adipogenic effect of apigetrin is due to its cytotoxicity, the effect of apigetrin on cell viability was measured by MTT assay. Our results showed that apigetrin does not exhibit cytotoxicity up to 100 μ M (Fig. 1b) suggesting that anti-adipogenic of apigetrin is not from cytotoxicity, but through regulation of other molecular mechanism. Together, these results suggest that apigetrin has anti-adipogenic effect in 3T3-L1 preadipocytes.

Apigetrin inhibits early stage of differentiation

To investigate the mechanism of anti-adipogenic effect of apigetrin during early phase of differentiation, 3T3-L1

cells were treated in the presence of different concentrations of apigetrin over 0–2 days (early stage), 2–4 days (middle stage), 6–8 days (late stage).

As shown in Fig. 2a, apigetrin exhibited anti-adipogenic effects essentially in the early stage. During the middle and late stages, its effect was very low, with no significant difference seen between the control and the treated cells.

Effect of apigetrin on the clonal expansion and cell cycle progression of 3T3-L1 cells during the early stage of differentiation

As described above, Ap7G displayed its main effect during the early stage of differentiation. We thus anticipated that this compound would affect the preadipocyte proliferation step.

Trypan blue assay result showed that following 24 h and 48 h exposure, apigetrin at 100 μ M decreased MDI-induced clonal expansion and the cell number remained lower in the treated culture (Fig. 2b). Next, cell cycle profile was examined by FACS analysis. Our results showed that apigetrin treatment caused a significant delay in the progression of the cell cycle and increased G0/G1 and S population in a dose-dependent manner (Fig. 2c) without any effect in the detection of dividing cells (G2M).

qRT-PCR analysis

Several transcription factors, such as the C/EBP and PPAR families, are sequentially and cooperatively expressed during differentiation. In this study, we evaluated whether the decreases in intracellular lipid contents were associated with lower levels of PPAR- γ and C/EBP- α , expressed in the early stage of adipogenesis. As shown in Fig. 3, Ap7G (100 μ M) markedly suppressed MDI-induced up-regulation of PPAR- γ and C/EBP- α with no significant effect at 50 μ M (Fig. 3a). Expression of both adipogenic marker proteins was not detected after 2 days of MDI treatment, representing the early stage of adipogenesis. Similarly, this compound was able to decrease the mRNA level of SREBP-1c and FAS (Fig. 3b). Moreover,

Table 1 Primers for RT-PCR. PCR was performed using the primers indicated as below under optimal amplification condition (95 °C for 5 min; 22–35 cycles of 95 °C for 30 s, 58 °C for 30 s, 72 °C for 30 s; 72 °C for 7 min) for each gene

Name	Forward	Reverse
GAPDH	5'-TGGTGAAGGTCGGTGTGAACGG-3'	5'-TGCCGTTGAATTTGCCGTGAGT-3'
PPAR γ	5'-AAACTCTGGGAGATTCTCT-3'	5'-TGGCATCTCTGTGTCAAC-3'
C/EBP α	5'-GCCAACTGAGACTCTCT-3'	5'-TGGCATCTCTGTGTCAAC-3'
SREBP-1c	5'-GCTTAGCCTCTACACCAACTGGC-3'	5'-ACAGACTGGTACGGGCCACAAG-3'
FAS	5'-TGGAGCCTGTGTAGCCTTCGAG-3'	5'-ACAGCCTGGGGTCATCTTTGCC-3'
IL-6	5'-GGTGACAACACGCGCTTCCC-3'	5'-GCCACTCTCTGTGACTCCAGC-3'
TNF- α	5'-AAATGGGCTCCCTCTCATCAGTTC-3'	5'-TCTGCTGGTGGTTTGTACGAC-3'




Fig. 1 a Effect of apigenin on adipocyte differentiation. Adipogenesis was induced in the presence or absence of several dose of apigenin for 8 days. Then, intracellular lipid accumulation was stained by oil red O solution. The intracellular lipid accumulation was expressed as a percentage of control values. **b** Cytotoxicity of apigenin on 3T3-L1 preadipocytes. The cells were cultured until confluence and then, cells were treated with Ap7G at indicated concentrations for 24 h. **c** Cell viability was measured by MTT assay. The viability was expressed as a percentage of control values. Data were presented as means $\pm$ S.D. ($n = 3$). The asterisks (**) indicate a significant difference between control group and MDI-treated group ($p < 0.01$)

Ap7G treated 3T3-L1 cells decreased the level of the pro-inflammatory genes especially TNF- α and IL-6 (Fig. 3c).

Effect of Apigenin on ROS production

To investigate the capacity of the apigenin-7-O-glucoside to reduce H_2O_2 induced ROS production, we use the fluorescence probe DCFH-DA. Our results showed that the adipocytes cells exposed to H_2O_2 showed an increase in the intracellular level of ROS compared to the untreated cells used as a control (Fig. 4). However, treated cells with apigenin reduced significantly ($p < 0.05$) the ROS level of about 21% at 100 μ M, respectively.

Discussion

Adipocyte differentiation and fat deposition are correlated to the emergence and development of obesity. Hence, many researchers advise that adipose tissue should become a main drug target for obesity and lipid metabolism dysregulation. The prevention and treatment of obesity with drugs is usually associated with the appearance of adverse reactions and drug dependence. For this reason, recent studies have been focusing on natural substances without side effects for the prevention or therapy of obesity [19]. Flavonoids are abundant in fruits, seeds and vegetables, and they have protective effects against various diseases [20]. Flavonoids showed different biological activities depending on their chemical structures; for example, the number and location of

hydroxyl groups, their conjugation pattern and the position of sugar in their structure [21].

In this study, we revealed for the first time the effect of apigenin in 3T3-L1 preadipocyte model. Previously, Rauter et al., [22] have reported the anti-hyperglycemic of Ap7G on streptozotocin-Induced diabetic rats. Our results showed that apigenin suppressed lipid accumulation in a dose-dependent manner. In fact, Ap7G inhibited triglyceride accumulation by around 5% and 37% at 50 and 100 μ M, respectively without any toxicity at high concentration. Our results are in concomitance with the report of Chao et al., 2013 who have demonstrated that flavonoids bounded to sugar present lower toxicity in either in vitro and in vivo models [23]. Although data shortage concerning the anti-obesity effect of Ap7G, Li et al., 2005 have revealed the anti-adipogenic effect of genistin, analogue of apigenin, in bone marrow stem cells (MSC) at 10 μ M [24]. Moreover, many studies showed that flavonol glycoside, isolated from plants, exhibited a strong inhibition against lipids accumulation, in adipocytes and hepatocytes cells [25, 26].

MCE is an important step in the adipocyte differentiation process. Thus, blocking DNA replication by various means prevents differentiation.

Time-course analysis of the effect of apigenin showed that this compound was able to exhibit its strongest effects during the early stages of differentiation, which runs in parallel with clonal expansion. Interestingly,




Fig. 2 a Effect of Ap7G on MDI induced cell number increase and cell cycle progression (**b** and **c**). Differentiation of 3T3-L1 preadipocytes was initiated in the presence of Ap7G (0, 50, 100 μ mol/L). After 24 h and 48 h, the cells were trypsinized and counted. Final concentration of DMSO was 0.1%. Change of cell cycle was analyzed by flow cytometry (**b**) and plotted on graph (**c**). The flow cytometry was performed 3 independent times. Data were presented as means $\pm$ S.D. ($n = 3$). The asterisks (*) and (**) indicate a significant difference between control group and MDI-treated group ($p < 0.05$) and ($p < 0.01$), respectively

Ap7G decreased DMI-induced clonal expansion of 3T3-L1 cells and delayed the cell cycle progression in a dose-dependent manner. These data are consistent with several previous studies, which showed that arresting or delaying the cell cycle of 3T3-L1 cells by phenolic compounds during the first 2 days of differentiation, decreased cell number and inhibited the differentiation rate of adipocytes. In this context, Hsu et al., [27] have reported that genistein, resveratrol and kaempferol, inhibits adipocyte differentiation during mitotic clonal. Ono et al., [16] have demonstrated that apigenin was able to decrease the accumulation of intracellular lipids in 3T3-L1 cells and regulated the expression of adipogenic genes during the first days.

PPAR γ is involved in the regulation of fat cell genes adipocytes and is the key nuclear transcription factor

that regulate different cellular functions, such as adipocyte lipogenesis, fatty acid transport, and energy metabolism. Furthermore, C/EBP α was found to be a regulator of cell differentiation. It is expressed at the adipogenic initiation stage and synergistically triggers adipocyte-specific gene expression with PPAR- γ after the growth arrest stage [28]. Our findings showed that Ap7G inhibited the gene expression of PPAR γ and C/EBP α , which lead to a reduction in fat deposition. Previous studies have confirmed that, apigenin suppressed adipocyte differentiation by reducing PPAR γ , C/EBP α mRNA levels and activating AMPK-activated protein kinase [16, 29]. Similarly, Park et al., [30] have reported that luteolin-7-O-glucoside inhibited the expression of adipogenic gene and down-regulated the expression of proteins involved in adipocytes differentiation.





Fig. 3 **a** and **b** Effect of apigenin on gene expression of PPAR γ , CEBP- α , SREBP-1c and FAS. **c** Effect of Ap7G on TNF- α and IL-6 gene expression. 3T3-L1 cells were cultured 8 days after initiation of differentiation. Cells were treated with 0–100 $\mu\text{mol/L}$ of Ap7G for 8 days at 37 $^{\circ}\text{C}$ in a humidified 5% CO_2 incubator. The relative expression level of PPAR γ , CEBP- α , SREBP-1c, FAS, TNF- α and IL-6 was quantified by qRT-PCR. Final concentration of ethanol was 0.1%. Data were presented as means $\pm$ S.D. ($n = 3$). The asterisks (**) indicate a significant difference between control group and MDI-treated group ($p < 0.01$)

Previous reports have shown that FAS can be a potential drug target because its inhibition can reduce food intake and obesity in mice [31]. The SREBP-1c appears to be involved and early expressed in adipocyte differentiation. It is the main regulator of FAS expression in liver and adipose tissues [32]. As shown in our results, apigenin treatment inhibited significantly FAS mRNA expression at 100 μM . Our findings suggest that blocking SREBP-1c activation by Ap7G resulted in a significant reduction in FAS expression. These results are in accordance with the research of Morikawa et al., who have demonstrated that flavonoids glycoside inhibits the expression of the transcription factors involved in the adipocyte differentiation [25].

The obesity is usually associated with increased mitochondrial ROS production, causing the oxidized lipids, synthesis of faulty proteins, and mtDNA mutations, which are related to decreased metabolic activity, mitochondrial dysfunction, and cellular insulin sensitivity [33]. Moreover, it has been established that high ROS production leads to a dysregulation of adipokine secretion and contribute to the production of pro-inflammatory cytokines such as: tumor necrosis factor- α (TNF- α), interleukin-6 (IL-6), monocyte chemoattractant protein-1 (MCP-1) and the reduction on adiponectin secretion [34]. In the present work, we found that apigenin decreased the intracellular ROS level induced by the hydrogen



peroxide and decreased the mRNA gene levels of TNF- α and IL-6 at high concentration. Similarly, treated mature adipocytes with apigenin decreased TNF- α secretion (data not shown). Our findings are in accordance with those reported by Francisco et al., [35], who showed that flavonoids, such as quercetin, genistein and luteolin glycosides showed potential anti-inflammatory agents, due to their significant inhibitory effects on ROS and NO levels produced by LPS-induced inflammation in mouse macrophage cells. Similarly, Miguel et al., [36] have demonstrated that pretreatment of animals with 50 mg/kg of apigenin inhibited the LPS induced the production of TNF- α and IL-6. In the same context, Hamalainen et al., have reported that genistin, was less effective on the iNOS inhibition and NO production in murine J774 macrophage [37]. Previous studies have shown that the O-glycosylation of flavonoids decreased the antioxidant activity of isoflavonoids due to the presence of the sugar that may attenuate the absorption of the flavonoid aglycone and thus reduced its activity. However, no clear structural activity relationship depending on the positions or types of sugar substitution was found in the anti-inflammatory activity for flavonoids glycosides.

Conclusion

The present study reveals that apigenin-7-O-glucoside can serve as a potential candidate for the management of obesity. It exerted inhibitory effects on 3T3-L1 preadipocytes differentiation and caused cell delay. As shown in molecular analysis, apigenin treatment decreased the mRNA levels of PPAR γ , CEBP- α , SREBP-1c and FAS genes involved in the adipocyte differentiation. This compound was also able to reduce the pro-inflammatory gene expression. Further studies are being carried out to

validate its effects on other biological process related to adipocyte biology.

Abbreviations

C/EBP α : CCAAT enhancer binding protein alpha; DEX: Dexamethasone; DMEM: Dulbecco's modified Eagle's medium; FAS: Fatty acid synthase; IBMX: 3-isobutyl-1-methylxanthine; IL-6: Interleukine-6; PPAR γ : Peroxisome proliferator-activated receptor-gamma; SREBP-1c: Sterol regulatory element-binding proteins; TNF- α : Tumor necrosis factor

Acknowledgements

This study was supported by the Ministry higher education and Scientific Research, Tunisia.

Funding

The funders had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Availability of data and materials

All data generated or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

SS designed and directed the study. FH performed the experiments, data analysis and drafted the manuscript. Both authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare that there are no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 20 October 2017 Accepted: 9 April 2018

Published online: 25 April 2018

References

- Choi JH, Yun JW. Chrysin induces brown fat-like phenotype and enhances lipid metabolism 316 in 3T3-L1 adipocytes. *Nutr.* 2016;32:1002–10.
- Nica Bulló M, Casas-Agustench P, Amigó-Correig P, Aranceta J, Salas-Salvadó J. Inflammation, obesity and comorbidities: the role of diet. *Public Health Nutr.* 2017;10:1164–72.
- Kotsis V, Nilsson P, Grassi G, Mancia G, Redon J, Luft F, et al. New developments in the pathogenesis of obesity-induced hypertension. *J Hypertens.* 2015;33:1499–508.
- Chae SY, Seo SG, Yang H, Yu JG, Suk SJ, Jung ES, et al. Anti-adipogenic effect of erucin in early stage of adipogenesis by regulating Ras activity in 3T3-L1 preadipocytes. *J Funct Foods.* 2015;19:700–9.
- Watanabe M, Inukai K, Katagiri H, Awata T, Oka Y, Katayama S. Regulation of PPAR γ transcriptional activity in 3T3-L1 adipocytes. *Biochem Biophys Res Commun.* 2003;300:429–36.
- Tang QQ, Lane MD. Adipogenesis: from stem cell to adipocyte. *Annu Rev Biochem Annual Rev.* 2012;81:715–36.
- Gregoire FM, Smas CM, Sul HS. Understanding adipocyte differentiation. *Physiol Rev.* 1998;78:783–09.
- Wang S, Moustaid-Moussa N, Chen L, Mo H, Shastri A, Su R, et al. Novel insights of dietary polyphenols and obesity. *J Nutr Biochem.* 2014;25:1–18.
- Zhang H, Tsao R. Dietary polyphenols, oxidative stress and antioxidant and anti-inflammatory effects. *Curr Opin Food Sci.* 2016;8:33–42.
- Pandey K, Rizvi S. Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative Med Cell Longev.* 2009;2:270–8.
- Rodriguez-Mateos A, Vauzour D, Krueger CG, Shanmuganayagam D, Reed J, Calani L, et al. Bioavailability, bioactivity and impact on health of dietary flavonoids and related compounds: an update. *Arch Toxicol.* 2014;88:1803–53.

12. Kim J, Lee I, Seo J, Jung M, Kim Y, Yim N, et al. Vitexin, orientin and other flavonoids from *Spirodela polyrhiza* inhibit adipogenesis in 3T3-L1 cells. *Phyther. Res.* 2010;24:1543–8.
13. Guo X, Liu J, Cai S, Wang O, Ji B. Synergistic interactions of apigenin, naringin, quercetin and emodin on inhibition of 3T3-L1 preadipocyte differentiation and pancreas lipase activity. *Obes Res Clin Pract.* 2016; 10:327–39.
14. Funakoshi-Tago M, Nakamura K, Tago K, Mashino T, Kasahara T. Anti-inflammatory activity of structurally related flavonoids, Apigenin, Luteolin and Fisetin. *Int Immunopharmacol.* 2011;11(9):1150.
15. Clere N, Faure S, Carmen Martinez M, Andriantsitohaina R. Anticancer properties of flavonoids: roles in various stages of carcinogenesis. *Cardiovasc Hematol Agents Med Chem.* 2011;9:62–77.
16. Ono M, Fujimori K. Antiadipogenic effect of dietary Apigenin through activation of AMPK in 3T3-L1 cells. *J Agric Food Chem.* 2011;59:13346–52.
17. Hadrich F, Garcia M, Maalej A, Moldes M, Isoda H, Feve B, et al. Oleuropein activated AMPK and induced insulin sensitivity in C2C12 muscle cells. *Life Sci.* 2016;151:167–73.
18. Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem.* 1987; 162:156–9.
19. Mohamed GA, Ibrahim SRM, Elkhatay ES, El Dine RS. Natural anti-obesity agents. *Bull Fac Pharmacy, Cairo Univ.* 2014;52:269–84.
20. Nijveldt RJ, van Nood E, van Hoorn DE, Boelens PG, van Norren K, van Leeuwen PA. Flavonoids: a review of probable mechanisms of action and potential applications. *Am J Clin Nutr.* 2001;74:418–25.
21. Corcoran MP, McKay DL, Blumberg JB. Flavonoid basics: chemistry, sources, mechanisms of action, and safety. *J Nutr Gerontol Geriatr.* 2012;31:176–89.
22. Rauter AP, Martins A, Borges C, Mota-Filipe H, Pinto R, Sepodes B, et al. Antihyperglycaemic and protective effects of flavonoids on streptozotocin-induced diabetic rats. *Phyther Res.* 2010;24:S133–8.
23. Chao S-C, Huang S-C, Hu D-N, Lin H-Y. Subtoxic levels of Apigenin inhibit expression and secretion of VEGF by uveal melanoma cells via suppression of ERK1/2 and PI3K/Akt pathways. *Evid Based Complement Alternat Med.* 2013;2013:817674.
24. Li X, Zhang J, Sui S, Yang M. Effect of daidzin, genistin, and glycitin on osteogenic and adipogenic differentiation of bone marrow stromal cells and adipocytic transdifferentiation of osteoblasts. *Acta Pharmacol Sin.* 2005; 26:1081–6.
25. Morikawa T, Ninomiya K, Miyake S, Miki Y, Okamoto M, Yoshikawa M, et al. Flavonol glycosides with lipid accumulation inhibitory activity and simultaneous quantitative analysis of 15 polyphenols and caffeine in the flower buds of *Camellia sinensis* from different regions by LCMS. *Food Chem.* 2013;140:353–60.
26. Bansal P, Paul P, Mudgal J, Nayak PG, Pannakal ST, Priyadarsini KI, et al. Antidiabetic, antihyperlipidemic and antioxidant effects of the flavonoid rich fraction of *Pilea microphylla* (L.) in high fat diet/streptozotocin-induced diabetes in mice. *Exp Toxicol Pathol.* 2012;64:651–8.
27. Hsu C-L, Yen G-C. Effects of flavonoids and phenolic acids on the inhibition of Adipogenesis in 3T3-L1 adipocytes. *J Agric Food Chem.* 2007;55:8404–10.
28. Ji S, Doumit ME, Hill RA, Catzeflis F, Stehelin D, Yoshihara Y. Regulation of Adipogenesis and key Adipogenic gene expression by 1, 25-Dihydroxyvitamin D in 3T3-L1 cells. *PLoS One.* 2015;10:e0126142.
29. Zhang Y, Liu X, Han L, Gao X, Liu E, Wang T. Regulation of lipid and glucose homeostasis by mango tree leaf extract is mediated by AMPK and PI3K/AKT signaling pathways. *Food Chem.* 2013;141:2896–905.
30. Park H-S, Kim S-H, Kim YS, Ryu SY, Hwang J-T, Yang HJ, et al. Luteolin inhibits adipogenic differentiation by regulating PPAR γ activation. *Biofactors.* 2009;35:373–9.
31. Ronnett G, Kim E, Landree L, Tu Y. Fatty acid metabolism as a target for obesity treatment. *Physiol Behav.* 2005;85:25–35.
32. Rosen ED, Walkey CJ, Puigserver P, Spiegelman BM. Transcriptional regulation of adipogenesis. *Genes Dev.* 2000;14:1293–307.
33. Sivitz WI, Yorek MA. Mitochondrial dysfunction in diabetes: from molecular mechanisms to functional significance and therapeutic opportunities. *Antioxid Redox Signal.* 2010;12:537.
34. Shoelson SE, Herrero L, Naaz A. Obesity, inflammation, and insulin resistance. *Gastroenterology.* 2007;132:2169–80.
35. Francisco V, Figueirinha A, Costa G, Liberal J, Lopes MC, García-Rodríguez C, et al. Chemical characterization and anti-inflammatory activity of luteolin glycosides isolated from lemongrass. *J Funct Foods.* 2014;10:436–43.
36. Miguel FG, Cavaleiro AH, Spinola NF, Ribeiro DL, Barcelos GRM, Antunes LMG, et al. Validation of a RP-HPLC-DAD method for chamomile (*Matricaria recutita*) preparations and assessment of the marker, Apigenin-7-glucoside, safety and anti-inflammatory effect. Evidence-Based Complement Altern Med. 2015:1–9. <https://doi.org/10.1155/2015/828437>.
37. Hämäläinen M, Nieminen R, Vuorela P, Heinonen M, Moilanen E. Anti-inflammatory effects of flavonoids. Genistein, kaempferol, quercetin, and daidzein inhibit STAT-1 and NF-kappaB activations, whereas flavone, isorhamnetin, naringenin, and pelargonidin inhibit only NF-kappaB activation along with their inhibitory effect on iNOS expression and NO production in activated macrophages. *Mediators Inflamm.* 2007:45673. <https://doi.org/10.1155/2007/45673>.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

