

Parabolic double cosets in Coxeter groups

Sara C. Billey^{1†} and Matjaž Konvalinka^{2‡} and T. Kyle Petersen^{3§} and William Slofstra^{4¶} and Bridget E. Tenner^{3||}

¹Department of Mathematics, University of Washington, Seattle, WA 98195, USA

²Department of Mathematics, University of Ljubljana, Jadranska 21, Ljubljana, Slovenia

³Department of Mathematical Sciences, DePaul University, Chicago, IL 60614, USA

⁴Institute for Quantum Computing, Quantum-Nano Centre, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada

received 2015-11-16, revised 16th November 2015, accepted tbd.

Abstract. Parabolic subgroups W_I of Coxeter systems (W, S) and their ordinary and double cosets W/W_I and $W_I \backslash W/W_J$ appear in many contexts in combinatorics and Lie theory, including the geometry and topology of generalized flag varieties and the symmetry groups of regular polytopes. The set of ordinary cosets wW_I , for $I \subseteq S$, forms the Coxeter complex of W , and is well-studied. In this extended abstract, we look at a less studied object: the set of all double cosets $W_I w W_J$ for $I, J \subseteq S$. Each double coset can be presented by many different triples (I, w, J) . We describe what we call the lex-minimal presentation and prove that there exists a unique such choice for each double coset. Lex-minimal presentations can be enumerated via a finite automaton depending on the Coxeter graph for (W, S) . In particular, we present a formula for the number of parabolic double cosets with a fixed minimal element when W is the symmetric group S_n . In that case, parabolic subgroups are also known as Young subgroups. Our formula is almost always linear time computable in n , and the formula can be generalized to any Coxeter group.

Résumé. Sous-groupes paraboliques W_I de systèmes de Coxeter (W, S) et de leur classes ordinaires et doubles W/W_I et $W_I \backslash W/W_J$ apparaissent dans de nombreux contextes dans combinatoire et la théorie de Lie, dont la géométrie et la topologie de variétés de drapeaux généralisées et les groupes de symétrie de polyèdres réguliers. L'ensemble de classes ordinaires wW_I , pour $I \subseteq S$, forme le complexe de Coxeter de W , et est bien étudié. Dans ce résumé étendu, nous regardons un objet moins étudié: l'ensemble des tous les classes doubles $W_I w W_J$ pour $I, J \subseteq S$. Chaque classe double peut être présenté par de nombreux triples différents (I, w, J) . Nous décrivons ce que nous appelons la présentation lex-minimal et prouvons qu'il existe un tel choix unique pour chaque double classe. L'énumération des présentations lex-minimal peut être trouvé en utilisant un automate fini selon le graphe de Coxeter pour (W, S) . En particulier, nous présentons une formule pour le nombre de classes paraboliques doubles avec un élément minimal fixé dans le cas W est le groupe symétrique S_n . Dans ce cas, les sous-groupes paraboliques sont également connu sous le nom des sous-groupes de Young. Notre formule est presque toujours calculable dans un temps linéaire en n . La formule peut être généralisée à tout groupe de Coxeter.

[†]Email: billey@math.washington.edu. Partially supported by the National Science Foundation grant DMS-1101017.

[‡]matjaz.konvalinka@fmf.uni-lj.si. Supported by Research Program Z1-5434 and Research Project BI-US/14-15-026 of the Slovenian Research Agency.

[§]tpeter21@depaul.edu. Partially supported by a Simons Foundation collaboration grant.

[¶]weslofst@uwaterloo.ca.

^{||}bridget@math.depaul.edu. Partially supported by a Simons Foundation collaboration grant.

Keywords: Coxeter group, parabolic subgroup, double cosets, enumeration

1 Introduction

Let G be a group with subgroups H and K . The group G is partitioned by the collection of double cosets $H \backslash G / K = \{HgK \mid g \in G\}$. The double cosets are usually more complicated than the one-sided cosets. For example, different double cosets HgK and $Hg'K$ can have different sizes. In this work, we investigate the parabolic double cosets of a finitely generated Coxeter group. That is, given a Coxeter system (W, S) of finite rank $|S|$, we consider cosets $W_I w W_J$ where I and J are subsets of the generating set S , and $W_I = \langle s : s \in I \rangle$ denotes the standard parabolic subgroup of W generated by the subset I . Such double cosets are well-studied, e.g., they play a prominent role in the paper of Solomon that first defines the *descent algebra* of a Coxeter group [11]. For finite Coxeter groups, these double cosets are intervals in Bruhat order, and these intervals have a rank-symmetric generating function with respect to length [7]. Such intervals correspond to the cell decomposition of certain rationally smooth Richardson varieties.

If we fix I and J , then the double quotient $W_I \backslash W / W_J$ is also well-studied. For example, Stanley [12] shows the Bruhat order on such a double quotient is strongly Sperner (for finite W), and Stembridge [14] has characterized when the natural root coordinates corresponding to elements in the quotient give an order embedding of the Bruhat order (for any finitely generated W). The number of elements in the quotient is a product of characters [13, Ex 7.77a],

$$|W_I \backslash W / W_J| = \left\langle \text{ind}_{W_I}^W 1_{W_I}, \text{ind}_{W_J}^W 1_{W_J} \right\rangle, \quad (1)$$

where 1_{W_J} denotes the trivial character on W_J .

In this paper, we are interested in a basic problem about parabolic double cosets that appears to have been unexamined until now; namely, how many double cosets $W_I w W_J$ does W have as I and J range across subsets of S ? Part of the motivation for this question comes from the analogous problem for ordinary cosets, where the set of such wW_I as I ranges across subsets of S is equal to the set of cells of the Coxeter complex. When W is the symmetric group S_n , the number of such cells is the n th ordered Bell number, see [8, A000670]. One fact that makes the ordinary case substantially simpler than the double version is that each ordinary parabolic coset is of the form wW_I for a unique subset $I \subseteq S$. If we take w to be the minimal element in the coset, then the choice of w is also unique. While double cosets do have unique minimal elements, different sets I and J often give the same double cosets $W_I w W_J$. Thus our question cannot be answered by simply summing equation (1) over all I and J .

To make our problem more tractable, we restrict to the set of double cosets with a fixed minimal element w . Since W has finite rank this set is always finite. Our main contribution is a formula for the number c_w of double cosets with minimal element w when $W = S_n$. This formula is efficient in most cases. The key to the formula is a condition on pairs of sets (I, J) that guarantees that each double coset $W_I w W_J$ arises exactly once. In other words, we get a unique presentation $W_I w W_J$ for each double coset with minimal element w . This criterion holds for any Coxeter group, and thus in principle our formula for c_w can also be extended to any Coxeter group. Specific enumerative consequences for other Coxeter groups are deferred to the full version of this paper.

When $W = S_n$, we can use our formula for c_w to calculate

$$p_n = \sum_{w \in S_n} c_w,$$

the number of double cosets in W . Although this requires summing up $n!$ terms, the approach seems to be a significant improvement over what was previously known. The terms p_n for $1 \leq n \leq 10$ are

$$1, 3, 19, 167, 1791, 22715, 334031, 5597524, 105351108, 2200768698.$$

For example, in $S_2 = \langle s : s^2 = e \rangle$ the three parabolic double cosets are $\{e\}$, $\{s\}$, and $\{e, s\}$. The sequence (p_n) previously did not appear in the OEIS; we have added it as sequence number A260700. In contrast, summing over I and J in equation (1) gives the sequence

$$1, 5, 33, 281, 37277, \dots$$

This counts the number of “two-way contingency tables” ([8, A120733], [13, Ex. 7.77], [4], [5, Section 5]). It is also the number of cells in a two-sided analogue of the Coxeter complex recently studied by the third author [9].

In the rest of this abstract, we first describe our enumeration formula for the symmetric group in some detail (Theorem 1, Section 2), review some basic properties of double cosets (Section 3), and then explain how to choose a unique presentation for each double coset (Section 4). The proof of the enumeration formula is sketched in Section 5.

2 The marine model and the formula for c_w

Let $W = S_n$, a Coxeter group with generating set $S = \{s_1, \dots, s_{n-1}\}$, where s_i denotes the i th adjacent transposition. Write a permutation $w = w_1 \cdots w_n \in S_n$ in one-line notation. Call j a *right ascent* of w if $w_j < w_{j+1}$, and i a *left ascent* of w if the value i appears to the left of $i + 1$ in $w_1 \cdots w_n$ (equivalently, if i is a right ascent of w^{-1}). An index in $\{1, 2, \dots, n - 1\}$ that is not an ascent is a *descent*. Let $\text{Asc}(w)$ denote the right ascent set of w . By symmetry, $\text{Asc}(w^{-1})$ is the left ascent set. The right descent set of w is the complement of $\text{Asc}(w)$ in S .

In terms of simple generators, j is a right ascent (resp., i is a left ascent) if and only if $\ell(ws_j) > \ell(w)$ (resp., $\ell(s_i w) > \ell(w)$), where ℓ is the length function. The element w is the minimal element of a double coset C if and only if C has the form $W_I w W_J$, where I and J are subsets of the left and right ascent sets of w respectively.

In addition to distinguishing between left and right ascents, we want to highlight ascents that are “small.” A *small right ascent* of w is an index j such that $w_{j+1} = w_j + 1$. If j is a small right ascent, then $ws_j w^{-1} = s_i$, where $i = w_j$, and we say that i is a *small left ascent* of w . Any ascent that is not a small ascent is a *large ascent*. For example, $w = 245613$ has $\text{Asc}(w) = \{1, 2, 3, 5\}$ and $\text{Asc}(w^{-1}) = \{2, 4, 5\}$. Its small right ascents are 2 and 3, while its small left ascents are 4 and 5. Both 1 and 5 are large right ascents, and 2 is a large left ascent.

Our formula for c_w involves structures and terminology that we will refer to as the *marine model*. As part of this model, we make the following definitions.

1. A *raft* of w is an interval $[i, j] \subseteq [n - 1]$ such that $i, i + 1, \dots, j$ are all small right ascents, while $i - 1$ and $j + 1$ are not.

2. A *(right) float* of w is a large right ascent that is not adjacent to any rafts. In other words, an index h is a right float of w if $w(h) < w(h+1) \neq w(h)+1$, and both $w(h) \neq w(h-1)+1$ and $w(h+1)+1 \neq w(h+2)$.
3. A *(right) rope* of w is a large right ascent that is adjacent to exactly one raft.
4. A *(right) tether* of w is a large right ascent that is adjacent to two rafts.

The sets of (right) rafts, floats, ropes, and tethers of w will be denoted by $\text{Rafts}(w)$, $\text{Floats}(w)$, $\text{Ropes}(w)$, and $\text{Tethers}(w)$ respectively. An explanation for this “maritime” terminology is provided in Section 5. Observe that every right ascent of w is either a tether, a rope, a float, or part of a raft. Also, because rafts are in some sense maximal, distinct rafts cannot be adjacent. For example, if $w = \text{id} \in S_n$ then w has one raft $R = [1, n-1]$, and no floats, ropes, or tethers. The rafts of 123567849 are $[1, 2]$ and $[4, 6]$, the only right float is at 8, and the only right tether is at 3.

Theorem 1 *There is a finite family of sequences $b_m^{\mathcal{T}}$, such that for any permutation w , the total number of parabolic double cosets with minimal element w is equal to*

$$c_w = 2^{|\text{Floats}(w)| + |\text{Floats}(w^{-1})|} \sum_{\substack{S \subseteq \text{Tethers}(w^{-1}) \\ T \subseteq \text{Tethers}(w)}} \left(\prod_{R \in \text{Rafts}(w)} b_{|R|}^{\mathcal{T}(R, S, T)} \right).$$

The sequences $b_m^{\mathcal{T}}$ satisfy a linear recurrence, and thus can be easily computed in time linear in m . The definition of these sequences requires a fair bit of notation, and we defer it to Section 5. The set $\text{Tethers}(w)$ is typically small (on the order of $1/n$), leading to an efficient formula for c_w . Unfortunately, there are some permutations for which $|\text{Tethers}(w)| + |\text{Tethers}(w^{-1})|$ is quite large. For example, the permutation

$$w = (1, 2, 17, 18, 3, 4, 19, 20, \dots, 15, 16, 31, 32)$$

has 8 tethers, and its inverse has 14, leading to a sum with 2^{22} terms (see Figure 2). In this case, the value $c_w = 632371867544102$ can be determined on a computer within a few minutes.

3 Parabolic double cosets

As in the introduction, let (W, S) be a Coxeter system of finite rank $|S| < \infty$. The generating set S is the *set of simple reflections* of W . The simple reflections and their relations are encoded in an edge-labeled graph on S called the *Coxeter graph*. Two generators $s, s' \in S$ are *adjacent* if the corresponding vertices are adjacent in the Coxeter graph.

The set of *reflections* is the set of conjugates of S , denoted by $T = \{ws w^{-1} : s \in S, w \in W\}$. Every element $w \in W$ can be written as a product of elements of S , and the *length* $\ell(w)$ of w is the minimal number of simple reflections in such a product. The *Bruhat order* on W is defined by taking the transitive closure of the relations $w < wt$, where $t \in T$ and $\ell(w) < \ell(wt)$.

As in the case of symmetric group, we say that a *right ascent* of w is a simple reflection $s \in S$ such that $\ell(ws) > \ell(w)$. Similarly, if $\ell(sw) > \ell(w)$, then s is a *left ascent*. We denote the set of right ascents by $\text{Asc}(w)$. The set of left ascents is equal to $\text{Asc}(w^{-1})$. The descent set of w is the complement of $\text{Asc}(w)$.

in S , i.e., it records the simple reflections s such that $\ell(ws) < \ell(w)$. We refer the reader to [1, 6] for further background on Coxeter groups.

As in the introduction, the Coxeter group generated by $I \subseteq S$ is a *standard parabolic subgroup* of W , denoted by W_I . We will identify I with the induced subgraph on vertices I of the Coxeter graph for W . For example, this I is *connected* if the corresponding induced subgraph is connected. The left cosets in W/W_I each have a unique minimal length element, and thus W/W_I can be identified with the set W^I of all minimal length left W_I -coset representatives. An element $w \in W$ belongs to W^I if and only if I contains no right descents of w ; that is, if $I \subseteq \text{Asc}(w)$.

Every element $w \in W$ can be written uniquely as $w = w^I v$, where $w^I \in W^I$ and $v \in W_I$. This is the *parabolic decomposition* of w [6, Sect. 5.12]. The product $w = w^I v$ is a *reduced factorization*, meaning that $\ell(w) = \ell(w^I) + \ell(v)$. As a poset under Bruhat order, every coset wW_I is isomorphic to W_I . Consequently, if W_I is finite, then every coset wW_I is also finite, and in addition has a unique maximal element, implying that wW_I is a Bruhat interval. Analogous statements can be made for the right cosets $W_I w$. Write ${}^I W$ for the set of minimal length right coset representatives for $W_I \backslash W$.

A *parabolic double coset* is a subset $C \subseteq W$ of the form $C = W_I w W_J$ for some $w \in W$ and $I, J \subseteq S$. Parabolic double cosets inherit some of the nice properties of ordinary cosets mentioned previously, including the following:

Proposition 2 ([2, 7]) *Let (W, S) be a Coxeter system, and fix $I, J \subseteq S$.*

- (a) *Every parabolic double coset in $W_I \backslash W / W_J$ has a unique minimal element with respect to Bruhat order. This element is also the unique minimal length element.*
- (b) *An element $w \in W$ is the minimal length element of a double coset in $W_I \backslash W / W_J$ if and only if w belongs to both ${}^I W$ and W^J . Thus $W_I \backslash W / W_J$ can be identified with*

$${}^I W^J := {}^I W \cap W^J.$$

- (c) *The parabolic double cosets in $W_I \backslash W / W_J$ are finite if and only if W_I and W_J are both finite. In this case, each $C \in W_I \backslash W / W_J$ has a unique maximal length element which is also the unique maximal element with respect to Bruhat order. In particular, if C is finite then it is a Bruhat interval.*

Corollary 3 (Double Parabolic Decomposition) *Fix $I, J \subseteq S$ and $w \in {}^I W^J$. Set $H := I \cap (wJw^{-1})$. Then $xw \in W^J$ for $x \in W_I$ if and only if $x \in W_I^H$. Consequently, every element of $W_I w W_J$ can be written uniquely as uwv , where $u \in W_I^H$ and $v \in W_J$, and $\ell(uwv) = \ell(u) + \ell(w) + \ell(v)$.*

4 Canonical presentations of double cosets

Fix a parabolic double coset C . A *presentation* of C is a choice of I, J and w such that $C = W_I w W_J$. Presentations are not unique; for instance, $W = W_S e W_\emptyset = W_\emptyset e W_S = W_S w W_S$ for any $w \in W$. To solve the problem that presentations are not unique, we define and characterize three types of presentations: maximal, minimal, and lex-minimal. We show that every parabolic double coset has unique maximal and lex-minimal presentations. The lex-minimal presentations are those that we have found most suitable for enumeration.

Definition 4 Given a parabolic double coset C , set

$$M_L(C) := \bigcup_{C=W_I w W_J} I \quad \text{and} \quad M_R(C) := \bigcup_{C=W_I w W_J} J$$

where both unions are over pairs (I, J) such that $C = W_I w W_J$.

Proposition 5 Let C be a parabolic double coset.

- (a) The coset C has a presentation $C = W_{M_L(C)} w W_{M_R(C)}$, and this is the largest possible presentation in the sense that if $C = W_I w' W_J$ then $I \subseteq M_L(C)$ and $J \subseteq M_R(C)$.
- (b) The sets $M_L(C)$ and $M_R(C)$ can be determined by

$$\begin{aligned} M_L(C) &= \{s \in S : sx \in C \text{ for all } x \in C\} \text{ and} \\ M_R(C) &= \{s \in S : xs \in C \text{ for all } x \in C\}. \end{aligned}$$

The presentation $W_{M_L(C)} w W_{M_R(C)}$ is the *maximal presentation* for C . We now look at presentations that are as small as possible.

Definition 6 A presentation $C = W_I w W_J$ is minimal if

- (a) $w \in {}^I W^J$,
- (b) no connected component of I is contained in $w J w^{-1} \cap S$, and
- (c) no connected component of J is contained in $w^{-1} I w \cap S$.

If a connected component I_0 of I is contained in $w J w^{-1} \cap S$ for $w \in {}^I W^J$, then $W_I w W_J = W_{I \setminus I_0} w W_J$. In other words, every presentation can be reduced to a minimal presentation. In Proposition 9, we will show that our nomenclature is appropriate, in that minimal presentations have minimum size.

Lemma 7 Let $C = W_I w W_J$ be a minimal presentation of C . Then

$$\begin{aligned} M_L(C) &= I \cup \{s \in (w J w^{-1}) \cap S : s \text{ is not adjacent to } I\} \text{ and} \\ M_R(C) &= J \cup \{s \in (w^{-1} I w) \cap S : s \text{ is not adjacent to } J\}. \end{aligned}$$

That $M_L(C)$ contains the set on the right hand side is clear. The other inclusion takes only slightly more work. The equality for $M_R(C)$ is analogous.

Corollary 8 Suppose that $C = W_I w W_J$ is a minimal presentation of C . If T is any connected subset of $M_L(C)$, then either $T \subseteq I$, or T is disjoint and non-adjacent to I and $T \subseteq (w J w^{-1}) \cap S$.

Given subsets $X, Y, Z \subseteq S$, write $X = Y \dot{\sqcup} Z$ to mean that X is the disjoint union of Y and Z , and Y and Z are non-adjacent. (In other words, the vertex-induced subgraph of X is isomorphic to the disjoint union of the vertex-induced subgraphs of Y and Z .) The following proposition is, roughly speaking, obtained by repeated application of Corollary 8.

Proposition 9 *Let $w \in {}^I W^J$. Then a presentation $C = W_I w W_J$ is minimal if and only if $|I| + |J| \leq |I'| + |J'|$ for all other presentations $C = W_{I'} w W_{J'}$ of C . Furthermore, if $C = W_I w W_J$ and $C = W_{I'} w W_{J'}$ are both minimal presentations, then there is a sequence of connected components I_1, \dots, I_m of I and J_1, \dots, J_n of J such that*

$$I' = \left(I \overset{\curvearrowright}{\sqcup} w J_1 w^{-1} \overset{\curvearrowright}{\sqcup} \dots \overset{\curvearrowright}{\sqcup} w J_n w^{-1} \right) \setminus \bigcup_{i=1}^m I_i \text{ and}$$

$$J' = \left(J \overset{\curvearrowright}{\sqcup} w^{-1} I_1 w \overset{\curvearrowright}{\sqcup} \dots \overset{\curvearrowright}{\sqcup} w^{-1} I_m w \right) \setminus \bigcup_{j=1}^n J_j.$$

(Note that the order of operations in these two identities is significant, in that $w J_j w^{-1}$ is disjoint and non-adjacent to I_i for all i, j .)

We now define lex-minimal presentations and give our desired characterization as a corollary to Proposition 9.

Definition 10 *A presentation $C = W_I w W_J$ of a parabolic double coset C is lex-minimal if w is the minimal element of C , and $(|I|, |J|)$ is lexicographically minimal among all presentations of C .*

In other words, if $C = W_I w W_J$ is lex-minimal and $C = W_{I'} w' W_{J'}$ is another presentation of C , then $w \leq w'$, and either $|I| < |I'|$, or $|I| = |I'|$ and $|J| \leq |J'|$.

Theorem 11 *Let $w \in {}^I W^J$. Then $C = W_I w W_J$ is lex-minimal if and only if*

- (a) *no connected component of J is contained in $(w^{-1} I w) \cap S$, and*
- (b) *if a connected component I_0 of I is contained in $w S w^{-1}$, then some element of I_0 is adjacent to but not contained in $w J w^{-1}$.*

Furthermore, every parabolic double coset has a unique lex-minimal presentation.

When w is the identity, Theorem 11 implies that lex-minimal presentations are two-level staircase diagrams in the sense of [10]. Note that while [10] addresses the enumeration of staircase diagrams, two-level staircase diagrams are not considered.

5 The Marine Model for S_n

We now return to the case $W = S_n$, and fix a permutation $w = w_1 \cdots w_n \in S_n$. Given that sets I and J are contained in the left and right ascent set of w , i.e., $I \subseteq \text{Asc}(w)$ and $J \subseteq \text{Asc}(w^{-1})$, we want to determine when $W_I w W_J$ is lex-minimal.

The Coxeter graph G of S_n is a path with vertices labelled $\{1, 2, \dots, n-1\}$. To the permutation w , we associate a diagram called the *marine model*, as follows. First, take two isomorphic copies G^L and G^R of G , and let B_w be the graph obtained from G^L and G^R , along with edges connecting the small left ascents of w to the associated small right ascents of w : vertex i in G^L and vertex j in G^R are connected whenever $w s_j w^{-1} = s_i$. The marine model is the graph A_w obtained by deleting from B_w all edges of G^L and G^R that are not incident to any small ascents.

As in the introduction, it is useful to have some terminology to refer to the parts of A_w . We use the following terminology:

1. Plank – an edge in A_w connecting a small left ascent to a right small ascent.
2. Raft – a maximal connected component of adjacent planks, where two planks are *adjacent* if the induced subgraph of A_w containing both of them is connected. As shown below, rafts are drawn as a series of parallel line segments of the same length connected across the top and bottom, hence the name. As in the introduction, we can identify the raft by the corresponding interval $[i, j]$ of small right ascents of w .
3. Float – a large left or right ascent of w not adjacent to any rafts.
4. Rope – a large left or right ascent of w adjacent to exactly one raft.
5. Tether – a large left or right ascent of w connected to two rafts.

It is helpful to draw A_w as follows. Draw two rows of $n - 1$ dots representing the possible right (top row) and left (bottom row) ascents of w . Denote small ascents and large ascents by small and large dots respectively. For each raft $[i, j]$ of w , draw a line from dot k on the top to dot w_k on the bottom for $k = i, \dots, j$, and connect the dots i, \dots, j on the top and w_i, \dots, w_j on the bottom. For example, for $w = (1, 3, 4, 5, 7, 8, 2, 6, 14, 15, 16, 9, 10, 11, 12, 13)$, we have rafts $[2, 3]$, $[5, 5]$, $[9, 10]$, and $[12, 15]$. Thus we draw nine lines from upper dots to lower dots, three lines on the top (2 to 3, 9 to 10, and 12 to 15), and three lines on the bottom (3 to 4, 9 to 12, and 14 to 15). If i is a rope or tether, draw an edge or edges horizontally from the larger dot in position i to the adjacent small ascent or ascents. In our example, the tethers are the edges incident to 4 on the top and 8 on the bottom. The ropes are the edges incident to 1 and 8 on the top and 5 on the bottom. We also circle any dot corresponding to a float, so in our example we circle dot 7 on the top and dot 1 on the bottom. The isolated dots that are not circled are the descents. This example is shown in Figure 1.




Fig. 1: Rafts, tethers, floats and ropes of $w = (1, 3, 4, 5, 7, 8, 2, 6, 14, 15, 16, 9, 10, 11, 12, 13)$.

Theorem 11 can be rephrased for the symmetric group as follows.

Theorem 12 *Let $w \in S_n$ and let I and J be subsets of the left and right ascent sets of w respectively. Then $W_I w W_J$ is a lex-minimal presentation of a parabolic double coset of S_n if and only if*

- (a) *there is no interval $[a, b] \subseteq J$ of small right ascents of w such that $\{w_a, \dots, w_b\}$ is contained in I and neither $a - 1$ or $b + 1$ are in J ; and*
- (b) *there is no interval $[a, b] \subseteq I$ of small left ascents of w such that $\{w_a^{-1}, \dots, w_b^{-1}\}$ is either contained in J or disjoint and non-adjacent to J , and neither $a - 1$ or $b + 1$ are in I .*




Fig. 2: The marine diagram for the element $w = (1, 2, 17, 18, 3, 4, 19, 20, \dots, 15, 16, 31, 32)$ that appears in the introduction. From this figure, we can verify that w has 22 tethers on the top and the bottom combined.

The enumeration formula in Theorem 1 counts the number of lex-minimal presentations of double cosets with fixed $w \in S_n$ as the minimal element. This is equivalent to the number of ways of choosing subsets I and J such that the conditions in Theorem 12 are satisfied. If the marine model of w consists of a single raft, then the conditions on I and J can be checked using a finite state automaton, pictured in Figure 3, that scans from left to right. The complete description of this automaton is deferred to the full paper. As a result of this machinery, the number of choices for I and J is given by a recurrence. A raft with ropes attached can be handled similarly, using different combinations of starting and ending states in the automaton. To make this specific, let $(a_m^k)_{m \in \mathbb{N}}$, $k \in \{0, 1, 2, 2', 2'', 3, 4\}$, be the family of sequences defined by the recurrence

$$a_m = 6a_{m-1} - 13a_{m-2} + 16a_{m-3} - 11a_{m-4} + 4a_{m-5} \quad \text{for } m \geq 5,$$

with initial conditions a_0^k, \dots, a_4^k given by the following table.

$k \backslash m$	0	1	2	3	4
0	1	2	6	20	66
1	1	3	9	28	89
2	1	4	12	36	112
2'	1	3	11	37	119
2''	1	4	12	37	118
3	1	4	14	46	148
4	1	4	16	56	184

Remark The characteristic polynomial corresponding to the recurrence, $1 - 6t + 13t^2 - 16t^3 + 11t^4 - 4t^5$, factors as $(1 - t + t^2)(1 - 5t + 7t^2 - 4t^3)$, and the sequences $(a_m^k)_{m \in \mathbb{N}}$ for $k = 0, 1, 2, 3, 4$ actually satisfy a recurrence of order 3:

$$a_m = 5a_{m-1} - 7a_{m-2} + 4a_{m-3} \quad \text{for } m \geq 3.$$

However, a single recurrence for all sequences makes the definition more concise.

Given $i_1, i_2, i_3, i_4 \in \{0, 1\}$, let

$$k(i_1, i_2, i_3, i_4) = \begin{cases} i_1 + i_2 + i_3 + i_4 & : i_1 + i_2 + i_3 + i_4 \neq 2 \\ 2 & : i_1 + i_2 + i_3 + i_4 = 2 \text{ and } i_1 = i_2 \\ 2' & : i_1 + i_2 + i_3 + i_4 = 2 \text{ and } i_1 = i_3 \\ 2'' & : i_1 + i_2 + i_3 + i_4 = 2 \text{ and } i_1 = i_4 \end{cases}.$$

The tuple (i_1, i_2, i_3, i_4) represents the lower-left, upper-left, lower-right, and upper-right outer corners of a raft: $\begin{smallmatrix} i_2 & i_4 \\ i_1 & i_3 \end{smallmatrix}$. The indicators are 1 (selected) or 0 (not selected) depending on whether or not a rope or tether attached at that point is selected for I or J . By symmetry, all four configurations with exactly one indicator equal to 1 are equivalent, as are the four configurations with exactly one indicator equal to 0. There are three configurations up to symmetry with exactly two indicators equal to 1, and these correspond to the states $k = 2, 2'$, and $2''$. Then $a_m^{k(i_1, i_2, i_3, i_4)}$ is the number of choices of I and J for a raft of length m . The automaton we use to recognize pairs (I, J) has 8 states (see Figure 3), and thus the generating functions for the sequences a_m^k can be computed as a sum of terms $(-1)^{i+j} \frac{\det(I-tA)^{j_i}}{\det(I-tA)}$, $1 \leq i, j \leq 8$, where A is the 8×8 transition matrix of the automaton. For instance,

$$\sum_m a_m^0 t^m = \frac{1 - 3t + 3t^2}{1 - 5t + 7t^2 - 4t^3} = 1 + 2t + 6t^2 + 20t^3 + 66t^4 + \dots$$




Fig. 3: The finite automaton (loops are omitted).

For a general permutation w , we have to handle floats, rafts, ropes, and tethers. Floats can be chosen or rejected completely independently of the other elements of I and J , leading to the term

$$2^{| \text{Floats}(w) | + | \text{Floats}(w^{-1}) |}.$$

Ropes and tethers can also be chosen independently, provided that this choice is incorporated into the calculation for the attached rafts as above. Once these choices are fixed, the calculation can be made independently for each raft. Because ropes are attached to exactly one raft, we gain some efficiency by incorporating the choice of ropes into the calculation for each raft. However, tethers must be chosen in advance, because they appear in two rafts. Thus the formula for c_w in Theorem 1 involves a sum across choices S and T of tethers on the bottom and the top respectively, after which we take a product of sequences $b_{|R|}^{\mathcal{I}(R, S, T)}$ corresponding to the rafts.

To make this specific, we define a family of sequences $b_m^{\mathcal{I}}$ indexed by tuples $\mathcal{I} = (I_1, I_2, I_3, I_4)$, where I_i is a non-empty subset of $\{0, 1\}$ indicating attached ropes or tethers in the same position as used for the

sequences $a_n^{k(i_1, i_2, i_3, i_4)}$. Here $I_i = \{0\}$ means that no rope or tether can be selected, $\{1\}$ means that an attached tether must be selected, and $\{0, 1\}$ means that an attached rope is available for selection. With this convention,

$$b_m^{\mathcal{I}} = \sum a_m^{k(i_1, i_2, i_3, i_4)},$$

where the sum is across $(i_1, i_2, i_3, i_4) \in I_1 \times I_2 \times I_3 \times I_4$. For example,

$$b_m^{\{\{0\}, \{0, 1\}, \{1\}, \{0, 1\}\}} = a_m^{k(0, 0, 1, 0)} + a_m^{k(0, 0, 1, 1)} + a_m^{k(0, 1, 1, 0)} + a_m^{k(0, 1, 1, 1)} = a_m^1 + a_m^2 + a_m^{2''} + a_m^3.$$

Although the family $b_m^{\mathcal{I}}$ contains 81 sequences, the sequences can be precomputed easily in time linear in m . (By symmetry, it is also possible to reduce the number of sequences to 27).

To describe which sequence $b_m^{\mathcal{I}}$ to use for each raft, suppose that $R = [i, j] \in \text{Rafts}(w)$, $S \subseteq \text{Tethers}(w^{-1})$, and $T \subseteq \text{Tethers}(w)$. Then $\mathcal{I}(R, S, T)$ is the 4-tuple (I_1, I_2, I_3, I_4) , where:

$$\begin{aligned} I_1 &= \begin{cases} \{1\} & : w_i - 1 \in S \\ \{0\} & : w_i - 1 \in \text{Tethers}(w^{-1}) \setminus S \text{ or } w_i - 1 \notin \text{Asc}(w^{-1}) \\ \{0, 1\} & : \text{otherwise} \end{cases} \\ I_2 &= \begin{cases} \{1\} & : i - 1 \in T \\ \{0\} & : i - 1 \in \text{Tethers}(w) \setminus T \text{ or } i - 1 \notin \text{Asc}(w) \\ \{0, 1\} & : \text{otherwise} \end{cases} \\ I_3 &= \begin{cases} \{1\} & : w_j + 1 \in S \\ \{0\} & : w_j + 1 \in \text{Tethers}(w^{-1}) \setminus S \text{ or } w_j + 1 \notin \text{Asc}(w^{-1}) \\ \{0, 1\} & : \text{otherwise} \end{cases} \\ I_4 &= \begin{cases} \{1\} & : j + 1 \in T \\ \{0\} & : j + 1 \in \text{Tethers}(w) \setminus T \text{ or } j + 1 \notin \text{Asc}(w) \\ \{0, 1\} & : \text{otherwise} \end{cases} \end{aligned}$$

This completes the description of the formula in Theorem 1.

Example 13 Let us compute the number of parabolic double cosets whose minimal element is our running example $w = (1, 3, 4, 5, 7, 8, 2, 6, 14, 15, 16, 9, 10, 11, 12, 13)$. There are two floats, so the first term is 2^2 . We have tether 4 on top and tether 8 on the bottom, so $S \subseteq \{8\}$, $T \subseteq \{4\}$, and there are four terms in the sum. For $S = T = \emptyset$, the raft $[2, 3]$ is of size 2 and has two ropes on opposite sides (the tether on top right was not selected), and contributes $b_2^{\{\{0\}, \{0, 1\}, \{0, 1\}, \{0\}\}} = a_2^0 + 2a_2^1 + a_2^{2''} = 6 + 2 \cdot 9 + 12 = 36$. The raft $[5, 5]$ is of size 1 and has ropes attached, and contributes $b_1^{\{\{0\}, \{0\}, \{0\}, \{0\}\}} = a_1^0 = 2$. The raft $[9, 10]$ has size 2 and has no ropes attached, so it contributes $b_2^{\{\{0\}, \{0, 1\}, \{0\}, \{0\}\}} = a_2^0 + a_2^1 = 6 + 9 = 15$. Finally, the raft $[12, 15]$ has size 4 and no ropes and therefore contributes $b_4^{\{\{0\}, \{0\}, \{0\}, \{0\}\}} = a_4^0 = 66$. The total contribution of $S = T = \emptyset$ is therefore $36 \cdot 2 \cdot 15 \cdot 66 = 71280$. For $S = \emptyset$, $T = \{4\}$, the raft $[2, 3]$ has two ropes and one attached tether, so the contribution is $b_2^{\{\{0\}, \{0, 1\}, \{0, 1\}, \{1\}\}} = a_2^1 + a_2^2 + a_2^{2'} + a_2^3 = 9 + 12 + 11 + 14 = 46$. The total contribution of $S = \emptyset$, $T = \{4\}$ is

$$b_2^{\{\{0\}, \{0, 1\}, \{0, 1\}, \{1\}\}} \cdot b_1^{\{\{0\}, \{1\}, \{0\}, \{0\}\}} \cdot b_2^{\{\{0\}, \{0, 1\}, \{0\}, \{0\}\}} \cdot b_4^{\{\{0\}, \{0\}, \{0\}, \{0\}\}} = 46 \cdot 3 \cdot 15 \cdot 66 = 136620.$$

Similarly, selecting $S = \{8\}$, $T = \emptyset$ contributes

$$b_2^{\{\{0\}, \{0, 1\}, \{0, 1\}, \{0\}\}} \cdot b_1^{\{\{0\}, \{0\}, \{1\}, \{0\}\}} \cdot b_2^{\{\{0\}, \{0, 1\}, \{0\}, \{0\}\}} \cdot b_4^{\{\{1\}, \{0\}, \{0\}, \{0\}\}} = 36 \cdot 3 \cdot 15 \cdot 89 = 144180$$

and selecting $S = \{8\}$, $T = \{4\}$ contributes

$$b_2^{(\{0\},\{0,1\},\{0,1\},\{1\})} \cdot b_1^{(\{0\},\{1\},\{1\},\{0\})} \cdot b_2^{(\{0\},\{0,1\},\{0\},\{0\})} \cdot b_4^{(\{1\},\{0\},\{0\},\{0\})} = 46 \cdot 4 \cdot 15 \cdot 89 = 245640.$$

$$\text{So } c_w = 2^2(71280 + 136620 + 144180 + 245640) = 2390880.$$

References

- [1] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [2] Nicolas Bourbaki. *Lie groups and Lie algebras. Chapters 4–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley.
- [3] C. Chevalley. Sur les décompositions cellulaires des espaces G/B . In *Algebraic Groups and their generalizations: Classical methods*, volume 56, Part 1 of *Proceedings of Symposia in Pure Mathematics (University Park, PA, 1991)*, pages 1–23. American Mathematical Society, 1994.
- [4] Persi Diaconis and Anil Gangolli. Rectangular arrays with fixed margins. In *Discrete probability and algorithms (Minneapolis, MN, 1993)*, volume 72 of *IMA Vol. Math. Appl.*, pages 15–41. Springer, New York, 1995.
- [5] Gérard Duchamp, Florent Hivert, and Jean-Yves Thibon. Noncommutative symmetric functions. VI. Free quasi-symmetric functions and related algebras. *Internat. J. Algebra Comput.*, 12(5):671–717, 2002.
- [6] James E. Humphreys. *Reflection groups and Coxeter groups*, volume 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.
- [7] Masato Kobayashi. Two-sided structure of double cosets in Coxeter groups, June 14, 2011. [Online; accessed 28-September-2015].
- [8] OEIS Foundation Inc. The On-Line Encyclopedia of Integer Sequences, 2015. Online. <http://oeis.org>.
- [9] T. Kyle Petersen. A two-sided analogue of the Coxeter complex. *In preparation*.
- [10] Edward Richmond and William Slofstra. Staircase diagrams and enumeration of smooth Schubert varieties. *Preprint*. arXiv: 1510.06060.
- [11] Louis Solomon. A Mackey formula in the group ring of a Coxeter group. *J. Algebra*, 41(2):255–264, 1976.
- [12] Richard P. Stanley. Weyl groups, the hard Lefschetz theorem, and the Sperner property. *SIAM J. Alg. Disc. Meth.*, 1(2):168–184, 1980.
- [13] Richard P. Stanley. *Enumerative Combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999.
- [14] John R. Stembridge. Tight quotients and double quotients in the Bruhat order. *Electron. J. Combin.*, 11(2):Research Paper 14, 41 pp. (electronic), 2004/06.