

Cryptosporidiosis: A zoonotic disease concern

Natapol Pumipuntu and Supawadee Piratae

One Health Research Unit, Faculty of Veterinary Sciences, Mahasarakham University, Maha Sarakham, Thailand.

Corresponding author: Supawadee Piratae, e-mail: bios.tah@gmail.com

Co-author: NP: natapol.p@msu.ac.th

Received: 23-01-2018, **Accepted:** 23-04-2018, **Published online:** 23-05-2018

doi: 10.14202/vetworld.2018.681-686 **How to cite this article:** Pumipuntu N, Piratae S (2018) Cryptosporidiosis: A zoonotic disease concern, *Veterinary World*, 11(5): 681-686.

Abstract

Cryptosporidiosis is considered to be a crucial zoonotic disease caused by worldwide distributing parasitic protozoa called *Cryptosporidium* spp. Cryptosporidiosis becomes a major public health and veterinary concern by affecting in human and various host range species of animals. Essentially, its importance of infection is increasing because of the high incidence in young children, immunocompromised persons, or immunodeficiency syndrome patients, especially in HIV/AIDS, and it is also one of the most causes of mortality in those patients who infected with *Cryptosporidium* spp. as well as young animals. All domestic animal, livestock, wildlife, and human can be potential reservoirs that contribute *Cryptosporidium* spp. to food and surface waters and transmitted to other hosts through fecal-oral route. The oocyst stage of *Cryptosporidium* spp. can remain infective and resistant to various environmental exposure and also resistant to many general disinfecting agents including chlorination which normally used in water treatment. Therefore, the understanding of these zoonotic pathogens is very essential in both animal and human health. This review focuses on the biology, life cycle, transmission, diagnosis, treatment, prevention, and control of this protozoan infection to emphasize and remind as the significant One Health problem.

Keywords: cryptosporidiosis, diarrhea, waterborne disease, zoonosis.

Introduction

Cryptosporidiosis is considered to be one of the most important diseases causing an intestinal infection to humans and animals [1]. The illness is primarily in mild diarrhea to severe in immune-compromised person, especially in HIV/AIDS, and patients who received immunosuppressive drugs. The etiological agent is the protozoan from *Cryptosporidium* genus [2] which has been reported that approximately 60% of them were the most common causative protozoan parasites of food and waterborne disease outbreak worldwide during 2004-2010 [3]. Human and animals get infected when they consume food and drink containing oocysts of these protozoa. Due to food and water transmission, incidence and prevalence of cryptosporidiosis are higher in less developed and developing countries where people are insufficient of basic infrastructure or fundamental facilities to avoid food and drinking water contaminated by infectious oocysts in feces [4]. In addition, the oocysts play a potential role to contribute *Cryptosporidium* spp. because they are tolerant to several chemicals and disinfectants including chlorine which is commonly used to treat in drinking water processing, also swimming pools, and water parks [5].

Twenty-three species and 61 valid genotypes of *Cryptosporidium* spp. have been described from a wide

range of vertebrates including humans, mammals, wildlife, domestic livestock, reptile, birds, amphibians, and fish with causing asymptomatic or mild-to-severe gastrointestinal disease in its host species [6]. Several species have been isolated from immunocompromised humans such as

Cryptosporidium parvum and *Cryptosporidium hominis* [7]. The most common three species of *Cryptosporidium* in human are

Cryptosporidium parvum, *Cryptosporidium hominis*, and *Cryptosporidium felis*; however, *Cryptosporidium parvum* and *Cryptosporidium hominis* are responsible for more than 90% of cryptosporidiosis cases [8]. Interestingly, almost 155 species of other mammals were reported as the non-human hosts of *Cryptosporidium* [9] which indicated that the parasites are adapting and developing to infect many diverse hosts and able to be one of the major zoonotic diseases problems.

Cryptosporidiosis has been informed as a significant cause of diarrhea worldwide. In Southeast Asia, cryptosporidiosis is frequently reported in both humans and animals from many countries such as Laos, Vietnam, Philippines, Myanmar, Indonesia, Cambodia, Malaysia, and Thailand [10]. In Thailand, 6% of ocean water samples and 11% of river samples are contaminated with *Cryptosporidium* spp. [11]. Moreover,

Cryptosporidium parvum and *Cryptosporidium hominis* have been isolated from HIV/AIDS cases in Bangkok [12] with the prevalence at 19-34% from 1996 to 2009 [13]. In other regions, cryptosporidiosis is also recognized as a major concern by the increasing cases of immune-compromised individuals and children in many countries of

the Americas such as Costa Rica, Argentina, Brazil, and the USA while the incidence of cryptosporidiosis in Europe is increasing by climate changes as heavy rainfall or flood and this pathogen can contaminate in drinking water. Therefore, the reports of cryptosporidiosis cases in Europe are higher than previous years, and *Cryptosporidium parvum* was identified as the most common pathogen [1]. *Cryptosporidium parvum* spp. is associated with severe diarrhea, mortality increasing, and negative impact of development and growth to the children in Africa. More than that, HIV/AIDS outbreak and malnutrition status of some developing countries in Africa can contribute the high prevalence of the contagion in this region [14].

Cryptosporidiosis in livestock is becoming the significant problems for animal health (both subclinical and clinical) and economic losses [15] because of increasing veterinary services and labor costs, increasing animal health-care cost, and decreasing a growth rate of animals and mortality of severe animals. Previous reports of cryptosporidiosis in livestock in Thailand were 31.5%, 5.7%, and 8.7% in dairy farms, individual animals, and dairy herd, respectively [16]. Moreover, 30% of buffalo farms were infected with *Cryptosporidium parvum* spp., and *Cryptosporidium parvum* was the most species contributed in livestock animal [17]. For other animals, the prevalence of cryptosporidiosis showed 2.1% in dog and 2.5% in cat. For wildlife animal such as in long-tailed macaques which live closely and normally contact to human communities in Thailand, it showed a prevalence about 1% but pose an important human health risk because infection in this animal can be occurred although less number of oocysts [18]. Here, we review the general information of *Cryptosporidium parvum* spp. which researches are needed for understanding the biology, life cycle, transmission, diagnosis, treatment, prevention, and control of these protozoa.

Morphology and Life Cycle of *Cryptosporidium* Species

Cryptosporidium parvum has a complex life cycle involving both sexual (meiosis) and asexual replication (mitosis) but monoxenous cycle [19]. Furthermore, it has many morphology formations to complete in its life cycle. Oocysts are excreted in the environment from human and animals through the feces. After the host ingests infective oocyst, excystation will occur to release four sporozoites. The sporozoites will invade the gastrointestinal tract and develop to merozoites, gamonts, and oocysts. Oocysts exist in two forms: Thin wall will reinfect in the gastrointestinal tract and thick wall will excrete in the environment through feces which shows in Figure-1. A detailed account of the life cycle starts from sporulated oocyst (which is rarely morphometric differences among different species) released by the infected host. After that, vertebrate host ingests sporulated oocyst through consumption of contaminated food or drink, and the process of excystation

will occur to release 4 infectious sporozoites. This progression occurs in the gastrointestinal tract triggered by CO₂, the temperature of 37°C, pancreatic enzymes, and pH acidic of bile salts [20].

Oocyst has two types, one is the thin wall (one layer of a protein-lipid-carbohydrate matrix), and another is the thick wall (consisting of inner and outer oocyst walls). Only the thick wall is shedding into the environment. The size of oocysts typical have nearly spherical with small size around 4-6 µm on the average and have obscure internal structures. The hallmark characteristic of mature oocysts contains four sporozoites but no sporocyst.

The sporozoites are nucleated, spindle shape with 5.0 µm×0.5 µm. The apical complex part plays function in the gliding motility to access the target cell. The nucleus is on the center of the cell. Sporozoites recognize and penetrate to the target host cells including stomach (*Cryptosporidium parvum* and *Cryptosporidium hominis*) and intestine (such as *Cryptosporidium parvum* and *Cryptosporidium hominis*) [7]. Invaded sporozoites are from parasitophorous vacuole surroundings where they can differentiate into trophozoites.

The trophozoites commence three mitosis divisions (merogony) to produce type I meronts which contain 8 merozoites (morphologically similar to

Type I merozoites can invade host cell again and reproduce asexual to form either type I meronts which contains eight merozoites or type II meronts which contain only four merozoites. Though, merozoites release from type II meronts are less uniform in shape, slightly larger and less active when compared to merozoites release from type I meronts.

Type II merozoites can be developed to microgamont (male) or macrogamont (female) which undergoes sexual reproduction (gametogony). Each microgamont produces sixteen microgametes which

size. It will fertilize with a unicellular adjacent macrogamont, a spherical to an oval structure of 4 to 6

two mitosis divisions, zygote develops either a thin-walled oocyst covering only a single layer membrane or a thick-walled oocyst containing two-layered membranes. The thick-walled oocysts are released through feces and tolerance for months in the unsuitable environment, whereas the thin-walled oocysts can cause reinfection within the gastrointestinal tract of the same host by rupturing and releasing infective sporozoites [23].

Transmission

Transmission routes of this parasite can be divided into two methods, direct and indirect transmissions. Direct transmission occurs through a fecal-oral route by accidental ingestion, and the *Cryptosporidium parvum* m oocytes excreted from feces. The

transmission happens between animal to animal, ani

DNA sequencing of PCR products [34], PCR restriction fragment length polymorphism [35,36], real-time PCR assays, nested PCR [37], multiplex PCR [38], and other modified PCR. Several genetic regions such as the 18s rDNA, oocyst wall protein, heat shock protein 70, Acetyl-CoA gene, b-tubulin gene, Cp15 gene, Cp 11 gene, dihydrofolate reductase inhibitors (), microsatellite loci, and 60 kDa glycoprotein () have been commonly used to identify isolates in animal and human infections [33,39]. Moreover, detection of cryptosporidiosis can be done with intestinal biopsy [40]. Infection part of the intestine can be stained with hematoxylin and eosin which will demonstrate parasites in each stage attached to the epithelial cells.

Treatment

According to the previous information, most patients and infected animals which have vigorous immune response can convalesce themselves without any treatments. Some clinical symptoms including watery fever, vomiting, nausea, stomach cramp, and dehydration can be managed by some supportive treatments as fluid and electrolyte replacement, anti-nausea drugs, antiemetic drugs, or analgesic drugs. Those medications can relieve the symptoms from cryptosporidiosis, but an antiprotozoal therapy is necessary in some cases.

Nitazoxanide is the most effective drugs for treating patients who infected with spp., and it is the only anti-cryptosporidial agent which has been approved for the treatment of cryptosporidiosis in humans by the US Food and Drug Administration. However, it is not commercially available or not yet widely used at the present time [1]. In addition, nitazoxanide cannot effective without a good immune response of the host so that it cannot be used effectively in immunocompromised patients [41]. For animals, they have few reports that studied the effect of nitazoxanide against clinical infections of in animals which demonstrated that nitazoxanide could decrease oocysts excretion [42,43]. Notwithstanding, it is also not yet generally used in animals.

Prevention

spp. can transmit and infect susceptible hosts by a major route of fecal to oral as cross-contamination in raw food and water from reservoir animals in community, farms, or abattoirs and some mechanical vectors such as cockroaches and flies [44]. Then, humans can get infected by their oocysts by ingesting this contaminated food and water with unprocessed or improper hygiene. Therefore, in a case of human prevention, the best strategy to prevent transmission of spp. is a practice of good personal hygiene including handwashing before preparing and consuming food, after using a toilet, and after contacting with diarrhea patients, children, and

some animals or livestock. Raw food and water must be suitably cleaned, washed, heated, cooked, or boiled before consumption and drinking, respectively. More than that, patients who have diarrhea symptom should realize not to swim in a public swimming pool, public water park, or river for preventing a transmission to others and people who swim in a swimming pool, water park, or river should recognize a potential risk of disease infection if they swallow the water [45].

Preventive measures that diminish transmission of spp. among animals, especially in livestock, should be emphasized on a primary prevention which reduce or eliminate causative risk factors by limiting the amount of animals density in the farms or stocks, minimizing a contraction between personnel, calves, and other herds, keeping young animals or susceptible hosts that have high risk of infection separated from adult animals, and keeping a short calving period of animals which may decrease the opportunities for spp. to spread within animal herds [46]. In human living areas, livestock husbandry, and their drinking water, the destruction of oocysts can be managed by heat or chemical disinfection such as hydrogen peroxide, sterilization processes using steam, ethylene oxide, chlorine dioxide, ozone (O₃), and ultraviolet light (UV light) which have been used to sterilize drinking water, but some of the disinfectants are not commonly practical procedure. However, all of the chemical disinfection can be used to prevent and control the occurrence of cryptosporidiosis and to reduce mortality and morbidity rate in both infected in both human and animals [47,48].

Conclusion

spp. distributes worldwide, especially in undeveloped and developing countries. The distribution has arisen when 's oocyst is defecated to water surface from human and animals (wildlife, domestic animals, and livestock) through the feces. The excreted oocysts are sustainable and tolerance with disinfectant, dilute bleach, and chlorine. Transmission occurs when hosts expose with 's oocyst mainly by fecal-oral route through contaminated food and water or contact directly with animal feces and indirectly by cross contamination. Laboratory diagnosis of infection normally uses oocyst staining and examines under microscopy. However, these methods cannot differentiate morphology of each species, so immunological methods and molecular techniques are playing more role for identification to species. Most of robust infected human and animals are asymptomatic and mild diarrhea, but violent symptoms will occur in the immunocompromised host. The best strategy to prevent and narrow down the spreading of this disease is keeping good personal hygiene in human coupled with reduction, control, or elimination the causative risk factors for other animals.

Authors' Contributions

NP and SP conceived the data, wrote the manuscript, reviewed and edited the manuscript. Both authors read and approved the final manuscript.

Acknowledgments

We thank One Health Research Unit (OHRU) and Faculty of Veterinary Sciences, Mahasarakham University for knowledge and data support. We also thank Miss Thamonphan Wimonrinarachai for picture design.

Competing Interests

The authors declare that they have no competing interests.

References

- Bamaiyi, P. and Redhuan, N. (2017) Prevalence and risk factors for cryptosporidiosis: A global, emerging, neglected zoonosis. *Journal of Parasitology*, 10: 309-325.
- Karanis, P., Kourenti, C. and Smith, H. (2007) Waterborne transmission of protozoan parasites: A worldwide review of outbreaks and lessons learnt. *Journal of Parasitology*, 5: 1-38.
- Baldursson, S. and Karanis, P. (2011) Waterborne transmission of protozoan parasites: Review of worldwide outbreaks—An update 2004–2010. *Journal of Parasitology*, 45: 6603-6614.
- Burnet, J.B., Penny, C., Ogorzaly, L. and Cauchie, H.M. (2014) Spatial and temporal distribution of *Cryptosporidium* in a drinking water resource: Implications for monitoring and risk assessment. *Water Science and Technology*, 472: 1023-1035.
- Fayer, R., Trout, J.M., Xiao, L., Morgan, U.M., Lal, A.A. and Dubey, J.P. (2001) *Cryptosporidium* n. sp. from domestic dogs. *Journal of Parasitology*, 87: 1415-1422.
- Ryan, U., Fayer, R. and Xiao, L. (2014) *Cryptosporidium* species in humans and animals: current understanding and research needs. *Journal of Parasitology*, 141: 1667-1685.
- Fayer, R. (2010) Taxonomy and species delimitation in *Cryptosporidium*. *Journal of Parasitology*, 124: 90-97.
- Xiao, L. and Ryan, U.M. (2004) Cryptosporidiosis: An update in molecular epidemiology. *Journal of Parasitology*, 17: 483-490.
- Šlapeta, J. (2013) Cryptosporidiosis and species in animals and humans: a thirty colour rainbow? *Journal of Parasitology*, 43: 957-970.
- Lim, Y.A., Mahdy, M.A. and Surin, J. (2013) Unravelling *Cryptosporidium* and *Isospora* in Southeast Asia. *Parasites and their Vectors*. Southeast Asia, and have been reported in countries such as Cambodia, Indonesia. p77-102.
- Koompaong, K. and Sukthana, Y. (2012) Seasonal variation and potential sources of *Cryptosporidium* contamination in surface waters of Chao Phraya River and Bang Pu nature reserve pier, Thailand. *Journal of Parasitology*, 43: 832-840.
- Srisuphanunt, M., Saksirisampant, W. and Karanis, P. (2011) Prevalence and genotyping of *Cryptosporidium* isolated from HIV/AIDS patients in urban areas of Thailand. *Journal of Parasitology*, 105: 463-468.
- Berger, S. (2017) Infectious Diseases of Thailand: 2017 Edition. GIDEON Informatics Inc., Thailand.
- Squire, S.A. and Ryan, U. (2017) *Cryptosporidium* and *Isospora* in Africa: Current and future challenges. *Journal of Parasitology*, 10: 195.
- Santin, M. (2013) Clinical and subclinical infections with *Cryptosporidium* in animals. *Journal of Parasitology*, 61: 1-10.
- Jittapalpong, S., Pinyopanuwat, N., Chimnoi, W., Siripanth, C. and Stich, R.W. (2006) Prevalence of *Cryptosporidium* among dairy cows in Thailand. *Journal of Parasitology*, 1081: 328-335.
- Inpankaew, T., Jiyipong, T., Wongpanit, K., Pinyopanuwat, N., Chimnoi, W., Kengradomkij, C., Xuan, X., Igarashi, I., Xiao, L. and Jittapalpong, S. (2014) Molecular detection of *Cryptosporidium* spp. infections in water buffaloes from northeast Thailand. *Journal of Parasitology*, 46: 487-490.
- Srichareern, W., Inpankaew, T., Keawmongkol, S., Supanam, J., Stich, R.W. and Jittapalpong, S. (2016) Molecular detection and prevalence of *Cryptosporidium* and *Isospora* spp. among long-tailed macaques (*Macaca fascicularis*) in Thailand. *Journal of Parasitology*, 40: 310-314.
- Bouazid, M., Hunter, P.R., Chalmers, R.M. and Tyler, K.M. (2013) *Cryptosporidium* pathogenicity and virulence. *Journal of Parasitology*, 26: 115-134.
- O'Donoghue, P.J. (1995) *Cryptosporidium* and cryptosporidiosis in man and animals. *Journal of Parasitology*, 25: 139-195.
- Cacciò, S.M. and Widmer, G., editors. (2013) *Cryptosporidium*: Parasite and Disease. Springer Science & Technology.
- Current, W. and Reese, N.C. (1986) A comparison of endogenous development of three isolates of *Cryptosporidium* in suckling mice. *Journal of Parasitology*, 33: 98-108.
- Tzipori, S. and Ward, H. (2002) Cryptosporidiosis: Biology, pathogenesis and disease. *Journal of Parasitology*, 4: 1047-1058.
- Hubalek, Z. (2003) Emerging human infectious diseases: Anthroponoses, zoonoses, and sapronoses. *Journal of Parasitology*, 9: 403-404.
- Xiao, L. and Feng, Y. (2008) Zoonotic cryptosporidiosis. *Journal of Parasitology*, 52: 309-323.
- Pedersen, C., Danner, S., Lazzarin, A., Glauser, M.P., Weber, R., Katlama, C., Barton, S.E. and Lundgren, J.D. (1996) Epidemiology of cryptosporidiosis among European AIDS patients. *Journal of Parasitology*, 72: 128-131.
- Jiang, J., Alderisio, K.A. and Xiao, L. (2005) Distribution of *Cryptosporidium* genotypes in storm event water samples from three watersheds in New York. *Journal of Parasitology*, 71: 4446-4454.
- Alles, A.J., Waldron, M.A., Sierra, L.S. and Mattia, A.R. (1995) Prospective comparison of direct immunofluorescence and conventional staining methods for detection of *Cryptosporidium* and *Isospora* spp. in human fecal specimens. *Journal of Parasitology*, 33: 1632-1634.
- Chalmers, R.M. and Katzer, F. (2013) Looking for *Cryptosporidium*: The application of advances in detection and diagnosis. *Journal of Parasitology*, 29: 237-251.
- Koonakositt, R., Sriurairatna, S. and Petchclai, B. (1992) Microscopic examination of *Cryptosporidium* oocysts in diarrhoeal stools. *Journal of Parasitology*, 75: 180-184.
- Campbell, A.T., Haggart, R., Robertson, L.J. and Smith, H.V. (1992) Fluorescent imaging of *Cryptosporidium* using a cooled charge couple device (CCD). *Journal of Parasitology*, 16: 169-174.
- Pacheco, F.T., Silva, R.K., Martins, A.S., Oliveira, R.R., Alcântara-Neves, N.M., Silva, M.P., Soares, N.M. and Teixeira, M.C. (2013) Differences in the detection of *Cryptosporidium* and *Isospora* oocysts according to the fecal concentration or staining method used in a clinical laboratory. *Journal of Parasitology*, 99: 1002-1008.
- Fayer, R., Morgan, U. and Upton, S.J. (2000) Epidemiology of *Cryptosporidium*: Transmission, detection and identification. *Journal of Parasitology*, 30: 1305-1322.
- Simonato, G., Frangipane di Regalbano, A., Cassini, R., Traversa, D., Tessarin, C., Di Cesare, A. and Pietrobelli, M. (2017) Molecular detection of *Cryptosporidium* and *Isospora* spp. in canine faecal samples contaminating public areas in Northern Italy. *Journal of Parasitology*, 116: 3411-3418.
- Xiao, L., Escalante, L., Yang, C., Sulaiman, I., Escalante, A.A., Montali, R.J., Fayer, R. and Lal, A.A. (1999) Phylogenetic analysis of *Cryptosporidium* parasites

- based on the small subunit rRNA gene locus. *Journal of Clinical Microbiology*, 65: 1578-1583.
36. Xiao, L., Morgan, U.M., Limor, J., Escalante, A., Arrowood, M., Shulaw, W., Thompson, R.C., Fayer, R. and Lal, A.A. (1999) Genetic diversity within *Shigella* and related species. *Journal of Clinical Microbiology*, 65: 3386-3391.
 37. Abdelsalam, I.M., Sarhan, R.M. and Hanafy, M.A. (2017) The impact of different copro-preservation conditions on molecular detection of *Shigella* species. *Journal of Clinical Microbiology*, 12: 274-283.
 38. Mero, S., Kirveskari, J., Antikainen, J., Ursing, J., Rombo, L., Kofoed, P.E. and Kantele, A. (2017) Multiplex PCR detection of *Shigella* sp., *Shigella* and *Shigella* directly from dried stool samples from Guinea-Bissauan children with diarrhoea. *Journal of Clinical Microbiology*, 49: 655-663.
 39. Khan, A., Shaik, J.S. and Grigg, M.E. (2017) Genomics and molecular epidemiology of *Shigella* species. *Journal of Clinical Microbiology*, 176: 349-354.
 40. Doganci, T., Araz, E., Ensari, A., Tanyuksel, M. and Doganci, L. (2002) Detection of *Shigella* infection in childhood using various techniques. *Journal of Clinical Microbiology*, 8: MT223-MT226.
 41. Gargala, G. (2008) Drug treatment and novel drug target against *Shigella*. *Journal of Clinical Microbiology*, 15: 275-281.
 42. Viel, H., Rocques, H., Martin, J. and Chartier, C. (2007)