



NIH PUBLIC ACCESS

Author Manuscript

Circ Res. Author manuscript; available in PMC 2010 September 11.

Published in final edited form as:

Circ Res. 2009 September 11; 105(6): 585–594. doi:10.1161/CIRCRESAHA.109.200451.

Loss of cardiac microRNA-mediated regulation leads to dilated cardiomyopathy and heart failure

Prakash K. Rao¹, Yumiko Toyama², H. Rosaria Chiang¹, Sumeet Gupta¹, Michael Bauer³, Rostislav Medvid⁴, Ferenc Reinhardt¹, Rongliu Liao³, Monty Krieger², Rudolf Jaenisch^{1,2}, Harvey F. Lodish^{1,2,5}, and Robert Blümel^{4,5}

¹Whitehead Institute for Biomedical Research, Cambridge, MA 02142²Dept of Biology, MIT, Cambridge, MA 02142³Division of Cardiology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115⁴Eli and Edythe Broad Center for Regeneration Medicine and Stem Cell Biology, Department of Urology, University of California, San Francisco, California 94143, USA.

Abstract

RATIONALE—Heart failure is a deadly and devastating disease that places immense costs on an aging society. In order to develop therapies aimed at rescuing the failing heart, it is important to understand the molecular mechanisms underlying cardiomyocyte structure and function.

OBJECTIVE—microRNAs are important regulators of gene expression and we sought to define the global contributions made by microRNAs toward maintaining cardiomyocyte integrity.

METHODS AND RESULTS—First, we performed deep sequencing analysis to catalog the miRNA population in the adult heart. Secondly, we genetically deleted, in cardiac myocytes, an essential component of the machinery that is required to generate miRNAs. Deep sequencing of miRNAs from the heart revealed the enrichment of a small number of microRNAs with one, miR-1, accounting for 40% of all microRNAs. Cardiomyocyte-specific deletion of *dgc8*, a gene required for microRNA biogenesis, revealed a fully penetrant phenotype that begins with left ventricular malfunction progressing to a dilated cardiomyopathy and premature lethality.

CONCLUSIONS—These observations reveal a critical role for microRNAs in maintaining cardiac function in mature cardiomyocytes and raise the possibility that only a handful of microRNAs may be ultimately responsible for the dramatic cardiac phenotype seen in the absence of *dgc8*.

Introduction

A subset of endogenous non-coding small RNAs, known as microRNAs (miRNAs or miRs) ¹ are ~22 nucleotides long and modulate gene expression by targeting mRNAs for post-transcriptional repression. There are nearly 500 and 800 microRNAs in mice and humans respectively² (<http://microrna.sanger.ac.uk>). In animals, repression is achieved through imperfect base-pairing between the microRNA and its target mRNA. Although there are certain

⁵**Contact information for corresponding authors:** Harvey F. Lodish, Ph.D. Professor of Biology, MIT; Member, Whitehead Institute; Professor of Bioengineering, MIT Whitehead Institute for Biomedical Research 9 Cambridge Center Cambridge, MA 02142 lodish@wi.mit.edu Robert Blümel, M.D., Ph.D. The Eli and Edythe Broad Center of Regeneration Medicine and Stem Cell Research, Center for Reproductive Sciences, and Department of Urology, University of California, San Francisco, San Francisco, California, USA. blumel@stemcell.ucsf.edu

DISCLOSURES None

rare instances in which microRNAs have been reported to upregulate target gene expression^{3,4}, repression is the most well-documented direct effect. The target mRNA is rendered labile through mechanisms involving deadenylation/decapping, translational repression or both. Target specificity is largely governed by the highly conserved seed region (nucleotides 2-8) of the miRNA⁵. Various target prediction programs have relied on this fact and an estimated 30% of the mRNAs are susceptible to miRNA-mediated regulation⁶. Although this number is likely an overestimate, as it does not take into account the requirement for coexpression of miRNAs and mRNAs, it puts into perspective the enormous regulatory potential possessed by microRNAs. Not surprisingly, a number of studies have revealed the importance of the microRNA pathway as a whole while others have pinpointed specific roles for individual microRNAs in various tissues⁷⁻¹⁸.

Although mature microRNAs are only ~22 nucleotides in length, they are generated from longer precursors whose length distribution is similar to that of a mRNA. Indeed, the primary transcripts (pri-miRNAs) are transcribed by RNA Polymerase II, capped, polyadenylated and regulated by transcription factors like protein-coding mRNAs¹⁹⁻²¹. Unlike mRNAs, miRNAs — because of their stem-loop structure — are cleaved within the nucleus by a Drosha/Dgcr8 containing complex into ~60-80 bp precursor miRNAs (pre-miRNAs)²²⁻²⁴. The precursor miRNAs are transported out of the nucleus by Exportin-5²⁵ and subsequently processed by a cytoplasmic RNaseIII--Dicer²⁶ — which also resides in a multiprotein complex. Since the PAZ domain of Dicer recognizes the two-nucleotide 3'OH overhang generated by Drosha/Dgcr8, it is believed that the nuclear Drosha/Dgcr8 cleavage is required for Dicer-mediated cytoplasmic cleavage of pre-miRNAs. Exceptions to this general dependence on Drosha/Dgcr8 occurs in miRtrons and endogenous shRNAs²⁷⁻²⁹ and in these rare cases, other nucleases generate the necessary ends for subsequent Dicer recognition and cleavage. Importantly, spatial segregation of Drosha/Dgcr8 and Dicer substrates allows for the two cleavage events to occur in a sequential manner.

We sought to uncover the regulatory potential of miRNAs in the heart by using two complementary approaches. First we catalog the known miRNA population of murine adult heart using deep (Solexa/Illumina) sequencing of a small RNA library. Secondly, we disrupt microRNA regulation by deleting *dgcr8* and hence canonical microRNA biogenesis. We chose to focus on **mature** muscle tissue in order to establish the importance of microRNA function in the *maintenance* (as opposed to the development) of cardiac tissue. Mice lacking *dgcr8* in muscle tissue die prematurely with signs of heart failure and dilated cardiomyopathy. Identification of the depleted microRNAs in *dgcr8* deficient hearts led to the refined list of microRNA targets that may collectively play an important role in the development of the pathological state. Thus, the importance of the microRNA regulation in maintaining cardiomyocyte function is revealed by the fatal outcome associated with lack of *dgcr8* in cardiomyocytes.

Materials and Methods summary

Details are included in the online supplement. Briefly a library of small RNAs were generated and sequenced using the Illumina platform³⁰. For the generation of conditional *dgcr8* knockout, floxed *dgcr8* mice was crossed with Muscle Creatine Kinase (MCK)-Cre mice³¹; mutant mice were genotyped using tail DNA by a PCR-based approach. Age and sex-matched mutant (2lox/2lox; Cre positive) and control (2lox/+; Cre positive) mice were analyzed pathologically; physiological studies were performed using telemetry and echocardiography. Molecular analyses were carried out using total RNA isolated from the heart; Northern blots were used to detect depletion of miR-1, miR-133 and miR-208. Array-based methods were employed to assess global loss of microRNAs.

Results

Deep sequencing of microRNAs from heart tissue

High throughput deep sequencing produces quantitative data with an extensive dynamic range, thereby enabling detailed insight into the relative levels of different microRNA in a particular tissue. Therefore, to gain such insight into the microRNA profile of the adult heart, we isolated small RNAs (16-24 nucleotides) from 6-8 week old male and female hearts, built tagged cDNA libraries and sequenced the libraries on a Illumina Genome analyzer producing over seven million reads from each sample. As has been reported previously, miR-1 and miR-133a were highly abundant (Fig 1A)--however the *relative* abundance of miR-1 reads was quite striking. miR-1 accounted for nearly 40% of all known microRNA reads. Also noteworthy is the fact that other microRNAs including miR-29a, miR-26a, let-7 family members, were more abundant than miR-133a. MicroRNAs from non-cardiomyocytes (miR-29a and miR-29c from fibroblasts 32, miR-126 from endothelial cells^{9,33}) also contributed to the library as expected since the different cell types in the heart were not separated. Within the cardiac-specific miR-208 subsets, 50-100 times more reads were obtained for miR-208a (encoded within an intron of Myh6) when compared to miR-208b (encoded within an intron of Myh7), consistent with the relative overexpression of Myh6 compared to Myh7 in adult mice. miR-22 was highly expressed and showed gender-based differences in expression levels. While sexually dimorphic gene expression patterns in somatic tissues ³⁴ has been established³⁴; follow-up experiments will need to be carried out to confirm sex-based differences in miRNA expression in the heart. Reads from the miR-378 hairpin were also high; miR-378/378* (miR-378* is the same as miR-422b) is encoded within an intron of the PGC1b gene. Since PGC1a and PGC1b regulate mitochondrial biogenesis and the heart is a mitochondria-rich organ, the high expression levels of miR-378/miR-378* probably reflects the high endogenous levels of PGC1b transcription. Since the ability of a miRNA to repress target gene expression is largely dependent on the 5' end of the miRNA, multiple miRNAs with identical 5' ends are expected to function in a similar manner. This seed identity is the basis by which microRNAs are grouped into families. Therefore, we tabulated all the microRNA reads within individual families (as defined by TargetScan 4.1; www.targetscan.org) (Fig 1B). By this analysis, the miR-1/206 family still emerged as the most dominant microRNA family (the reads from miR-206 were insignificant). Since a considerable number of reads were obtained individually from members of the let7/miR-98 and miR-30a-5p family, these families were respectively, the second and third most abundant microRNA families in the heart.

Muscle-specific Dgcr8 knockout

The importance of the microRNA pathway during development has been largely inferred from studies in which Dicer has been deleted ^{18,35-39}. As dicer has roles outside of the canonical miRNA pathway, we sought to block microRNA maturation (and therefore microRNA-mediated regulation) using another component of the microRNA biogenesis pathway, namely Dgcr8. Dgcr8 deletion in embryonic stem cells has revealed that it is essential for microRNA biogenesis and implicate microRNAs in regulating efficient ES-cell differentiation ⁴⁰. Using MCK-Cre mice³¹ and a conditional floxed allele of *dgcr8*, we generated mice with a muscle-specific deletion of the *dgcr8* gene. Endogenous MCK expression reportedly peaks around birth and declines to 40% of peak levels by day ten ³¹. This Cre line was deliberately chosen to match our interest in specifically disrupting microRNA biogenesis in mature, differentiated muscle as this allowed us to determine the importance of the microRNA pathway in muscle homeostasis. Genotyping analysis showed that although mutant (2lox/2lox; Cre positive) mice were slightly underrepresented at the time of genotyping, most mutant mice survived to at least 12 days after birth. We did not observe any pathology on 4-chamber sections (H & E stained) at two weeks of age. At three weeks of age, we detected fibrosis in the ventricular wall in all mice examined, and loss of ventricular function (as revealed by trans-thoracic

echocardiography—see below). Subsequently, all mutant mice died before two months of age and the median survival was 31 days (Figs. 2A & 2B). At end stage, the hearts of mutant mice showed marked decreases in the thickness of the left and right ventricular walls. Therefore the development of the pathology is quite rapid and highly penetrant. This demonstrates the stringent requirement for a threshold level of microRNAs below which heart function rapidly deteriorates.

In order to determine the extent of microRNA depletion in the heart, we performed Northern Blot and quantitative RT-PCR analyses to quantify cardiomyocyte-specific microRNAs (Fig. 2C) with RNA derived from the heart tissue of mutant and control (2lox/+; Cre positive) mice. At the time of sacrifice (when mutant mice were moribund), Northern Blot analysis showed that three cardiac-enriched mature microRNAs (miR-1, miR-133a, and miR-208) were dramatically depleted, but not completely absent, in mutant heart tissue. Quantification of the Northern blots revealed that depending on the microRNA, the mature forms were depleted 10 to 60 fold (Fig. 2C, **bottom**). Their precursor miRNAs (the ~ 60 base pair product of Drosha/Dgcr8 cleavage) were detectable in the control lanes and absent in the mutant lanes (Fig. 2D; shown for miR-208). The complete loss of the short-lived ~60 bp precursor, but not mature miRNA favors the argument that the residual amount of mature miRNA detected is due to its long half-life, rather than an incomplete excision of *dgcr8* in these tissues.

The hearts of the mutant animals exhibited a variety of abnormalities that suggest cardiac dysfunction was responsible for their premature death. Preliminary ECG analysis of revealed dramatic drops in the heart rate of mutant mice along with an increased PQ interval and QRS width (all at end stage) indicative of a cardiac conduction defect (Online Fig S4). Histopathological analysis revealed that the hearts obtained from end-stage mutant mice were considerably enlarged with notable thinning of the ventricular walls (Fig. 3A & 3B; note end-stage mutant hearts). Fibrosis was also evident (Fig. 3C), an early and consistent pathological finding as it is observed in all mice at about three weeks of age (at which time there was no histopathologically obvious defect in the thickness of the ventricular wall; Fig. 3B and 3C). Quantitative RT-PCR analyses of cardiomyocyte-specific microRNAs was also carried out at end stage and at two weeks after birth. Precipitous decreases in miR-1, miR-133a and miR-208 levels was detected in two week old mice (Fig. 3D) and this preceded any pathophysiological changes that we observed.

In order to assess left ventricular function, we performed echocardiography. We conducted these studies at two timepoints: three weeks and four weeks after birth as histopathological analysis showed a dramatic progression between these two timepoints from mild fibrosis with otherwise no overt ventricular/wall defects (at three weeks) to extensive dilation (at four weeks). Accordingly, measurement of fractional shortening (FS) revealed that the mutant mice had dramatically reduced ventricular function at four weeks (Online Fig. S6); although wall thickness was not significantly different. This finding was not surprising considering the clear histopathological defects at this timepoint. The expectation at three weeks (Fig. 4B) was more ambiguous since we noted fibrosis at this point but did not see an obvious defect in wall thickness or ventricular volume in tissue sections. However, echocardiography at three weeks after birth, revealed that ventricular function (as assessed by FS readings) was decreased in mutant mice and the trend toward increased ventricular volume was already evident (see numbers for EDD at three weeks in the table in Fig. 4B).

Given the defects in ventricular function, one plausible explanation is that the myofibrillar apparatus was disorganized to the extent that contraction was ineffective. Such disarray has been noted in mice bearing a cardiac-specific loss of function allele of *dicer*¹³. Ultrastructural analysis (Online Fig. S5) revealed mild myofibrillar disarray mostly related to misalignment of the contractile apparatus. In order to determine whether pathology-associated cardiac

markers are expressed and fetal genes are activated, we performed real time RT-PCR analysis. Nppa and Nppb were expressed at higher levels in mutant heart (Fig. 5). Myh7, a fetal myosin whose re-expression in adulthood is associated with heart failure, was also expressed at higher levels in the mutant hearts. Myh6, the normal adult cardiac myosin, within which miR-208 is encoded, was expressed at similar levels in control and mutant hearts; thus the decrease seen in mature miR-208 is not attributable to differences in the regulation of the host gene. These molecular assays complement the pathological and echocardiographic observations and are consistent with a diagnosis of DCM.

Next, we isolated RNA from mutant and control hearts to examine the expression of marker genes expressed in striated muscle (Fig. 5). Cardiac markers were uniformly low in the mutant heart. In contrast, *fast skeletal* muscle markers were uniformly upregulated. One of the three slow skeletal markers (Tnni1) was also upregulated, while two (Tnnt1 and Tnnc1) were not; intriguingly Tnni1 also has a miR-133 binding site in its 3' UTR and part of its upregulation may be attributable to the loss of miR-133. The upregulation of skeletal muscle genes has been previously noted in other miR knockout mice^{8,10}. As misexpression of skeletal muscle isoforms in the heart can lead to impaired cardiac function⁴¹, at least part of the observed pathology may be attributed to increased expression of fast skeletal muscle transcripts at the expense of cardiac genes.

An array-based profiling approach was carried out to compare relative levels of mature microRNAs in RNA derived from the hearts of mutant and control mice. MicroRNAs that are less abundant in the mutant heart when compared to the control heart are likely to be those enriched in cardiomyocytes (as *dgcr8* is knocked out only in cardiomyocytes). Hence this analysis allows us to indirectly detect cardiomyocyte-enriched microRNAs. As expected, we detected precipitous declines in the levels of cardiomyocyte-specific miR-1, miR-133, miR-208 and miR-499 in the mutant hearts (Online Fig S7, **compare to** Fig 2C). Others that were decreased by greater than 2 fold, and therefore likely to be enriched within the cardiomyocytes include miR-378/miR-378* (aka miR-422b), miR-22, miR-486, miR-30e*, miR-149, miR-709, miR-345 and members of the miR-30a-5p family (Online Fig. S7).

To uncover the scope of regulation that is disrupted by the loss of the microRNA pathway, we carried out an *in silico* analysis. We chose ten microRNAs that were downregulated the most and used Targetscan to obtain a target list of mRNAs with conserved miRNA binding sites. Next we extracted a published dataset⁴² that had compiled the list of genes that are expressed in the human heart. The intersection of these two lists (Online Suppl Table 1) provided us with a list of genes whose expression could be upregulated in the hearts of mutant mice. This analysis suggests that approximately 14% (1140/7896) of the genes expressed in the heart could be potentially upregulated due to the loss of these ten microRNAs that we determined to be cardiomyocyte-enriched. Included amongst this list of targets are genes that are involved in GPCR signaling (endothelin receptors), calcium signaling (Calcineurin subunits), smooth muscle contraction (Mylk) and calcification (Runx2). Thus it is likely that the complex phenotype is at least in part due to the misexpression of a subset of these genes.

DISCUSSION

miR-1 and miR-133a

While many studies utilizing inter-tissue comparisons can attest to the abundance of miR-1 within muscle tissue, our study, by focusing on the intra-tissue abundance has revealed a wide disparity between the levels of miR-1 and all the other microRNAs. This is especially significant when we compare miR-1 to miR-133 which are co-regulated⁴³, (albeit differentially spliced⁴⁴) microRNAs. These results suggest that mechanisms other than transcription (eg. processing or stability) can dramatically alter steady state levels of mature

miRNAs. Recent evidence from other labs have shown such post-transcriptional regulation for let-7 and miR-2145⁴⁸. A second testament to the relative abundance of miR-1 is evident when comparing the levels of mature miR-1 and miR-208a. Since miR-208a is resident within a cardiac myosin, its levels should be representative of a highly transcribed miRNA. The fact that miR-1 levels far exceed that of miR-208a provides further indirect evidence for post-transcriptional mechanisms governing microRNA stability. Lastly, as we are sampling a multitude of cell types in the heart, it is possible that within cardiomyocytes, the percentage of miR-1 in relation to other microRNAs is even higher and further studies, using purified cardiomyocytes will be needed to verify this possibility. The very high levels of miR-1 suggests it plays a central role in sustaining heart muscle function; indeed previous analysis of a miR-1-2 knockout mouse ⁸ confirmed the importance of miR-1 dosage in maintaining proper cardiac function. The homozygous loss of miR-1-2, one of two mir-1 loci, causes multiple defects in heart function⁸. We await the generation of appropriate conditional knockout mice lacking both miR-1-1 & miR-1-2 mice to ascertain its singular importance in adult mouse myocardium. However, recent findings evaluating the mir-133a knockout mice clearly define its importance in cardiac development. Taken together, miR-1 and miR-133 appear to be attractive candidates for rescuing the phenotype associated with *Dgcr8* loss.

Our sequencing revealed that the miRBase annotation for miR-133a is offset by one nucleotide at the 5' end. The mature miRNA (which is the one that has the most common 5' end and is read most often) is UUGGUCCCCUUAACCAGCUGU; the miRBase annotated miR-133a is UUUGGUCCCCUUAACCAGCUG. Given the importance of the 5' end in determining target repression, this also changes the putative targets that may be repressed by miR-133a. We did obtain significant number of counts for the miR-133 species annotated on miRbase; however based on our criteria for miRNA classification, we denote the species with one less U as the mature miRNA. Other independent sequencing data (R.C, unpublished data) confirm this 5' heterogeneity of mature miR-133a.

We also performed microarrays to determine the fold change in microRNA levels in mutant versus wild-type hearts. This strategy is particularly powerful as it enables the specific identification of microRNAs that are present in the cells expressing the cre transgene. Of note, the ranking of genes that were most dramatically reduced in the mutant hearts as determined by microRNA microarrays did not directly match with the relative amounts of individual microRNAs uncovered in the sequencing data. This difference is likely the result of a number reasons. First, the sequencing data represents the miRNAs in all the cell types of the heart, rather the cells expressing the transgene. Second, the absolute level of any microRNA will not necessarily correlate with the fold decrease following *Dgcr8* loss as different miRNAs in the cardiomyocytes will almost certainly have different half-lives. Third, we isolated RNA from the hearts of slightly different aged mice for sequencing (6-8 week) versus the microarrays (4-5 weeks). Hence age-related differences may partly explain the differences between the array and the sequencing data sets.

Phenotype of muscle-specific *Dgcr8* KO mice

We have utilized a loss of function allele of *dgcr8* to uncover the importance of the microRNA pathway in cardiac integrity. The phenotypic outcome is similar to the cardiac-specific *dicer* deficient mice ¹⁸ and this similarity in phenotypes has also been shown in mice bearing conditional alleles of *dgcr8* and *dicer* in the skin⁴⁹. However, *dgcr8* deficient mice have an advantage over *dicer* deficient as the former can potentially be rescued by a Dicer-substrate shRNA designed to produce mature miRNAs. Hence it will be possible to define, in a fairly straightforward manner, the “minimal microRNA” requirements for different cell types derived from these mice. This approach has been successfully employed to reveal microRNAs

important in the cell cycle regulation of murine ES cells⁵⁰ and such an approach should be feasible in other cell types too.

The results from the muscle-specific *dgcr8* knockout mice demonstrate the essential role of microRNA regulation in cardiac function. Although we have not identified the root cause of DCM and HF in our mice, our data clearly demonstrates a role for the microRNA pathway in proper functioning of the heart. Changes in ventricular diameters were further visible in trans-thoracic echocardiography. This resulted in a significant decrease in mutant ventricular function as assessed by fractional shortening at four weeks of age compared to control littermates. Furthermore trans-thoracic echocardiography revealed functional deterioration was already present at three weeks of age with mutant mice showing markedly reduced fractional shortening. This functional deterioration preceded dilation seen at 4 weeks of age (Fig. 4). Heart rates and wall thickness were not significantly different in mutant mice (both at three weeks and at four weeks) supporting our histological observations that ruled out a hypertrophic phase prior to dilation. Ventricular walls from mutant mice did however, exhibit a thinning trend that was detectable by echocardiography at four weeks (Fig. 4B) and was very obvious histologically in end stage mice (Fig. 3B). Changes in microRNA levels have been noted to be secondary consequences of a stressed heart⁵¹. We demonstrate that cardiomyocyte-specific microRNA levels are depleted prior to the occurrence of pathophysiological changes (Fig. 3D). These data are consistent with the microRNA loss being causative and representing a primary event in the emergence of the phenotype we have described.

In comparison to the other single-microRNA knockouts^{8,10,52}, the *dgcr8* knockout exhibits a much more severe and penetrant phenotype. This is to be expected as a number of microRNAs are affected. Using a candidate gene approach and incorporating results from previously published work with the miR-208 Δ/Δ mice¹⁰, we interrogated and detected the upregulation of several fast skeletal muscle genes in the heart. As cardiac muscle is more akin to a “slow” muscle the aberrant activation of fast skeletal genes could be pathological. Even though myofibrillar proteins are homologous, each striated muscle tissue has evolved to meet its particular needs and previous studies have demonstrated that cardiac-specific overexpression of skeletal muscle specific protein can cause loss of cardiac function⁴¹. Clearly, a widespread increase in skeletal gene expression, suggested by our candidate gene analysis (Fig 5), could contribute to the loss of cardiac function. Another aspect of pathological remodeling is the re-establishment of a fetal gene program in failing cardiomyocytes. Clearly this is also a consequence of *Dgcr8* loss as exemplified by an increase in *Myh7*, a fetal myosin.

Comparison with other microRNA deficient mice

In comparing our phenotype to the heart-specific Dicer deficient mice (using the alpha-MHC promoter driven Cre)¹³ we note the following important differences. We always detect fibrosis (Fig. 3C) and see marked increases in MYH7 expression (Fig. 5) in mutant mice. These could be due to differences in the timing of Cre-mediated excision, implying that loss of microRNA regulation at different times lead to different phenotypic outcomes. We also note that the recent report describing the knockout of *dicer* in the adult myocardium¹⁸ shows a broadly similar, but not an identical phenotype, to the *dgcr8* KO mice described herein. Importantly, we did not detect any hypertrophy during routine pathological staining and the pathology of our mutant mice is more consistent with dilated cardiomyopathy that eventually leads to a phenotype that resembles human heart failure. Since the timing of the cre-mediated excision can cause different phenotypic outcomes¹⁸, direct comparison of the two *dicer* KO studies with this *dgcr8* KO (utilizing different promoters driving Cre) is complicated; nonetheless all these studies point to the clear importance for the microRNA pathway in cardiac function. A recent report from the Olson group reported the phenotype of a complete knockout of miR-133a⁵². Pathologically, the *dgcr8* muscle KO mice are very similar to the fraction of 133a-1^{ko};

133aa-2^{ko}(133a dKO) mice that survive to 2 or 4 months of age with fibrosis and ventricular wall thinning as common features. Similar to the 133adKO mice, we did not detect any gross hypertrophy prior to the advent of dilation; however, in contrast to the 133a dKO mice, we detect only mild myofibrillar disarray in our ultrastructural analysis. In both the 133a dKO mice and the cardiac-specific dicer KO mice, deletion of the cognate genomic region occurs early and hence may have a more profound effect on the arrangement of myofibrils. When *dgcr8* expression (and therefore microRNA-mediated regulation) is perturbed **after** the establishment of the myofibrillar array, the requirement for an intact microRNA pathway may be less stringent (as is the case here). Nonetheless, overall pathological similarity suggests that one reason for the myopathy seen in the *dgcr8* mice could be due to its lack of mature miR-133a. Fast skeletal gene expression has also been noted in the hearts of miR-208^{-/-} mice and this particular phenotypic characteristic could be a consequence of low levels of miR-208. However, miR-208^{-/-} mice do not lose cardiac function and additional microRNAs have to be implicated in describing the complete phenotype associated with loss of *Dgcr8* in the heart.

Cardiac heterochrony as a possible mechanism underlying the dramatic phenotype

An equally feasible and alternative (albeit speculative) explanation for the sustained expression of fetal gene markers in adulthood is that the lack of the microRNA pathway leads to an arrest in the development of the heart such that late embryonic/prenatal gene expression patterns are maintained in the adult. Indeed, such heterochronic events in *C.elegans* were instrumental in the identification of the founding members of microRNAs⁵³⁻⁵⁵ and recent reports have confirmed that this heterochronic pathway is conserved⁵⁶⁻⁵⁸. Two candidate genes that we have examined are normally repressed in adult tissues but continue to be expressed when *Dgcr8* is absent: *Myh7* (a fetal myosin) and *Tnni* (a slow skeletal muscle-specific troponin-complex subunit). Under normal physiological conditions, *Tnni* levels are downregulated in the heart following birth⁵⁹, but it continues to be expressed at relatively higher levels in the *dgcr8* KO heart. Importantly, *Tnni*, as opposed to *Myh7*, is NOT expressed at higher levels in a pathological state⁵⁹ and, therefore, its overexpression cannot be attributed to the cardiac reprogramming that occurs secondary to a failing heart. This observation suggests that the lack of the *dgcr8* gene can cause a heterochronic phenotype (which in turn is incompatible with adult heart function). In addition, our *in silico* analysis (Online Suppl Table 1) suggests that a large number of genes that are expressed in the heart are susceptible to microRNA-mediated regulation. Comparison of temporal mRNA expression profiles in mutant mice and wild type mice will aid in the identification of the primary targets of the microRNA pathway as well as to provide evidence for the existence of cardiac heterochrony in the mutant mice.

In summary, we have, through complementary approaches, ascertained the importance of the microRNA pathway in maintaining cardiomyocyte function. We have identified high abundance microRNAs in the heart by performing intra-tissue comparisons and the preeminent position of miR-1 within muscle tissue has been quantitatively established. In addition, the requirement of the microRNA pathway in cardiac muscle maintenance has been unequivocally established (as its absence is lethal). Based on the results described, we suggest two distinct, but related, mechanisms to explain the drastic loss of cardiac function. The first one implicates fast skeletal muscle gene expression as a plausible causative factor in loss of cardiac function. Second, the loss microRNA function maybe causing cardiac heterochrony, which ultimately leads to heart failure. In addition, our data from the deep sequencing suggests that loss of a few microRNAs-including miR-1 and miR-133a may ultimately be responsible for the dramatic loss of function seen in *Dgcr8* deficient cardiomyocytes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Ron Kahn (Joslin Diabetes Center) for the MCK-Cre line and members of the Lodish and Bartel lab for their insightful comments. We are indebted to Carsten Russ and the Broad Sequencing Platform for carrying out Illumina/Solexa sequencing runs at the Broad Institute.

FUNDING SOURCES This work was supported by the following grants: **PKR** (Muscular Dystrophy Association-3882), **RB** (NIH K08 NS48118-01, NIH RO1 NS057221, Stem Cell Research Foundation and the Pew Scholars Program in the Biomedical Sciences); **HFL** (NIH-R01 DK068348-04 and a SPARC grant from the Broad Institute); **MK** (NIH-HL52212); **HFL & MK** (NIH/NHLBI P01 - HL066105); **RJ** (NIH RO1-CA087869, NIH R37-CA084198, and NIH RO1-HD0445022); **RL** (NIH RO1s, HL071775, HL088533, HL090884, and HL093148); **MB** (Max Kade Foundation, Austria); **HRC** (NIH RO1 GM067031 to David Bartel).

Non-standard Abbreviations and Acronyms

Dgcr8	DiGeorge syndrome critical region 8
PAZ	Piwi Argonaut and Zwillie
shRNA	short hairpin RNA
PGC1 α	PPAR γ coactivator-1a
PGC1 β	PPAR γ coactivator-1b
Myh6	Myosin Heavy Chain 6
Myh7	Myosin Heavy Chain 7
RT-PCR	Reverse Transcription-Polymerase Chain Reaction
DCM	Dilated Cardiomyopathy
3' UTR	3' Untranslated Region
KO	knock-out

REFERENCES

1. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 2004;116:281–97. [PubMed: 14744438]
2. Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ. miRBase: tools for microRNA genomics. *Nucleic Acids Res* 2008;36:D154–8. [PubMed: 17991681]
3. Vasudevan S, Tong Y, Steitz JA. Switching from repression to activation: microRNAs can up-regulate translation. *Science* 2007;318:1931–4. [PubMed: 18048652]
4. Orom UA, Nielsen FC, Lund AH. MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation. *Mol Cell* 2008;30:460–71. [PubMed: 18498749]
5. Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB. Prediction of mammalian microRNA targets. *Cell* 2003;115:787–98. [PubMed: 14697198]
6. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 2005;120:15–20. [PubMed: 15652477]
7. Xiao C, Calado DP, Galler G, Thai TH, Patterson HC, Wang J, Rajewsky N, Bender TP, Rajewsky K. MiR-150 controls B cell differentiation by targeting the transcription factor c-Myb. *Cell* 2007;131:146–59. [PubMed: 17923094]
8. Zhao Y, Ransom JF, Li A, Vedantham V, von Drehle M, Muth AN, Tsuchihashi T, McManus MT, Schwartz RJ, Srivastava D. Dysregulation of cardiogenesis, cardiac conduction, and cell cycle in mice lacking miRNA-1-2. *Cell* 2007;129:303–17. [PubMed: 17397913]
9. Wang S, Aurora AB, Johnson BA, Qi X, McAnally J, Hill JA, Richardson JA, Bassel-Duby R, Olson EN. The endothelial-specific microRNA miR-126 governs vascular integrity and angiogenesis. *Dev Cell* 2008;15:261–71. [PubMed: 18694565]

10. van Rooij E, Sutherland LB, Qi X, Richardson JA, Hill J, Olson EN. Control of stress-dependent cardiac growth and gene expression by a microRNA. *Science* 2007;316:575–9. [PubMed: 17379774]
11. Thai TH, Calado DP, Casola S, Ansel KM, Xiao C, Xue Y, Murphy A, Frendewey D, Valenzuela D, Kutok JL, Schmidt-Supprian M, Rajewsky N, Yancopoulos G, Rao A, Rajewsky K. Regulation of the germinal center response by microRNA-155. *Science* 2007;316:604–8. [PubMed: 17463289]
12. Yi R, O'Carroll D, Pasolli HA, Zhang Z, Dietrich FS, Tarakhovsky A, Fuchs E. Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nat Genet* 2006;38:356–62. [PubMed: 16462742]
13. Chen JF, Murchison EP, Tang R, Callis TE, Tatsuguchi M, Deng Z, Rojas M, Hammond SM, Schneider MD, Selzman CH, Meissner G, Patterson C, Hannon GJ, Wang DZ. Targeted deletion of Dicer in the heart leads to dilated cardiomyopathy and heart failure. *Proc Natl Acad Sci U S A* 2008;105:2111–6. [PubMed: 18256189]
14. Finnegan EJ, Margis R, Waterhouse PM. Posttranscriptional gene silencing is not compromised in the Arabidopsis CARPEL FACTORY (DICER-LIKE1) mutant, a homolog of Dicer-1 from *Drosophila*. *Curr Biol* 2003;13:236–40. [PubMed: 12573220]
15. Cuellar TL, Davis TH, Nelson PT, Loeb GB, Harfe BD, Ullian E, McManus MT. Dicer loss in striatal neurons produces behavioral and neuroanatomical phenotypes in the absence of neurodegeneration. *Proc Natl Acad Sci U S A* 2008;105:5614–9. [PubMed: 18385371]
16. Kobayashi T, Lu J, Cobb BS, Rodda SJ, McMahon AP, Schipani E, Merckenschlager M, Kronenberg HM. Dicer-dependent pathways regulate chondrocyte proliferation and differentiation. *Proc Natl Acad Sci U S A* 2008;105:1949–54. [PubMed: 18238902]
17. Koralov SB, Muljo SA, Galler GR, Krek A, Chakraborty T, Kanellopoulou C, Jensen K, Cobb BS, Merckenschlager M, Rajewsky N, Rajewsky K. Dicer ablation affects antibody diversity and cell survival in the B lymphocyte lineage. *Cell* 2008;132:860–74. [PubMed: 18329371]
18. da Costa Martins PA, Bourajaj M, Gladka M, Kortland M, van Oort RJ, Pinto YM, Molkentin JD, De Windt LJ. Conditional Dicer Gene Deletion in the Postnatal Myocardium Provokes Spontaneous Cardiac Remodeling. *Circulation*. 2008
19. Rao M. Conserved and divergent paths that regulate self-renewal in mouse and human embryonic stem cells. *Dev Biol* 2004;275:269–86. [PubMed: 15501218]
20. Cai X, Hagedorn CH, Cullen BR. Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *Rna* 2004;10:1957–66. [PubMed: 15525708]
21. Lee Y, Kim M, Han J, Yeom KH, Lee S, Baek SH, Kim VN. MicroRNA genes are transcribed by RNA polymerase II. *Embo J* 2004;23:4051–60. [PubMed: 15372072]
22. Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Radmark O, Kim S, Kim VN. The nuclear RNase III Drosha initiates microRNA processing. *Nature* 2003;425:415–9. [PubMed: 14508493]
23. Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ. Processing of primary microRNAs by the Microprocessor complex. *Nature* 2004;432:231–5. [PubMed: 15531879]
24. Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R. The Microprocessor complex mediates the genesis of microRNAs. *Nature* 2004;432:235–40. [PubMed: 15531877]
25. Yi R, Qin Y, Macara IG, Cullen BR. Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev* 2003;17:3011–6. [PubMed: 14681208]
26. Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 2001;409:363–6. [PubMed: 11201747]
27. Babiarz JE, Ruby JG, Wang Y, Bartel DP, Blelloch R. Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs. *Genes Dev* 2008;22:2773–85. [PubMed: 18923076]
28. Okamura K, Hagen JW, Duan H, Tyler DM, Lai EC. The mirtron pathway generates microRNA-class regulatory RNAs in *Drosophila*. *Cell* 2007;130:89–100. [PubMed: 17599402]
29. Ruby JG, Jan CH, Bartel DP. Intronic microRNA precursors that bypass Drosha processing. *Nature* 2007;448:83–6. [PubMed: 17589500]

30. Grimson A, Srivastava M, Fahey B, Woodcroft BJ, Chiang HR, King N, Degnan BM, Rokhsar DS, Bartel DP. Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature*. 2008
31. Bruning JC, Michael MD, Winnay JN, Hayashi T, Horsch D, Accili D, Goodyear LJ, Kahn CR. A muscle-specific insulin receptor knockout exhibits features of the metabolic syndrome of NIDDM without altering glucose tolerance. *Mol Cell* 1998;2:559–69. [PubMed: 9844629]
32. van Rooij E, Sutherland LB, Thatcher JE, DiMaio JM, Naseem RH, Marshall WS, Hill JA, Olson EN. Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. *Proc Natl Acad Sci U S A* 2008;105:13027–32. [PubMed: 18723672]
33. Fish JE, Santoro MM, Morton SU, Yu S, Yeh RF, Wythe JD, Ivey KN, Bruneau BG, Stainier DY, Srivastava D. miR-126 regulates angiogenic signaling and vascular integrity. *Dev Cell* 2008;15:272–84. [PubMed: 18694566]
34. Rinn JL, Snyder M. Sexual dimorphism in mammalian gene expression. *Trends Genet* 2005;21:298–305. [PubMed: 15851067]
35. Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, Mills AA, Elledge SJ, Anderson KV, Hannon GJ. Dicer is essential for mouse development. *Nat Genet* 2003;35:215–7. [PubMed: 14528307]
36. Harfe BD, McManus MT, Mansfield JH, Hornstein E, Tabin CJ. The RNaseIII enzyme Dicer is required for morphogenesis but not patterning of the vertebrate limb. *Proc Natl Acad Sci U S A* 2005;102:10898–903. [PubMed: 16040801]
37. Schaefer A, O'Carroll D, Tan CL, Hillman D, Sugimori M, Llinas R, Greengard P. Cerebellar neurodegeneration in the absence of microRNAs. *J Exp Med* 2007;204:1553–8. [PubMed: 17606634]
38. Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, Conlon FL, Wang DZ. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat Genet* 2006;38:228–33. [PubMed: 16380711]
39. O'Rourke JR, Georges SA, Seay HR, Tapscott SJ, McManus MT, Goldhamer DJ, Swanson MS, Harfe BD. Essential role for Dicer during skeletal muscle development. *Dev Biol* 2007;311:359–68. [PubMed: 17936265]
40. Wang Y, Medvid R, Melton C, Jaenisch R, Blueloch R. DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nat Genet* 2007;39:380–5. [PubMed: 17259983]
41. Huang QQ, Feng HZ, Liu J, Du J, Stull LB, Moravec CS, Huang X, Jin JP. Co-expression of skeletal and cardiac troponin T decreases mouse cardiac function. *Am J Physiol Cell Physiol* 2008;294:C213–22. [PubMed: 17959729]
42. Hannenhalli S, Putt ME, Gilmore JM, Wang J, Parmacek MS, Epstein JA, Morrissey EE, Margulies KB, Cappola TP. Transcriptional genomics associates FOX transcription factors with human heart failure. *Circulation* 2006;114:1269–76. [PubMed: 16952980]
43. Rao PK, Kumar RM, Farkhondeh M, Baskerville S, Lodish HF. Myogenic factors that regulate expression of muscle-specific microRNAs. *Proc Natl Acad Sci U S A* 2006;103:8721–6. [PubMed: 16731620]
44. Liu N, Williams AH, Kim Y, McAnally J, Bezprozvannaya S, Sutherland LB, Richardson JA, Bassel-Duby R, Olson EN. An intragenic MEF2-dependent enhancer directs muscle-specific expression of microRNAs 1 and 133. *Proc Natl Acad Sci U S A* 2007;104:20844–9. [PubMed: 18093911]
45. Davis BN, Hilyard AC, Lagna G, Hata A. SMAD proteins control DROSHA-mediated microRNA maturation. *Nature* 2008;454:56–61. [PubMed: 18548003]
46. Heo I, Joo C, Cho J, Ha M, Han J, Kim VN. Lin28 mediates the terminal uridylation of let-7 precursor MicroRNA. *Mol Cell* 2008;32:276–84. [PubMed: 18951094]
47. Newman MA, Thomson JM, Hammond SM. Lin-28 interaction with the Let-7 precursor loop mediates regulated microRNA processing. *Rna* 2008;14:1539–49. [PubMed: 18566191]
48. Viswanathan SR, Daley GQ, Gregory RI. Selective blockade of microRNA processing by Lin28. *Science* 2008;320:97–100. [PubMed: 18292307]
49. Yi R, Pasolli HA, Landthaler M, Hafner M, Ojo T, Sheridan R, Sander C, O'Carroll D, Stoffel M, Tuschl T, Fuchs E. DGCR8-dependent microRNA biogenesis is essential for skin development. *Proc Natl Acad Sci U S A*. 2008

50. Wang Y, Baskerville S, Shenoy A, Babiarz JE, Baehner L, Blueloch R. Embryonic stem cell-specific microRNAs regulate the G1-S transition and promote rapid proliferation. *Nat Genet* 2008;40:1478–83. [PubMed: 18978791]
51. Ikeda S, Kong SW, Lu J, Bisping E, Zhang H, Allen PD, Golub TR, Pieske B, Pu WT. Altered microRNA expression in human heart disease. *Physiol Genomics* 2007;31:367–73. [PubMed: 17712037]
52. Liu N, Bezprozvannaya S, Williams AH, Qi X, Richardson JA, Bassel-Duby R, Olson EN. microRNA-133a regulates cardiomyocyte proliferation and suppresses smooth muscle gene expression in the heart. *Genes Dev* 2008;22:3242–54. [PubMed: 19015276]
53. Lee RC, Feinbaum RL, Ambros V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 1993;75:843–54. [PubMed: 8252621]
54. Pasquinelli AE, Reinhart BJ, Slack F, Martindale MQ, Kuroda MI, Maller B, Hayward DC, Ball EE, Degan B, Muller P, Spring J, Srinivasan A, Fishman M, Finnerty J, Corbo J, Levine M, Leahy P, Davidson E, Ruvkun G. Conservation of the sequence and temporal expression of *let-7* heterochronic regulatory RNA. *Nature* 2000;408:86–9. [PubMed: 11081512]
55. Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G. The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 2000;403:901–6. [PubMed: 10706289]
56. Decembrini S, Andreazzoli M, Barsacchi G, Cremisi F. Dicer inactivation causes heterochronic retinogenesis in *Xenopus laevis*. *Int J Dev Biol* 2008;52:1099–103. [PubMed: 18956342]
57. Caygill EE, Johnston LA. Temporal regulation of metamorphic processes in *Drosophila* by the *let-7* and *miR-125* heterochronic microRNAs. *Curr Biol* 2008;18:943–50. [PubMed: 18571409]
58. Sokol NS, Xu P, Jan YN, Ambros V. *Drosophila let-7* microRNA is required for remodeling of the neuromusculature during metamorphosis. *Genes Dev* 2008;22:1591–6. [PubMed: 18559475]
59. Huang X, Lee KJ, Riedel B, Zhang C, Lemanski LF, Walker JW. Thyroid hormone regulates slow skeletal troponin I gene inactivation in cardiac troponin I null mouse hearts. *J Mol Cell Cardiol* 2000;32:2221–8. [PubMed: 11112997]




Figure 1. microRNA abundance in the murine adult heart

(A) The top 20 known microRNAs (in terms of normalized read number) from the male (dark bars) and the female (clear bars) heart small libraries were converted to percentage terms and plotted. Since the rank order differs slightly between the male and female libraries, the total number of microRNAs plotted is greater than 20. Note the abundance of miR-1 reads relative to other known microRNAs. (B). microRNAs belonging to the same family (as defined by Targetscan; www.targetscan.org) were summed and plotted together. This analyses reveals that aside from miR-1, the let-7 and miR-30 families are amongst the ones that are highly abundant in the heart.






Fig.2. Lethality and microRNA expression in muscle-specific *dgcr8* KO mice


(A) Actual numbers of Cre positive mice (and the expected numbers-based on Punnett square analysis for two independent loci) obtained from matings between 2lox/+; Cre positive mice are shown. Expected numbers are based on the assumption that Cre transgene is heterozygous although this is not known. (B). Postnatal lethality of muscle-specific *dgcr8* KO mice. Survival curves for Cre positive mice are shown and reveal the lethality when *dgcr8* is excised in muscle tissue. Moribund, “hunched over” mice that had to be sacrificed due to animal care committee specifications were considered dead for survival analysis. Survival curves were plotted using a built-in module in Prism software. (C). miR-1, miR-133 and miR-208 expression was determined using Northern Blots from total RNA derived from heart tissue. Tissues from three **mutant** (2lox/2lox; Cre positive) and three sex and age-matched **control** (2lox/+; Cre positive) siblings were used for total RNA isolation; ages were 29 days (males), 29 days and 38 days (females). The same blots were reprobed for U6 (bottom part of each set) to normalize for differences in loading. Quantified miR/U6 ratios are plotted below for miR-1, miR-133 and miR-208 and the indicated p values were obtained using two-sample (unequal variance) one-tailed t-tests. (D). A larger region of the miR-208 Northern blot is shown to reveal the absence of pre-miR-208 in total RNA derived from the mutant heart.

**Fig.3.**

(A). Intact, excised hearts from 30 day old mutant (2lox/2lox; Cre positive; left) and control (2lox/+; Cre positive; right) female sibling mice. (B). Representative long axis sections at different stages (as indicated) from mutant and control sex-matched sibling mice were stained with H & E (for end stage: left ventricle=LV and right ventricle=RV) or Masson's Trichrome (for d14/15 & d21/22). Bar = 500 μ M. The stage at which the mice were euthanized to reveal end-stage pathology was variable and defined by the health status of the mice and is d34 in this panel. (C). High magnification (20X) view of Masson's Trichrome stained sections from 3 week and end-stage (d43 in this panel) mutant and control female sibling mice (interstitial blue staining (bright green arrows) is indicative of fibrotic collagen deposits; Bars = 50 μ M. (D). RT-PCR assay to detect mature miR-1, miR-133a and miR-208 levels showing that the decline is already evident at two weeks after birth and continues to decrease by the time the mice are moribund ("end stage"). The ratio of mutant to control is shown on the y-axis and the pairs chosen for evaluation were age and sex matched.

**Fig.4.**

Trans-Thoracic Echocardiography. (A) Representative short axis B and M mode images for both mutant (left) and control (right) mice at three weeks of age showing dilation in the mutant mice. (B) Summary of echocardiographic data 3 weeks and 4 weeks after birth showing progressive dilation and reduction in ventricular function (see numbers for EDD and FS respectively) between three and four weeks in mutant mice. WT: Wall thickness, EDD: End-diastolic diameter, ESD: End-systolic diameter, FS: Fractional shortening, HR: Heart rate. *: $p < 0.05$ vs. 2lox/+; Cre pos; † $p < 0.05$ vs. 3 weeks.

**Fig.5.**

Gene expression patterns in *dgr8* KO heart. Total RNA was obtained from end stage (mutant) and sex and age-matched control mice. A minimum of seven pairs was used for the analysis of relative expression levels of the indicated genes. Box and whisker plots are represented for each gene. Expression is depicted as a ratio of mutant over control, (with each being first normalized to GAPDH to account for differences in total amount of RNA used). Lack of a difference should manifest itself as having a ratio of 1.0 (dotted line). While it is appreciated that some genes may fall into two or more categories during the embryonic and postnatal development, for convenience, they are grouped into a single category (as indicated) that is representative of adult gene expression.