Striatal Pre- and Postsynaptic Profile of Adenosine A_{2A} Receptor Antagonists

Marco Orru^{1,9}, Jana Bakešová^{2,9}, Marc Brugarolas² ´sar Quiroz¹, Vahri Beaumont³, Steven R. Goldberg¹, Carme Lluís², Antoni Cortés², Rafael Franco², Vicent Casadó², Enric I. Canela², Sergi Ferré^{1*}

1 National Institute on Drug Abuse, IRP, NIH, DHHS, Baltimore, Maryland, United States of America, 2 Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Faculty of Biology, University of Barcelona, Barcelona, Spain, 3 CHDI Management, CHDI Foundation, Los Angeles, California, United States of America

Abstract

Striatal adenosine A_{2A} receptors (A_{2A} Rs) are highly expressed in medium spiny neurons (MSNs) of the indirect efferent pathway, where they heteromerize with dopamine D_2 receptors (D_2 Rs). A_{2A} Rs are also localized presynaptically in corticostriatal glutamatergic terminals contacting MSNs of the direct efferent pathway, where they heteromerize with adenosine A_1 receptors (A_1 Rs). It has been hypothesized that postsynaptic A_{2A} R antagonists should be useful in Parkinson's disease, while presynaptic A_{2A} R antagonists could be beneficial in dyskinetic disorders, such as Huntington's disease, obsessivecompulsive disorders and drug addiction. The aim or this work was to determine whether selective A_{2A} R antagonists may be subdivided according to a preferential pre- *versus* postsynaptic mechanism of action. The potency at blocking the motor output and striatal glutamate release induced by cortical electrical stimulation and the potency at inducing locomotor activation were used as *in vivo* measures of pre- and postsynaptic activities, respectively. SCH-442416 and KW-6002 showed a significant preferential pre- and postsynaptic profile, respectively, while the other tested compounds (MSX-2, SCH-420814, ZM-241385 and SCH-58261) showed no clear preference. Radioligand-binding experiments were performed in cells expressing A_{2A} R heteromers. Heteromerization played a key role in the presynaptic profile of SCH-442416, since it bound with much less affinity to A_{2A} R when co-expressed with D_2 R than with A_1 R. KW-6002 showed the best relative affinity for A_{2A} R co-] TJETBT/F4 1 Tf8.96 involves $A_{2A}R$ and $D_{2}R$ that do not form heteromers, but most probably homomers [4]. In this interaction, which takes place at the level of adenylyl-cyclase (AC), stimulation of G_i-coupled D₂R counteracts the effects of Golf-coupled A2AR [4]. Due to a strong tonic effect of endogenous dopamine on striatal D₂R, this interaction keeps A_{2A}R from signaling through AC. However, under conditions of dopamine depletion or with blockade of D_2R , A_{2A}R-mediated AC activation is unleashed. This is biochemically associated with a significant increase in the phosphorylation of PKA-dependent substrates, which increases gene expression and the activity of the indirect MSN, producing locomotor depression (reviewed in ref. [4]). This interaction seems to be the main mechanism responsible for the locomotor depression induced by D₂R antagonists. Thus the motor depressant and most biochemical effects induced by genetic or pharmacologic blockade of D₂R are counteracted by the genetic or pharmacological blockade of $A_{2A}R$ [8–10]

Striatal A2ARs are not only localized postsynaptically but also presynaptically, in glutamatergic terminals, where they heteromerize with A1 receptors (A1Rs) and where their stimulation facilitates glutamatergic neurotransmission [5,11]. Interestingly, presynaptic $A_{2A}Rs$ are preferentially localized in glutamatergic terminals of cortico-striatal afferents to the direct MSN [5]. According to the widely accepted functional basal circuitry model [2,3], blockade of postsynaptic $A_{2A}R$ localized in the indirect MSN should produce motor activation (by potentiating D₂Rmediated effects by means of $A_{2A}R$ - D_2R receptor interactions). On the other hand, according to the same model, blockade of presynaptic A_{2A}R localized in the cortico-striatal glutamatergic terminals that make synaptic contact with the direct MSN should decrease motor activity (by inhibiting glutamate release). The preferential locomotor-activating effects of systemically administered $A_{2A}R$ receptor antagonists can be explained by a stronger influence of a tonic adenosine and A2AR receptor-mediated modulation of the indirect pathway versus the direct pathway under basal conditions. In any case, the potency at inducing locomotor activation can be used as an in vivo measure of the ability of an $A_{2A}R$ antagonist to block postsynaptic striatal $A_{2A}R$. Recently we have established an *in vivo* model that evaluates the efficacy of cortico-striatal glutamatergic neurotransmission to the direct MSN, by quantifying the correlation between the current delivered into the orofacial premotor cortex and the concomitant electromyographic response elicited in the jaw muscles [5]. In this model, $A_{2A}R$ or D_1R antagonists were able to counteract the motor output induced by cortical electrical stimulation, which can only be explained by blockade of striatal presynaptic A2AR or postsynaptic D_1R , respectively [5,12].

Receptor heteromer is defined as a macromolecular complex composed by at least two (functional) receptor units with biochemical properties that are demonstrably different from those of its individual components [13]. Specific ligand binding characteristics are one of those properties [13,14]. The aim of the present study was, first, to investigate the possible existence of different pre- and postsynaptic profiles of several A2AR antagonists. The potency at blocking the motor output and striatal glutamate release induced by cortical electrical stimulation and the potency at inducing locomotor activation were used as in vivo measures of pre- and postsynaptic activities, respectively. Second, we wanted to evaluate if the different pre- and postsynaptic profiles could be related to different affinities that A2AR could have for those compounds when forming heteromers with either A1R or D_2R . In fact, the results strongly suggest that heteromerization plays a key role in the pre- and postsynaptic profile of A_{2A}R antagonists.

Materials and Methods

Ethics Statement

All animals used in the study were handled in accordance with the National Institutes of Health Animal care guidelines. The animal research conducted to perform this study was approved by the NIDA IRP Animal Care and Use Committee (under the auspices of protocol 09-BNRB-73) on 12/7/2009.

Animals

Male Sprague-Dawley rats (Charles River Laboratories, Wilmington, MA) weighting between 300–350 g were used in these experiments. Rats were housed 2 per cage and they maintained at a temperature of $22\pm2^{\circ}$ C on a regular 12-h light–dark cycle. Food and water were available *ad libitum*.

Adenosine A_{2A}R antagonists

The following A_{2A}R antagonists were used: 2-(2-Furanyl)-7-[3-(4methoxyphenyl)propyl]-7H-pyrazolo[4,3-e][1,2,4]triazolo[1,5-c]pyrimidin-5-amine (SCH-442416), 2-(2-Furanyl)-7-(2-phenylethyl)-7Hpyrazolo[4,3-e][1,2,4]triazolo[1,5-c]pyrimidin-5-amine (SCH-58261), 2-(2-furanyl)-7-[2-[4-[4-(2-methoxyethoxy)phenyl]-1-piperazinyl]ethyl]-7H-pyrazolo[4,3-e][1,2,4]-triazolo[1,5-c]pyrimidin-5-amine (SCH-420814), 4-(2-[7-Amino-2-(2-furyl)[1,2,4]triazolo[2,3-a][1,3,5] triazin-5-ylamino]ethyl)phenol (ZM-241385), (E)-1, 3-diethyl-8-(3,4dimethoxystyryl)-7-methyl-3,7-dihydro-1H-purine-2,6-dione (**KW-**6002), (E)-3-(3-hydroxypropyl)-8-[2-(3-methoxyphenyl)vinyl]-7-methyl-1-prop-2-ynyl-3,7-dihydropurine-2,6-dione (MSX-2) and its watersoluble phosphate prodrug (E)-phosphoric acid mono-(3-{8-[2-(3methoxyphenyl)vinyl]-7-methyl-2,6-dioxo-1-prop-2-ynyl-1,2,6,7-tetrahydropurin-3-yl}propyl) ester disodium salt (MSX-3). MSX-3 is a water-soluble phosphate pro-drug of MSX-2; in vivo MSX-3 is readily converted to the A2AR antagonist MSX-2 (Sauer et al., 2002). For their systemic administration, the compounds were prepared as follows:

SCH-442416 and SCH 58261 were suspended in a solution of 5% dimethyl- sulfoxide (DMSO) (Sigma-Aldrich, St. Louis,MI), 5% TWEEN80 (Sigma-Aldrich, St. Louis, MI) and 90% ddH₂O; SCH-420814 was suspended in a solution of 20% PEG400, 40% β -cyclodextrin and 40% Lutrol 1% (in ddH₂O); ZM-241385 was suspended in a solution of 15% DMSO, 10% TWEEN80 and 75% ddH₂O; KW-6002 was suspended in a solution of 8% TWEEN80 and 92% ddH₂O; MSX-3 was dissolved in sterile saline (with 3 μ

Surgical procedures

Rats were anesthetized with 3 ml/kg of Equithesin (4.44 g of chloral hydrate, 0.972 g of Na pentobarbital, 2.124 g of MgSO₄, 44.4 ml of propylene glycol, 12 ml of ethanol and distilled H_2O up to 100 ml of final solution; NIDA Pharmacy, Baltimore, MD) and implanted unilaterally with bipolar stainless steel electrodes, 0.15 mm in diameter (Plastics One, Roanoke, VA), into the orofacial area of the lateral agranular motor cortex (3 mm anterior, 3 and 4 mm lateral, and 4.2 mm below bregma). The electrodes and a head holder (connected to a swivel during stimulation) were fixed on the skull with stainless steel screws and dental acrylic resin. For the experiments with electromyographic (EMG) recording, electrodes were also implanted in mastication muscles (during the same surgical procedure). Two 5 mm-long incisions were made in the skin on the upper and lower jaw areas to expose the masseter and the lateral pterygoid muscles. Two silicon rubber-coated coiled stainless steel recording electrodes (Plastics One, Roanoke, VA) were slipped below the skin from the incision in the skull until the tips showed up from the incisions in the jaw. The bare tips of the electrodes were then held in contact with the masseter and the lateral pterygoid muscles and the skin was closed with surgical staples. The other end of the recording electrodes was encased in a molded plastic pedestal with a round threaded post which was attached to an electrical swivel and then to a differential amplifier (Grass LP511, Grass Instruments, Warwick, RI). The pedestal was secured to the skull with dental cement together with the stimulation electrodes. For the in vivo microdialysis experiments, concentric microdialysis probes with 2mm long dialysis membranes (Eicom Corp., Tokio, Japan) were implanted respectively into the striatum ipsilateral to the stimulation electrodes (0.0 mm AP, 4.5 ML and 7.0 mm DV).

EMG recording and power correlation analysis

Rats were placed in individual bowl chambers. Both stimulation electrodes and recording electrodes were attached using flexible shielded cabling to a four channel electrical swivel. Stimulation electrodes were connected to two-coupled constant current isolation units (PSIU6X, Grass Instruments West Warwick, RI) driven by an electrical stimulator (Grass S88X; Grass Instruments). The recording electrodes were connected to a differential amplifier (Grass LP511, West Warwick, RI). This configuration allows the rat to move freely while the stimulation and EMG recordings are taking place. After 60 min of habituation, biphasic current pulse trains (pulse of 0.1 ms at 120-200 µA; 100 Hz, 160 ms trains repeating once per 2 seconds) were delivered. The current intensity was adjusted to the threshold level, defined as the minimal level of current intensity allowing at least 95% of the stimulation pulses to elicit a positive EMG response. Positive EMG response was defined as at least 100% increase of the peak to peak amplitude respect to the background tonic EMG activity lasting more than 100 ms or at least 70% increase in the power of the EMG signal respect to the baseline. Positive EMG responses always matched observable small jaw movements. The threshold level was different for each animal but it was very stable and reproducible once established. The threshold level was in the 100 to 150 μ A range for most cases and it reached 200 μ A in a few (6) animals. Animals that failed to show a positive EMG response with electrical cortical stimulation intensities of 200 µA were discarded from the experimental procedure (less than 10%). Both stimulator monitoring and the amplified and filtered EMG signal (20,000 times gain, bandwidth from 10 to 1,000 Hz with a notch filter set at 60 Hz) were directed to analogto-digital converter for recording (Lab-Trax-4, World Precision Instruments, Sarasota, FL) and backup (NI 9215, National Instruments, Austin, TX) and digitized at a sampling rate of

10,000 samples/second. Recordings of the digitized data were made using the software Data Trax2 software (World Precision Instruments) and LabVIEW SignalExpress (National Instruments). A power correlation analysis was used to quantify the correlation between the stimulation pulses of current delivered into the orofacial motor cortex (input signal; $\mu A)$ and the elicited EMG response in the jaw muscles (output signal; μV). Decrease in the power correlation coefficient (PCC) between these two signals is meant to describe a decrease in the efficacy of the transmission in the neural circuit. Off-line, both signals were rectified and the root mean square (RMS) over each period of the stimulation pulses was calculated in the recorded signals using Data Trax2 software. The transformed data (RMS) from the stimulator monitor and the EMG were then exported with a time resolution of 100 samples/second to a spreadsheet file. The stimulation signal values were used as a reference to select data in a time window of 320 ms starting at the beginning of each train of pulses. This time window was chosen to ensure the analysis of any EMG response whose occurrence or length was delayed from the onset of the stimulation trains and to maximize the exclusion from the analysis of spontaneous jaw movements not associated with the stimulation. Pearson's correlation between the RMS values from the stimulation and EMG signals was then calculated for each experimental subject. PCC was calculated using the data recorded 40 min after the administration of the dose of any compound or vehicle. The effects of the different doses of A2AR antagonists on PCC were analyzed by a one-way ANOVA, followed by Dunnett's post-hoc test.

In vivo microdialysis

The experiments were performed on freely moving rats 24 h after probe implantation. An artificial cerebrospinal solution of (in mM) 144 NaCl, 4.8 KCl, 1.7 CaCl₂, and 1.2 MgCl₂ was pumped through the microdialysis probe at a constant rate of 1 µl/min. After a washout period of 90 min, dialysate samples were collected at 20-min intervals. After 60 min of collecting samples for baseline, the rats were injected either with the $A_{2A}R$ antagonists KW-6002 or SCH-442416. Both compounds were compared to vehicle controls (5% DMSO, 5% of TWEEN80 and 90% of ddH_2O). After 20 min from drug or vehicle injection, electrical stimulation pulses were applied through the electrodes implanted in the orofacial motor cortex for 20 min (pulse of 0.1 ms at 50-150 µA; 100 Hz, 160 ms trains repeating once \times second) and samples were collected for 2 additional hours. Glutamate content was measured by reverse-phase HPLC coupled to a flourimetric detector (Shimadzu Inc., Tokio, Japan) [15]. Glutamate values were transformed as percentage of the mean of the three values before the drug or vehicle injection and transformed values were statistically analyzed. The effect of KW-6002, SCH-442416 and vehicle were analyzed using a one-way ANOVA for repeated measures followed by a Tukey's post-hoc test.

Cell clones

To obtain CHO cells expressing single receptors or coexpressing $A_{2A}R$ and A_1R or $A_{2A}R$ and D_2R , the human cDNAs for A_1R or D_2 1000 µg/ml hygromycin). After an appropriate number of days/ passes, several stable lines were selected and cultured in the presence of the selection antibiotic (600 µg/ml geneticin or 300 µg/ml hygromycin). To obtain clones co-expressing $A_{2A}R$ and A_1R or $A_{2A}R$ and D_2R , CHO cells expressing high affinity $A_{2A}R$ (obtained as above described) were transfected with the human cDNAs for A_1R or D_2R cloned in pcDNA3.1 vector using lipofectamine. After an appropriate number of days/passes stable lines were selected and cultured in the presence of the selection antibiotic. The receptor(s) expression in the cell clones was first detected by dot-blot of cell lysates using commercial available antibodies and wild-type CHO cells lysates as negative basal staining. Positively moderated stained clones were grown to obtain membranes in which the receptor expression was quantified by radioligand-binding experiments (see Results).

Bioluminescence Resonance Energy Transfer (BRET) assays

The fusion proteins A2AR-Renilla Luciferase (A2AR-RLuc), A1R-Yellow Fluorescence Protein (A1R-YFP) and D2R-YFP were prepared and characterized as described elsewhere [16]. The cDNA encoding serotonin 5HT2B-YFP receptor was kindly provided by Dr. Irma Nardi (University of Pisa, Italy). CHO cells were transiently transfected with the corresponding fusion protein cDNA (see Figure legends) using lipofectamine. Cells were incubated (4 h) with the corresponding cDNA together with lipofectamine and Opti-MEM medium (Invitrogen). After 4 hours, the medium was changed to a fresh complete culture medium. Twenty-four hours after transfection, cells were washed twice in quick succession in HBSS with 10 mM glucose and scraped in 0.5 ml of the same buffer. To control the cell number, sample protein concentration was determined using a Bradford assay kit (Bio-Rad, Munich, Germany) using bovine serum albumin dilutions as standards. To quantify fluorescence proteins, cells (20 µg protein) were distributed in 96-well microplates (black plates with a transparent bottom) and fluorescence was read at 400 nm in a Fluo Star Optima Fluorimeter (BMG Labtechnologies, Offenburg, Germany) equipped with a high-energy xenon flash lamp, using a 10 nm bandwidth excitation filter. Receptorfluorescence expression was determined as fluorescence of the sample minus the fluorescence of cells expressing protein-Rluc alone. For BRET measurements, the equivalent of 20 µg of cell protein were distributed in 96-well microplates (Corning 3600, white plates; Sigma) and 5 µM coelenterazine H (Molecular Probes, Eugene, OR) was added. After 1 minute of adding coelenterazine H, the readings were collected using a Mithras LB 940, which allows the integration of the signals detected in the 485 nm-short- (440-500 nm) and the 530 nm-long-(510-590 nm) wavelength filters. To quantify receptor-Rluc expression luminescence readings were performed after 10 minutes of adding 5 µM coelenterazine H. The net BRET is defined as [(long-wavelength emission)/(short-wavelength emission)]-Cf where Cf corresponds to [(long-wavelength emission)/(short-wavelength emission)] for the Rluc construct expressed alone in the same experiment.

Radioligand binding experiments

Cells were disrupted with a Polytron homogenizer (PTA 20 TS rotor, setting 3; Kinematica, Basel, Switzerland) for two 5 s-periods in 10 volumes of 50 mM Tris-HCl buffer, pH 7.4 containing a proteinase inhibitor cocktail (Sigma, St. Louis, MO, USA). Cell debris was removed by centrifugation at 1,500 g for 5 min at 4°C and membranes were obtained by centrifugation at 105,000 g (40 min, 4°C). Membranes were resuspended and centrifuged under the same conditions. The pellet was stored at -20° C, washed

once more as described above and resuspended in 50 mM Tris-HCl buffer for immediate use. Membrane protein was quantified by the bicinchoninic acid method (Pierce Chemical Co., Rockford, IL, USA) using bovine serum albumin dilutions as standard. For competition experiments, membrane suspensions (0.2 mg of protein/ml) were incubated for 2 h at 25°C in 50 mM Tris-HCl buffer, pH 7.4, containing 10 mM MgCl₂ and 0.2 U/ml of adenosine deaminase (ADA, EC 3.5.4.4; Roche, Basel, Switzerland) with the indicated free concentration of the A_1R , $A_{2A}R$, or D_2R antagonist [³H]DPCPX (GE Healthcare, UK), [³H]ZM-241385, or ³H]YM-09151-2, respectively (NEN Perkin Elmer, Wellesley, MA, USA) or the A₁Ragonist [³H](R)-PIA (Moravek Biochemicals Inc., Brea, CA, USA) and increasing concentrations of DPCPX, ZM-241385, YM-09151-2, the A2AR agonist CGS-21680 or the tested A2AR antagonist (all provided by CHDI Foundation Inc.). Nonspecific binding was determined in the presence of $11 \,\mu\text{M}$ of the corresponding non-radiolabelled ligand. Free and membranebound ligand were separated by rapid filtration of 500 µl aliquots in a cell harvester (Brandel, Gaithersburg, MD, USA) through Whatman GF/C filters embedded in 0.3% polyethylenimine that were subsequently washed for 5 s with 5 ml of ice-cold 50 mM Tris-HCl buffer. The filters were incubated with 10 ml of Ecoscint H scintillation cocktail (National Diagnostics, Atlanta, GA, USA) overnight at room temperature and radioactivity counts were determined using a Tri-Carb 1600 scintillation counter (PerkinElmer, Boston, MA, USA) with an efficiency of 62% [17]. All displacers were dissolved in DMSO and diluted in the binding medium. The DMSO concentration in the binding incubates was less than 0.5% and, at this concentration, it did not affect agonist or antagonist affinity for their respective receptors.

Binding data analysis

Radioligand competition curves were analyzed by nonlinear regression using the commercial Grafit curve-fitting software (Erithacus Software, Surrey, UK), by fitting the binding data to the mechanistic two-state dimer receptor model [18,19]. Since there is now abundant evidence for GPCR oligomerization, including A_1R , $A_{2A}R$ and D_2R [20–23] and the minimal functional unit of GPCRs in biological tissues seems to imply dimerization [23], this model considers a homodimer as the minimal structural unit of the receptor. Here, we also consider the possibility of a homodimer as the minimal structural unit of a receptor forming homomers or forming heteromers with another receptor. To calculate the macroscopic equilibrium dissociation constants the following equation for a competition binding experiment deduced previously [19,24] was considered:

$$\begin{array}{c} \sum_{\mathbf{r} \in \mathbf{1} : \mathbf{s}, \mathbf{s}} = \left\{ \mathbf{r} + \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{c} \mathbf{r} + \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{c} \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{r} + \mathbf{r} + \mathbf{r} \\ \mathbf{r} \\ \mathbf{r} + \mathbf{r} \\ \mathbf{r} \\ \mathbf{r} + \mathbf{r} \\ \mathbf{r} + \mathbf{r} \\ \mathbf{r} + \mathbf{r} \\ \mathbf{r} \\$$

where A represents free radioligand (the adenosine A_1R or $A_{2A}R$ or dopamine D_2R antagonist [³H]DPCPX, [³H]ZM-241385 or [³H]YM-09151-2, respectively or the A_1R agonist [³H](R)-PIA) concentration, R_T is the total amount of receptor dimers and K_{DA1} and K_{DA2} are the macroscopic equilibrium dissociation constants describing the binding of the first and the second radioligand molecule (A) to the dimeric receptor; B represents the assayed competing compound concentration, and K_{DB1} and K_{DB2} are, respectively, the macroscopic equilibrium dissociation constants for the binding of the first ligand molecule (B) to a dimer and for the binding of the second ligand molecule (B) to the semi-occupied dimer; K_{DAB} is the hybrid equilibrium radioligand/competitor dissociation constant, which is the dissociation constant of B binding to a receptor dimer semi-occupied by A.

When the radioligand A shows non-cooperative behaviour, eq. (1) can be simplified to eq. (2) due to the fact that $K_{DA2} = 4K_{DA1}$ [19,25] and, therefore, K_{DA1} is enough to characterize the binding of the radioligand A:

$$+ C_{\mathbf{L}} \cdot \sqrt{C_{\mathbf{L}} \cdot C_{\mathbf{L}}} + C_{\mathbf{L}} \cdot \sqrt{C_{\mathbf{L}} \cdot C_$$

Binding to GPCRs quite often displays negative cooperativity. Under these circumstances $K_{D2}/K_{D1}>4$ and then K_{D1} and K_{D2} represent the "high-affinity" and the "low-affinity" binding sites, respectively. On the other hand, for positive cooperativity, $K_{D2}/K_{D1}<4$ and then K_{D2} represents the "high-affinity" and K_{D1} represents the "low-affinity" binding sites [25]. The two-state dimer model also introduces a cooperativity index (D_{CB}). The dimer cooperativity index for the competing ligand B is calculated as [19,25]:

$$\mathbf{b} = \mathbf{f} \left(\mathbf{C} \mathbf{b} \cdot \mathbf{A} \right)$$

The way the index is defined is such that its value is "0" for noncooperative binding, positive values of D_C indicate positive cooperativity, whereas negative values imply negative cooperativity [14,19].

In experimental conditions when both the radioligand A and the competitor B (i.e., most adenosine A_{2A} receptor antagonist tested in the present study) show non-cooperativity, it results that $K_{DA2} = 4K_{DA1}$ and $K_{DB2} = 4K_{DB1}$, and eq. (1) can be simplified to:

$$\begin{array}{cccc} & +^{\mathbf{C}\,\mathbf{t}} & \bullet & \sqrt{\mathbf{C}\,\mathbf{t}} & \bullet & \gamma^{\mathbf{t}} & \bullet &$$

When both the radioligand A and the competitor B (DPCPX, ZM241385, SCH 23390 or YM-09151-2) are the same compound and the binding is non-cooperative, eq. (3) simplifies to:

Goodness of fit was tested according to reduced χ^2 value given by the nonlinear regression program. The test of significance for two different population variances was based upon the Fdistribution (see ref. [25] for details). Using this *F* test, a probability greater than 95% (p<0.05) was considered the criterion to select a more complex equation to fit binding data over the simplest one. In all cases, a probability of less than 70% (p>0.30) resulted when one equation to fit binding data was not significantly better than the other. Results are given as parameter values \pm S.E.M. of three-four independent experiments.

Results

Striatal pre- versus postsynaptic profile of A_{2A} receptor antagonists

Dose-response experiments with the six $A_{2A}R$ antagonists indicated that four compounds (SCH-420814, SCH-58261, MSX-3 and ZM-241385) had a similar potency (similar minimal significant effective doses) at inducing locomotor activation (Fig. 1) and at reducing PCC (Fig. 2). The other two compounds had a very different profile: KW-6002 produced a strong locomotor activation already at the dose of 0.3 mg/kg i.p., while it did not reduce PCC at the highest tested dose (10 mg/kg i.p.). On the other hand, SCH-442416 produced a very weak locomotor activation, only significant at doses higher than 3 mg/kg i.p., while it significantly decreased PCC already at the dose of 0.1 mg/kg i.p.

In vivo microdialysis with cortical electrical stimulation was used as an additional *in vivo* evaluation of the preferential pre- and postsynaptic activity of SCH-442416 and KW-6002, respectively. SCH-442416 significantly counteracted striatal glutamate release induced by cortical stimulation at a dose that strongly reduced PCC but did not induce locomotor activation (1 mg/kg i.p.; Fig. 3). On the other hand, KW-6002 did not modify striatal glutamate release induced by cortical stimulation at a dose that produced a pronounced locomotor activation but did not reduce PCC (1 mg/ kg i.p.; Fig. 3).

Development of CHO cell-lines expressing A_1 - A_{2A} or A_{2A} - D_2 receptor heteromers

Cell clones expressing $A_{2A}R$, $A_1R-A_{2A}R$ heteromers or $A_{2A}R$ -D₂R heteromers and control clones expressing A_1R or D₂R were generated (see Materials and Methods). First of all, the ability of $A_{2A}R$ to form heteromers with A_1R or D₂R in CHO cells was demonstrated by BRET experiments in cells transiently coexpressing $A_{2A}R$ -Rluc and A_1R -YFP or $A_{2A}R$ -Rluc and D₂R-YFP. A positive BRET signal for energy transfer was obtained (Fig. 4). The BRET signal increased as a hyperbolic function of the concentration of the YFP-fusion construct added reaching an asymptote. As a negative control the BRET pair formed by $A_{2A}R$ -Rluc and 5-HT_{2B}R-YFP was used. As shown in Figure 4, the negative control gave a linear non-specific BRET signal. The significant and hyperbolic BRET signal found for these fusion proteins indicates that the intermolecular interaction between $A_{2A}R$ and A_1R or $A_{2A}R$ and D_2R in CHO cells is specific.

 $A_{2A}R$ - D_2R and A_1R - $A_{2A}R$ heteromerization in stably transfected CHO cells was shown by ligand binding experiments. This is an indirect approach for the identification of a receptor heteromer in native tissues or cells [13]. In the $A_{2A}R$ - D_2R heteromer, an allosteric interaction between both receptors in the heteromer has been described, in which the dopamine D_2R agonist affinity decreases in the presence of an $A_{2A}R$ agonist [14]. In CHO cells stably expressing $A_{2A}R$ and D_2R , the affinity of the D_2R for dopamine was determined by competition experiments of the D_2R antagonist [³H]YM-09151-2 *versus* dopamine in the presence (Fig. 5a) or in the absence (Fig. 5b) of the $A_{2A}R$ agonist



Figure 1. Locomotor activation in rats induced by $A_{2A}R$ **antagonists.** Data represent means \pm S.E.M. of the locomotor activity (distance traveled, in cm, of total accumulated counts) in habituated rats (90 min) during 90 min following the drug administration (n = 6–8 per group). * and **: p<0.05 and p<0.01, respectively in comparison to vehicle-treated animals (0 mg/kg); ANOVA with *post-hoc* Newman–Keuls' comparisons, p<0.5 and p<0.01, respectively). doi:10.1371/journal.pone.0016088.g001

CGS-21680 (200 nM). By fitting data obtained in the absence of CGS-21680 to eq. 3 (Methods; considering K_{DA1} = 2.9 nM see below) the calculated K_{DB1} was 9±2 µM. In the presence of CGS-

21680, 5 μ M of dopamine was unable to decrease the radioligand bound and more than 50% of radioligand bound was found in the presence of 100 μ M of dopamine (Fig. 5b). A K_{DB1} >30 μ M was



Figure 2. Blockade by $A_{2A}R$ **antagonists of the motor output induced by cortical electrical stimulation.** Dose-dependent decrease in the Power Correlation Coefficient (PCC) induced by the administration of different $A_{2A}R$ antagonists. Results represent means \pm S.E.M. (n = 5-6 per group). * and **: p<0.05 and p<0.01, respectively in comparison to vehicle-treated animals (0 mg/kg); ANOVA with *post-hoc* Dunnett' comparisons, p<0.5 and p<0.01, respectively). doi:10.1371/journal.pone.0016088.g002

). PLoS ONE | www.plosone.org

estimated and it was shown that CGS-21680 induced a decrease in the dopamine affinity for D₂R. An allosteric interaction in the A1R-A2AR heteromer has also been described, in which the A1R agonist affinity decreases in the presence of an $A_{2A}R$ agonist [11]. As shown in Figure 6a, the displacement of the A1R agonist $[^{3}H]$ R-PIA by CGS21680 was significantly (p < 0.001) better fitted by a biphasic than by a monophasic curve. At low CGS-21680 concentrations, when it binds preferentially to A2AR (at concentrations of CGS-21680 <500 nM, the direct binding of CGS-21680 to A_1R is <1%, according to the calculated affinity of A_1R for CGS-21680), CGS-21680 decreased the binding of [3H]R-PIA to the A_1R with an IC_{50} value of $386\pm35~nM$ (n = 3). At high CGS-21680 concentrations (>10 μ M), the [³H]R-PIA binding displacement reflects the binding of CGS-21680 directly to the A₁R and the competition between CGS-21680 and R-PIA for the binding to the A_1R . In fact, in the control clone expressing only A_1R , the displacement by CGS-21680 of $[^3H]R$ -PIA only occurred at CGS-21680 concentrations higher than 10 µM (Fig. 6b).

A pharmacological characterization of selected cell clones was performed with competition experiments of radio-labeled antagonists of A₁, A_{2A} and D₂ receptors *versus* selective agonists or antagonists. In all cases, the competition curves of the A_{2A}R antagonist [³H]ZM-241385 (2 nM) *versus* ZM-241385 (0.1 nM to 11 μ M), the D₂R antagonist [³H]YM-09151-2 (0.2 nM) *versus* YM-09151-2 (0.01 nM to 11 μ M) or the A₁R antagonist [³H]DPCPX (2 nM) *versus* DPCPX (0.1 nM to 11 μ M), were monophasic, indicating the absence of cooperativity (see Materials and Methods). By fitting the binding data to eq. 4 (Materials and Methods), the K_D (K_{D1}) values obtained for the antagonists ZM-241385 or YM-09151-2 were 8±3 nM and 2.9±0.3 nM, respectively, for the chosen A_{2A}R-D₂R clone, the K_D values obtained for the A₁R and A_{2A}R antagonists were 8±2 nM (DPCPX) and 1.8±0.4 nM (ZM-241385), respectively, for the

chosen A1R-A2A R697 730[[sn693r7394/663 0.7(t89Trsus)Tj/F6 0 5.97607s]Tj/F6 0 a1697 TDecti2A 27bo-3189(112.18.9663)-587.3(chosen)-585.7(A)]T





Figure 4. Identification of receptor heteromers in CHO cells by BRET saturation curve. BRET experiments were performed with CHO cells co-expressing A2AR-RLuc and A1R-YFP (A) or A2AR-RLuc and D2R-YEP (B). Co-transfections were performed with increasing amounts of plasmid–YFP (0.25 to 4 μg cDNA corresponding to A1R-YFP and 0.5 to 8 µg corresponding to D_2R -YFP) whereas the $A_{2A}R$ -*RLuc* construct was maintained constant (0.5 µg cDNA). Both fluorescence and luminiscence of each sample were measured before every experiment to confirm similar donor expressions (about 100,000 luminescent units) while monitoring the increase acceptor expression (10,000-25,000 fluorescent units). As a negative control, linear BRET was obtained in cells expressing equivalent luminescence and fluorescence amounts corresponding to A_{2A}R-RLuc, (0.5 µg transfected cDNA) and serotonin $5HT_{2B}$ -YFP (0.5 to 8 µg transfected cDNA) receptors. The relative amount of acceptor is given as the ratio between the fluorescence of the acceptor minus the fluorescence value of cells expressing the donor alone (YFP) and the luciferase activity of the donor (Rluc). BRET data are expressed as means \pm S.D. of 4–6 different experiments grouped as a function of the amount of BRET acceptor. doi:10.1371/journal.pone.0016088.g004

fitting to eq. 3). Since the screened compounds are $A_{2A}R$ antagonists, competition curves were expected to be monophasic, assuming that antagonist binding is not cooperative. In fact, in all cell clones, MSX-2, KW-6002, SCH-420814, ZM-241385 and SCH-58261 gave monophasic competition curves (fitting binding data to eq. 2 was not better than fitting to eq. 3; see Methods and Fig. 7 a–c as an example). Accordingly, the pharmacological characterization for these compounds gave $D_{CB}=0$ and $K_{DB2}=4K_{DB1}$ (see Table 3). For all compounds, co-transfection with A_1R did not significantly modify their affinity for $A_{2A}R$. On the other hand, co-transfection with D_2R significantly reduced the affinity of $A_{2A}R$ for MSX-2, SCH-420814, SCH-58261 and ZM-241385, from two to about nine times, and did not significantly modify the affinity of $A_{2A}R$ for KW-6002 (Table 3).



Figure 5. Allosteric interaction between A_{2A}R and D₂R in A_{2A}R-D₂R CHO cells. Competition experiments were performed in membrane preparations from CHO cells expressing A_{2A}R and D₂R with 0.5 nM [³H]YM-09151-2 and increasing concentrations of dopamine (from 0.1 nM to 30 μ M) in the absence (a) or in the presence (b) of 200 nM CGS-21680 as indicated in Methods. Data represent means \pm S.E.M. of a representative experiment performed with triplicates. doi:10.1371/journal.pone.0016088.q005

For SCH-442416, a careful statistically-based analysis of the monophasic or biphasic nature of the competition curves led to an unexpected finding: in A2AR-D2R cells, competition curves of [³H]ZM-241385 (2 nM) binding versus increasing concentrations of SCH-442416 were biphasic (fitting to eq. 2 improves the fitting to eq. 3; see Methods) (Fig. 7d). Table 4 shows the deduced pharmacological parameters from competition experiments of ³H]ZM-241385 versus SCH-442416 in cells expressing A_{2A}R, $A_1R\text{-}A_{2A}R$ and $A_{2A}R\text{-}D_2R.$ In $A_{2A}R$ and $A_1R\text{-}A_{2A}R$ cells the curves were monophasic. Accordingly, the pharmacological characterization gave a D_{CB} values of 0 and a $K_{DB2} = 4K_{DB1}$. In contrast, as mentioned above, in cells expressing $A_{2A}R$ - D_2R , competition curves were biphasic, and binding data were then fitted to eq. 2 (Methods) and robust parameters were obtained (Table 4). Thus, in $A_{2A}R$ - D_2R cells, SCH-442416 binding showed a strong negative cooperativity and, consequently, with a marked loss of affinity (an increase of 600 times in K_{DB2}) respect to cells expressing $A_{2A}R$. This is reflected by the B_{50} value (concentration competing 50% of radioligand binding), which was more than 40 times higher in $A_{2A}R$ - D_2R cells than in A_1R - $A_{2A}R$ cells or $A_{2A}R$ cells.

juuse Ziebui (IVI)



Figure 6. Allosteric interaction between A₁R and A_{2A}R in A₁R-A_{2A}R CHO cells. Competition experiments were performed in membrane preparations from CHO cells expressing A₁R or A₁R and A_{2A}R with 12 nM [³H]R-PIA *versus* increasing concentrations of the A_{2A}R agonist CGS-21680 as indicated in Methods. Data represent means \pm S.E.M. of a representative experiment performed with triplicates. doi:10.1371/journal.pone.0016088.g006

Discussion

An important finding of the present study is that several A_{2A}R antagonists previously thought as being pharmacologically similar present different striatal pre- and postsynaptic profiles. Six compounds already known as selective A2AR antagonists were first screened for their ability to block striatal pre- and postsynaptic A_{2A}Rs with *in vivo* models. Locomotor activation was used to evaluate postsynaptic activity while PCC reduction was used to determine presynaptic activity (see Introduction). Two compounds, SCH-442416 and KW-6002, showed preferential preand postsynaptic profiles, respectively, and four compounds, MSX-3, SCH-420814, SCH-58261 and ZM-241385, showed mixed pre-postsynaptic profiles. Combining in vivo microdialysis with cortical electrical stimulation was used as an additional in vivo evaluation of presynaptic activity of SCH-442416 and KW-6002. In agreement with its preferential presynaptic profile, SCH-442416 significantly counteracted striatal glutamate release induced by cortical stimulation at a dose (1 mg/kg i.p.) that strongly reduced PCC but did not induce locomotor activation. On the other hand, according to its preferential postsynaptic profile, KW-6002 did not modify striatal glutamate release induced by cortical stimulation at a dose (1 mg/kg i.p.) that produced a pronounced locomotor activation but did not

Table 1. Pharmacological parameters for agonist binding to A_1R , $A_{2A}R$ and D_2R in A_1R , $A_{2A}R$ and D_2R CHO cells.

Parameters	A _{2A} R cells	A ₁ R cells	D ₂ R cells	
K _{DB1}	90±30 nM	13±3 nM	120±60 nM	
K _{DB2}	360±120 nM	1±0.3 mM	480±240 nM	
D _{CB}	0	-1.3	0	
B ₅₀	180±60 nM	110±30 nM	240±120 nM	

Binding data from competition experiments were fitted assuming that receptors form homodimers, and cooperativity ($D_{CB} \neq 0$, fitting to eq. 2; Materials and Methods) or non-cooperativity ($D_{CB} = 0$, fitting to eq. 3; Materials and Methods) in competitor ligand binding was statistically tested (F test). K_{DB1} and K_{DB2} are, respectively, the equilibrium dissociation constants of the first and second binding of B (the A₁R, A₂A_R, or D₂R agonists: R-PIA, CGS-21680 or quinpirole, respectively) to the dimer. D_{CB} is the "dimer cooperativity" index for the binding of the ligand B, and B₅₀ is the concentration providing half saturation for B. Data are mean \pm S.E.M. values of three experiments. doi:10.1371/journal.pone.0016088.t001

counteract PCC. In a previous study, we reported that intrastriatal perfusion of MSX-3 almost completely counteracted striatal glutamate release induced by cortical electrical stimulation [5], which agrees with its very effective reduction of PCC shown in the present study.

Another important finding of the present study is that at least part of these pharmacological differences between $A_{2A}R$ antagonists can be explained by the ability of pre- and postsynaptic $A_{2A}R$ to form different receptor heteromers, with A_1R and D_2R , respectively [4–6,11,14]. Radioligand-binding experiments were performed in CHO cells stably expressing $A_{2A}R$, $A_{2A}R$ - D_2R heteromers or A_1R - $A_{2A}R$ heteromers to determine possible differences in the affinity of these compounds for different $A_{2A}R$ heteromers. Co-expression with A_1R did not significantly modify the affinity of $A_{2A}R$ for the different ligands, but co-expression with D_2R decreased the affinity of all compounds, with the exception of KW-6002. The structural changes in the $A_{2A}R$ induced by heteromerization with the D_2R could be detected not only by antagonists but also by agonists. Indeed, the affinity of the selective $A_{2A}R$ agonist CGS-21680 was reduced in cells co-

Table 2. Pharmacological parameters for agonist binding to A_1R - $A_{2A}R$ and $A_{2A}R$ - D_2R CHO cells.

Parameters	A _{2A} R-D ₂ R cells		A _{2A} R-A ₁ R cells	
	A _{2A} R	D ₂ R	A _{2A} R	A ₁ R
K _{DB1}	200±40 nM*	1.2±0.6 μM	70±10 nM	0.7±0.3 nM
K _{DB2}	$0.8{\pm}0.4~\mu\text{M}$	$4.8{\pm}2.4~\mu\text{M}$	280±40 nM	$1.1\!\pm\!0.5~\mu M$
D _{CB}	0	0	0	-2.6
B ₅₀	$0.4{\pm}0.08~\mu\text{M}$	$2.4{\pm}1.2~\mu\text{M}$	140±20 nM	30±10 nM

Binding data from competition experiments were fitted assuming that receptors (also when heteromerizing) form homodimers, and cooperativity ($D_{CB} \neq 0$, fitting to eq. 2; Materials and Methods) or non-cooperativity ($D_{CB} = 0$, fitting to eq. 3; Materials and Methods) in competitor ligand binding was statistically tested (F test). K_{DB1} and K_{DB2} are, respectively, the equilibrium dissociation constants of the first and second binding of B (the A₁R, A_{2A}R, or D₂R agonists: R-PIA, CGS-21680 or quinpirole, respectively) to the dimer. D_{CB} is the "dimer cooperativity" index for the binding of the ligand B, and B₅₀ is the concentration providing half saturation for B. Data are mean \pm S.E.M. values of three experiments.

*: $p{<}0.05$ compared to K_{DB1} values in $A_1R{-}A_{2A}R$ and $A_{2A}R$ cells (Table 1); one-way ANOVA, followed by Newman-Keuls test.

doi:10.1371/journal.pone.0016088.t002



Figure 7. Binding of the A_{2A}R antagonists KW-6002 and SCH-442416 to A₁R-A_{2A}R and A_{2A}R-D₂R CHO cells. Competition experiments of [³H]ZM-241385 (2 nM) *versus* increasing concentrations of KW-6002 (a and c) or SCH-442416 (b and d) were performed as indicated in Methods in membrane preparations from CHO cells expressing A₁R and A_{2A}R (a and b) or A_{2A}R and D₂R (c and d). Data are means \pm S.E.M. of a representative experiment performed with triplicates. doi:10.1371/journal.pone.0016088.g007

Table 3. Pharmacological parameters for A _{2A} R antagonist
binding to $A_{2A}R$, $A_1R-A_{2A}R$ and $A_{2A}R-D_2R$ CHO cells.

K _{D1} (nM)	A _{2A} R cells	A ₁ R-A _{2A} R cells	A _{2A} R-D ₂ R cells
ZM241385	0.9±0.3	1.8±0.4	8±3*
SCH58261	3.3±0.3	4.7±0.6	23±8*
MSX2	3.2±0.2	4.2±0.3	7±2*
KW6002	100±10	100±20	160±70
SCH420814	0.5±0.1	1.1±0.1	2.7±0.8*

Competition experiments of [³H]ZM-241385 (2 nM) binding versus increasing concentrations of A_{2A} receptor antagonists were performed as indicated in Methods in membrane preparations from CHO cells expressing A_{2A}R or A₁R and A_{2A}R or A_{2A}R and D₂R. Binding data were fitted assuming that receptors (also when heteromerizing) form homodimers, and cooperativity (D_{CB} \neq 0, fitting to eq. 2; Materials and Methods) or non-cooperativity (D_{CB} =0, fitting to eq. 3; Materials and Methods) for competitor ligand binding was statistically tested (F test). Only K_{DB1} values (equilibrium dissociation constant of the first binding of B: ZM-241385, MSX-2, SCH-58261, SCH-420814 or KW-6002) are shown, since the analysis demonstrated non-cooperativity for the five A_{2A}R antagonists. Data are mean \pm S.E.M. values of three experiments.

*: $p{<}0.05$ compared to K_{DB1} values in $A_{2A}R$ cells; one-way ANOVA, followed by Newman-Keuls test.

doi:10.1371/journal.pone.0016088.t003

Table 4. Pharmacological parameters for SCH-442416 binding to $A_{2A}R$, A_1R - $A_{2A}R$ and $A_{2A}R$ - D_2R CHO cells.

Parameters	A _{2A} R cells	A ₁ R-A _{2A} R cells	A _{2A} R-D ₂ R cells
K _{DB1}	2.0±0.3 nM	2.4±0.4 nM	7±4 nM
K _{DB2}	8±2 nM	10±2 nM	$5\pm2~\mu M^{**}$
D _{CB}	0	0	-2.3
B ₅₀	4.0±0.6 nM	4.8±0.8 nM	190±80 nM**

Competition experiments of [³H]ZM-241385 (2 nM) binding *versus* increasing concentrations of SCH-442416 were performed as indicated in Methods in membrane preparations from CHO cells expressing $A_{2A}R$ or A_1R and $A_{2A}R$ or $A_{2A}R$ and D_2R . Results were fitted assuming that receptors (also when heteromerizing) form homodimers, and cooperativity ($D_{CB} \neq 0$, fitting to eq. 2; Materials and Methods) or non-cooperativity ($D_{CB} = 0$, fitting to eq. 3; Materials and Methods) of SCH-442416 binding was statistically tested (F test). K_{DB1} and K_{DB2} are, respectively, the equilibrium dissociation constants of the first and second binding of B (SCH-442416) to the dimer. D_{CB} is the "dimer cooperativity" index for the binding of the ligand B, and B₅₀ is the concentration providing half saturation for B. Data are mean \pm S.E.M. values of three experiments. **: p<0.01, respectively compared to the K_{DB2} and B₅₀ values in A₂R and A₁R-A₂A cells; Kruskal-Wallis, followed by Dunn's test.

doi:10.1371/journal.pone.0016088.t004

transfected with the D₂R. When trying to explain the differential action of SCH-442416 observed *in vivo*, it is interesting to note that SCH-442416 showed a much higher affinity for the A_{2A}R in a presynaptic-like than in a postsynaptic-like context. The binding of SCH-442416 to the A_{2A}R-D₂R heteromer displayed a strong negative cooperativity, phenomenon that was not observed for the binding of SCH-442416 to the A₁R-A_{2A}R heteromer. This negative cooperativity explains the pronounced decrease in affinity of A_{2A}R in cells expressing A_{2A}R-D₂R heteromers (B₅₀ values 40 times higher in cells expressing A_{2A}R-D₂R than A₁R-A_{2A}R heteromers).

The loss of affinity of $A_{2A}R$ upon co-expression of D_2R was much less pronounced for ZM-241385, SCH-58261, MSX2 or SCH-420814, for which the affinity was reduced from two to about nine fold. Taking into account that these $A_{2A}R$ antagonists behave similarly than the $A_{2A}R$ agonist CGS-21680 in terms of binding to A_1R - $A_{2A}R$ and $A_{2A}R$ - D_2R heteromers, it is expected that these four compounds compete equally for the binding of the endogenous agonist at pre- and at postsynaptic sites. This would fit with the *in vivo* data, which shows that these compounds have a non-preferred pre-postsynaptic profile. Yet, KW-6002 was the only antagonist whose affinity was not significantly different in cells expressing $A_{2A}R$, A_1R - $A_{2A}R$ heteromers or $A_{2A}R$ - D_2R heteromers. Thus, KW-6002 showed the best relative affinity for $A_{2A}R$ - D_2R heteromers of all coumpounds, which can at least partially explain its preferential postsynaptic profile.

The present results support the notion that receptor heteromers may be used as selective targets for drug development. Main reasons are the very specific neuronal localization of receptor heteromers (even more specific than for receptor subtypes), and a differential ligand affinity of a receptor depending on its partner (or partners) in the receptor heteromer. In the striatum, $A_{2A}R$ provides a particularly interesting target, eventually useful for a variety of neuropsychiatric disorders. $A_{2A}R$ -D₂R and A_1R - $A_{2A}R$ heteromers are segregated in different striatal neuronal elements. While $A_{2A}R$ -D₂R heteromers are located postsynaptically in the dendritic spines of the indirect MSNs [4–6,14], A_1R - $A_{2A}R$

References

- Gerfen CR (2004) Basal Ganglia. In: Paxinos G, ed. The Rat Nervous System. Amsterdam: Elsevier Academic Press. pp 445–508.
- Obeso JA, Rodr´guez-Oroz MC, Rodr´guez M, Arbizu J, Giménez-Amaya JM (2002) The basal ganglia and disorders of movement: pathophysiological mechanisms. News Physiol Sci 17: 51–55.
- DeLong MR, Wichmann T (2007) Circuits and circuit disorders of the basal ganglia. Arch Neurol 64: 20–24.
- 4. Ferre S, Quiroz C, Woods AS, Cunha R, Popoli P, et al. (2008) An update on

receptor heteromers are located presynaptically in glutamatergic terminals contacting the MSNs of the direct pathway [5,11,14]. Blocking postsynaptic A2AR in the indirect MSN should potentiate D_2R -mediated motor activation, which is a strategy already used in the development of anti-parkinsonian drugs [26-28]. However, blocking A2AR in glutamatergic terminals to the direct MSN could potentially be useful in dyskinetic disorders such as Huntington's disease and maybe in obsessive-compulsive disorders and drug addiction [5]. The present results give a mechanistic explanation to the already reported antiparkinsonian activity of KW-6002 [27,28] and suggest that SCH-442416 could be useful in dyskinetic disorders, obsessive-compulsive disorders and in drug addiction. Medicinal chemistry and computerized modeling should help understanding the molecular properties that determine the particular pharmacological profile of SCH-442416 and KW-6002, which may be used as lead compounds to obtain more effective antidyskinetic and antiparkinsonian compounds, respectively. It will also be of importance to take into account potential changes in the expression of pre- and postsynaptic A2ARs and in their respective heteromers which can occur in those mentioned neuropsychiatric disorders. For instance, dopamine denervation seems to differentially modify the expression of striatal A2AR, A1R and D_2R [28–31]. This could be addressed by applying the *in vivo* methodology here described to animal models.

Acknowledgments

We acknowledge the technical help obtained from Jasmina Jiménez (Molecular Neurobiology laboratory, Barcelona University).

Author Contributions

Conceived and designed the experiments: MO JB MB CQ VB SRG CL AC RF VC EIC SF. Performed the experiments: MO JB MB CQ AC VC. Analyzed the data: MO CQ CL AC VC SF. Wrote the paper: MO CL AC VC SF. assessment by fluorescence and bioluminescence energy transfer. J Neurochem 88: 726–734.

- Guo W, Urizar E, Kralikova M, Mobarec JC, Shi L, et al. (2008) Dopamine D2 receptors form higher order oligomers at physiological expresión levels. EMBO J 27: 2293–2304.
- Han Y, Moreira IS, Urizar E, Weinstein H, Javitch JA (2009) Allosteric Communications between protomers of dopamine class A GPCR dimers modulates activation. Nat Chem Biol 9: 688–695.
- Casadó V, Cortes A, Ciruela F, Mallol J, et al. (2007) Old and new ways to calculate the affinity of agonists and antagonists interacting with G-proteincoupled monomeric and dimeric receptors: The receptor-dimer cooperativity index. Pharmacol Ther 116: 343–354.
- Casadó V, Mart' T, Franco R, Lluis C, Mallol J, et al. (1990) A method for binding parameters estimation of A1 adenosine receptor subtype: a practical approach. Anal Biochem 184: 117–123.
- Schwarzschild MA, Agnati L, Fuxe K, Chen JF, Morelli M (2007) Targeting adenosine A2A receptors in Parkinson's disease Trends Neurosci 29: 647–654.

- Jenner P (2003) A2A antagonists as novel non-dopaminergic therapy for motor dysfunction in PD Neurology 61: S32–S8.
- Stacy M, Silver D, Mendis T, Sutton J, Mori A, et al. (2008) A 12-week, placebocontrolled study (6002-US-006) of istradefylline in Parkinson disease Neurology 70: 2233–2240.
- Kaasinen V, Ruottinen HM, Nøgren K, Lehikoinen P, Oikonen V, et al. (2000) Upregulation of putaminal dopamine D2 receptors in early Parkinson's disease: a comparative PET study with [11C] raclopride and [11C]N-methylspiperone J Nucl Med 41: 65–70.
- Pinna A, Corsi C, Carta AR, Valentini V, Pedata F, et al. (2002) Modification of adenosine extracellular levels and adenosine A(2A) receptor mRNA by dopamine denervation. Eur J Pharmacol 446: 75–82.
- Varani K, Vincenzi F, Tosi A, Gessi S, Casetta I, et al. (2010) A2A adenosine receptor overexpression and functionality, as well as TNF-alpha levels, correlate with motor symptoms in Parkinson's disease FASEB J 24: 587–598.