BMJ Open Diabetes Research & Care

New clinical screening strategy to distinguish *HNF1A* variant-induced diabetes from young early-onset type 2 diabetes in a Chinese population

Yumin Ma ^(D), ¹ Siqian Gong, ¹ Xirui Wang, ² Xiaoling Cai, ¹ Xinhua Xiao, ³ Weijun Gu, ⁴ Jinkui Yang, ⁵ Liyong Zhong, ⁶ Jianzhong Xiao, ⁷ Meng Li ^(D), ¹ Wei Liu, ¹ Simin Zhang, ¹ Xianghai Zhou, ¹ Yufeng Li, ⁸ Lingli Zhou, ¹ Yu Zhu, ¹ Yingying Luo, ¹ Qian Ren, ¹ Xiuting Huang, ¹ Xueying Gao, ¹ Xiuying Zhang, ¹ Rui Zhang, ¹ Ling Chen ^(D), ¹ Fang Wang, ⁶ Qiuping Wang, ⁹ Mengdie Hu, ¹ Xueyao Han, ¹ Linong Ji¹

ABSTRACT

To cite: Ma Y, Gong S, Wang X, *et al.* New clinical screening strategy to distinguish *HNF1A* variantinduced diabetes from young early-onset type 2 diabetes in a Chinese population. *BMJ Open Diab Res Care* 2020;**8**:e000745. doi:10.1136/ bmjdrc-2019-000745

Additional material is published online only. To view please visit the journal online (http://dx.doi.org/10.1136/ bmjdrc-2019-000745).

YM, SG and XW contributed equally. XHa and LJ contributed equally.

Received 9 July 2019

Revised 11 February 2020 Accepted 9 March 2020

Check for updates

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Linong Ji; jiln@bjmu.edu.cn and Dr Xueyao Han; xueyaohan@sina.com **Objective** Maturity-onset diabetes of the young caused by hepatocyte nuclear factor-1 alpha (*HNF1A*) variants (*HNF1A*-MODY) is a common form of monogenetic diabetes. Although patients with *HNF1A*-MODY might specifically benefit from sulfonylurea treatment, available methods for screening this specific type of diabetes are not cost-effective. This study was designed to establish an optimized clinical strategy based on multiple biomarkers to distinguish patients with *HNF1A*-MODY from clinically diagnosed early-onset type 2 diabetes (EOD) for genetic testing in a Chinese population.

Research design and methods A case–control study including 125 non-related young patients with EOD and 15 probands with *HNF1A*-MODY (cohort 1) was conducted to evaluate reported biomarkers for *HNF1A*-MODY. A cut-off for the fasting insulin (Fins) level, the 97.5 percentile of 150 healthy subjects with normal components of metabolic syndrome (cohort 2), was used to filter out individuals with obvious insulin resistance (Fins <102 pmol/L). An optimized clinical screening strategy (*HNF1A*-CSS) was established, and its effectiveness was assessed in another group of 410 young patients with EOD (cohort 3).

Results In cohort 1, body mass index (BMI), serum high-density lipoprotein cholesterol (HDL-c) and highsensitivity C reactive protein (hs-CRP) levels were confirmed to be useful for the differential diagnosis of *HNF1A*-MODY. In cohort 3, eight probands with *HNF1A*-MODY were identified. In cohort 3 and young relatives with *HNF1A*-MODY, meeting three of four criteria (BMI <28 kg/m², hs-CRP <0.75 mg/L, Fins <102 pmol/L and HDL-c >1.12 mmol/L), the sensitivity and specificity of *HNF1A*-CSS were 100% and 69.3%, respectively. In the pooled analysis of all young patients, *HNF1A*-CSS displayed 90.5% sensitivity and 73.6% specificity for identifying patients with *HNF1A*-MODY among those with clinically diagnosed EOD.

Conclusion Our *HNF1A*-CSS is useful for distinguishing patients with *HNF1A*-MODY from patients with EOD in a young Chinese population.

Significance of this study

What is already known about this subject?

- The overlap of phenotypes makes patients with maturity-onset diabetes of the young (MODY) difficult to distinguish from patients with early-onset type 2 diabetes (EOD) in the clinical setting.
- A potentially efficient clinical screening strategy (CSS) including readily available clinical and biochemical features confirmed to be associated with hepatocyte nuclear factor-1 alpha (*HNF1A*)-MODY is needed.

What are the new findings?

- Our study is the first to establish an HNF1A-CSS with the largest sample in an East Asian population. In this study, we confirmed the effectiveness of several reported biomarkers (high-sensitivity C reactive protein (hs-CRP), body mass index (BMI) and highdensity lipoprotein cholesterol (HDL-c)) for differentiating young patients with HNF1A-MODY from patients with EOD in a Chinese population.
- A cut-off for the fasting insulin (Fins) level derived from upper limit of normal reference range further excluded patients with type 2 diabetes.
- By including multiple biomarkers (hs-CRP, BMI, HDL-c and Fins), we established and evaluated a CSS with better performance than the reported models and a single biomarker for identifying HNF1A-MODY in a Chinese population with EOD.

How might these results change the focus of research or clinical practice?

The CSS composed of the four criteria established in our study is practical and easy to be widely used in the real world.

INTRODUCTION

Maturity-onset diabetes of the young (MODY) caused by hepatocyte nuclear factor-1 alpha



(HNF1A) gene variants is one of the most common types of MODY (HNF1A-MODY),¹² usually misdiagnosed with either type 1 (T1DM) or type 2 diabetes mellitus (T2DM), and often receive incorrect treatment.^{2–4} Patients with *HNF1A*-MODY are sensitive to sulfonylureas (SUs),⁵ and their therapy is often changed from insulin to SUs once the diagnosis is established.⁶

Early-onset T2DM (EOD, diagnosed before 40 years old) has become increasingly prevalent in the past decades worldwide. Because of the overlap of EOD phenotypes with the classical forms of MODY,⁷ clinicians experience challenges in distinguishing patients with MODY from EOD. Despite tremendous advances in DNA sequencing technology in recent years, the application of genetic testing for common MODY subtypes to all patients remains very costly. And the interpretation of genetic findings without a clinical context is challenging. Therefore, a cost-effective screening strategy to identify subjects for genetic testing is urgently needed. In fact, using basic clinical characteristics, a prediction model to determine the probability of MODY has been developed,⁸ and if other important clinical markers such as insulin secretion and insulin resistance were included, the model would be improved. Particularly, some MODY subtype-specific biomarkers have been discovered, and they make it possible to establish a strategy to clinically screen patients with MODY due to single gene.⁹¹⁰

Some researchers recommended clinical screening strategies (CSS) for identifying individuals with diabetes who have a higher probability of carrying causal variants in the *HNF1A* gene before molecular testing.²⁷ These strategies are very attractive since they have the potential to reduce both the labor and costs associated with diagnosing a disease condition. In the past, single index or compositional indices, including the absence of autoantibodies associated with T1DM, young age at diabetes diagnosis, a non-obese status, a strong family history of diabetes in multiple generations, high-sensitivity C reactive protein (hs-CRP) and high-density lipoprotein cholesterol (HDL-c) levels, have been used to screen for HNF1A-MODY.^{7 11–13} However, the performance of these approaches in clinical settings has been suboptimal. In the present study, we used multiple biomarkers to establish a new optimized screening strategy to identify HNF1A-MODY in patients with EOD.

METHODS

Study subjects

First, 125 patients with EOD and 15 patients with *HNF1A*-MODY were included (cohort 1). The cohort was used to evaluate the ability of the reported clinical biomarkers (body mass index (BMI), HDL-c and hs-CRP) associated with *HNF1A*-MODY to differentially diagnose *HNF1A*-MODY.

The patients with EOD were consecutively recruited from the Department of Endocrinology and Metabolism of Peking University People's Hospital (PUPH) in Beijing from June 2016 to January 2018. They were diagnosed at 14–40 years old. The 125 patients with EOD were confirmed not to carry any rare variants of *HNF1A*, *HNF4A*, *HNF1B*, *GCK* or mitochondrial DNA 3243A>G mutation by DNA sequencing.

The 15 patients with a confirmed genetic diagnosis of *HNF1A*-MODY by DNA sequencing were from other clinical centers in Beijing. Each carried one of 13 rare variants of *HNF1A* (p.Thr10Met, p.Pro33Leu, p.Arg131Trp, p.Gl-n176Ter, p.Arg203His, p.Gly292Argfs*25, p.Ala311Asp, p.Pro379Thr, IVS2 +1 G>A, p.Pro519Leu, p.Gln541Ter, p.Val567Ile and p.Ile618Met) (online supplementary table S1), which were pathogenic or likely pathogenic according to the standards and guidelines recommended by the American College of Medical Genetics (ACMG) (online supplementary table S2).

Second, 150 subjects (cohort 2) with normal glucose tolerance (NGT) and normal components of metabolic syndrome¹⁴ were used to determine the cut-off value for the fasting serum insulin (Fins) level to identify patients with *HNF1A*-MODY.

Increased insulin secretion from beta cells compensates for insulin resistance and does not result in hyperglycemia in normal individuals with NGT¹⁵; metabolic syndrome is strongly correlated with insulin resistance. Thus, in the subjects with NGT and no signs of insulin resistance (a condition with normal components of metabolic syndrome), an upper limit of the reference range of Fins levels would help to screen individuals with insulin resistance because an increase in Fins beyond the upper limit would indicate the occurrence of insulin resistance. In contrast to *HNF1A*-MODY, which is characterized by impaired beta cell function, insulin resistance is a major clinical feature of T2DM.

Cohort 2 were randomly selected from the 3345 subjects of the Pinggu cohort, which was described in our previous study.¹⁰ The subjects were aged \leq 45 years at examination. Their fasting plasma glucose (FPG) levels were less than 6.1 mmol/L, 2-hour plasma glucose levels during the oral glucose tolerance test were less than 7.8 mmol/L and hemoglobin A1c (HbA1c) levels were less than 42 mmol/mol (6.0%). The clinical characteristics of the 150 subjects are presented in online supplementary table S3.

Third, a group of patients with clinically diagnosed as EOD were recruited in PUPH from January 2014 to May 2016 and used to validate the effectiveness of the CSS established in this study for *HNF1A*-MODY.

To avoid the interference from other subtypes of monogenic diabetes, four patients with other common subtype monogenic diabetes (two mitochondrial 3243A>G, one *HNF4A*-MODY and one *GCK*-MODY) (shown in online supplementary table S4) were excluded, and the remaining 410 patients (cohort 3) were used for analysis.

Clinical examinations and laboratory tests

All participants were examined in the morning after an overnight fast of 10–12 hours. Their clinical information

copyright.

and blood samples were collected.¹⁰ HbA1c, C-peptide, insulin, plasma glucose, total cholesterol (TC), HDL-c, low-density lipoprotein cholesterol (LDL-c), triglyceride (TG) and hs-CRP levels were measured. Fresh urine samples were collected to assess the urinary albumin/ creatinine ratio (ACR).

The serum for quality control and serum samples from the patients were determined for glutamic acid decarboxylase antibody (GADA) (RSR Limited, UK), islet cell antibodies (ICA) (Biomerica, USA) and insulin antibody (IA) (Orgentec, Germany) using commercialized ELISA kit. The assays were performed according to the instructions from the manufacturer.

According to the Chinese-specific diagnostic criteria for obesity, a BMI $\geq 28 \text{ kg/m}^2$ was defined as obesity.¹⁶ The diagnosis of obesity in subjects aged between 14 and 18 years was based on weight-for-age curves for Chinese individuals.¹⁷

Inclusion and exclusion criteria for the study subjects with EOD

All subjects were of Chinese Han ethnicity and aged ≤45 years. Diabetes was clinically diagnosed before 40 years according to WHO criteria.¹⁸ Patients were excluded from this study according to the following criteria: (1) the presence of typical clinical features of T1DM (absolute insulin deficiency and insulin-dependent treatment after diagnosis); (2) other specific forms of diabetes (eg, chronic pancreatitis and steroid-induced diabetes; (3) if any autoantibodies (GADAs, ICAs and IAs) were positive, but for the patients with exogenous insulin treatment, if only IA was positive and they had no other typical features of T1DM, they were also included; (4) a fasting serum C-peptide level <0.1 ng/mL; (5) the presence of an autoimmune disease; (6) the presence of an acute infectious disease or the acute stage of myocardial infarction and stroke and (7) an hs-CRP level >10 mg/L.

DNA sequencing and genetic diagnosis

Genomic DNA was extracted from whole blood. The patients in cohort 1 were sequenced to screen for the variants in exons of HNF1A, GCK, HNF4A, HNF1B or mitochondrial 3243A>G mutation by Sanger method. The patients in cohort 3 were screened for mitochondrial 3243A>G mutation by Sanger sequencing, and their DNA samples were also used for exon capture and next-generation sequencing. An Agilent SureselectXT2 customized gene panel was designed for HNF1A and other genes for common MODY subtypes (GCK, HNF4A and HNF1B). A HiSeq2000 Sequencing System (Illumina, San Diego, California, USA) was used for sequencing. Rare variants (≤0.005 in the Exome Aggregation Consortium) with a sequencing depth of at least 20 and variants located in coding regions with non-synonymous and frameshift variants or in splicing sites near the introns were included and validated by PCR and Sanger sequencing.

Three hundred controls with NGT and normal HbA1c were used to screen the rare variants identified in this study by Sanger method, which is helpful for determining whether the variants are damaging or not.

The pathogenicity of these variants was evaluated using the standards and guidelines recommended by ACMG¹⁹ (online supplementary table S2). If patients carried pathogenic or likely pathogenic variants, they were diagnosed with *HNF1A*-MODY.

Selection of clinical biomarkers and establishment of a CSS to differentiate *HNF1A*-MODY from EOD

As reported in previous studies, most patients with *HNF1A*-MODY have no insulin resistance.^{8 20–22} However, a recent study pointed that the patients with both *HNF1A*-MODY and overweight/obesity should not be ignored.²³ They also present low insulin secretion, low serum hs-CRP levels, ^{9 24 25} low prevalence of dyslipidemia (such as a normal or higher HDL-c level than patients with T2DM) and a low rate of insulin resistance.^{4 7} All relevant biomarkers were used to establish the clinical screening model for identifying individuals with a higher probability of *HNF1A*-MODY. If different cut-off values were reported for the same variable, the cut-off value with higher sensitivity was used in this study. The flowchart showing the design of our study is presented in figure 1.

Statistical analysis

Statistical analyses were performed using the Statistical Package for Social Sciences (SPSS for Windows, V.23.0; Chicago, Illinois, USA). Student's t-test was used to analyze quantitative traits. The χ^2 test or Fisher's exact test was used to assess categorical traits. Normally distributed continuous variables are presented as means and SDs (means±SDs), and non-normally distributed variables are presented as medians (IQRs). Categorical variables are presented as medians (IQRs). Categorical variables are presented as mumbers and percentages. The ability of each biomarker and *HNF1A* score to discriminate *HNF1A*-MODY from EOD was assessed by calculating the area under the receiver operating characteristic curve (ROC) curve (C-statistics). A p value <0.05 was considered significant.

RESULTS

Evaluation of three reported clinical biomarkers for *HNF1A*-MODY in cohort 1

Compared with patients with EOD (table 1), a lower percentage of patients with *HNF1A*-MODY exhibited obesity and dyslipidemia; these patients were also diagnosed at a younger age and presented lower levels of BMI, HbA1c, TC, TGs, LDL-c, uric acid and hs-CRP, as well as higher levels of HDL-c.

As shown in online supplementary figure S1, the C-statistic for hs-CRP was 0.902 (95% CI 0.840 to 0.946), and the cut-off value was 0.72 mg/L, with 76.9% sensitivity and 92.0% specificity. The C-statistic for BMI was 0.811 (95% CI 0.736 to 0.872), and the cut-off value was 22.7 kg/m², with 60.0% sensitivity and 92.0% specificity. The



Figure 1 Flowchart showing the design of our study. CSS, clinical screening strategy; *HNF1A*, hepatocyte nuclear factor-1 alpha; MODY, maturity-onset diabetes of the young.

C-statistic for HDL-c was 0.763 (95% CI 0.684 to 0.831), and the cut-off value was 1.20 mmol/L, with 66.7% sensitivity and 88.0% specificity.

When the reported cut-off values for hs-CRP (<0.75 mg/L),⁹ HDL-c $(>1.12 \text{ mmol/L})^{26}$ and BMI (the cut-off for obesity in Chinese individuals is 28 kg/m^2)¹⁶ that have been used to identify *HNF1A*-MODY were applied to cohort 1, the sensitivity and specificity were 76.9% and 92.0% for hs-CRP, 73.3% and 78.4% for HDL-c, and 93.3% and 62.4% for BMI, respectively. These cut-off values displayed higher sensitivity than those determined in cohort 1.

Establishment of a cut-off value for the Fins level for diagnosing *HNF1A*-MODY

In cohort 2, the 97.5 percentile of the Fins level was 102 pmol/L. In cohort 1, all of 15 patients with *HNF1A*-MODY presented Fins levels less than 102 pmol/L or received insulin treatment, 88 individuals of 125 patients with EOD had Fins levels of <102 pmol/L. Among the subjects in the Pinggu cohort with Fins levels greater than 102 pmol/L (n=390, unpublished data), 95.1% were diagnosed with metabolic syndrome. Thus, using the Fins cut-off of 102 pmol/L, we were able to exclude most patients with T2DM presenting obvious insulin resistance from patients with EOD.

Establishment of the CSS for HNF1A-MODY (HNF1A-CSS)

Due to our limited sample size, the cut-off values defined in the current study might not be the most precise values for the clinical screening of *HNF1A*-MODY. Among the reported cut-off values and our values for screening patients with *HNF1A*-MODY, we selected those with the highest sensitivity and established an *HNF1A*-CSS. To maximally avoid missing patients with *HNF1A*-MODY, this *HNF1A*-CSS comprises the following four criteria: (1) non-obesity, BMI < $28 \text{ kg/m}^{2,16}$ (2) hs-CRP level <0.75 mg/L,⁹ (3) insulin deficiency or a lack of insulin resistance (insulin treatment or Fins <102 pmol/L in patients without insulin treatment) and (4) an HDL-c level >1.12 mmol/L.²⁶

Screening of patients with HNF1A-MODY in cohort 3

In cohort 3, nine rare heterozygous HNF1A variants (p.Arg131Trp, p.Arg200Trp, p.Arg263Cys, p.Gln324Ter, IVS8-1G>A, p.Pro112Leu, p.Glu240Val, p.Glu548Gly and p.Arg159fs (c.474_475insA)) (online supplementary table S1) were identified in nine study subjects. The mothers of the probands carrying p.Arg200Trp, p.Arg131Trp and IVS8-1G>A variants also carried the same variants and were diagnosed with EOD (online supplementary figure S2). The parents of the probands carrying the p.Arg263Cys variant did not have diabetes and did not carry the p.Arg263Cys variant. Unfortunately, we failed to recruit the family members of the other probands. According to the guidelines of the ACMG, eight rare variants (p.Arg131Trp, p.Arg200Trp, p.Arg263Cys, p.Gln324Ter, IVS8-1G>A, p.Glu240Val, p.Pro112Leu and p.Arg159fs (c.474_475insA)) are pathogenic or likely pathogenic (online supplementary table S2). Thus, eight patients with HNF1A-MODY were identified in cohort 3. Nine patients with HNF1A-MODY, including eight probands and one younger family member with the p.Arg200Trp variant and diabetes (clinical features are summarized in online supplementary table S5), were compared with patients with EOD. Similar to cohort 1, patients with HNF1A-MODY from cohort 3 were also characterized by a lower percentage of obesity and dyslipidemia, and displayed a lower age at diagnosis and lower

Table 1 Clinical characteristics of the patients with HNF1A-MODY and young patients with EOD in cohort 1					
Characteristic	EOD (N=125)	HNF1A-MODY (N=15)	P value		
Sex, male/female	91/34	8/7	0.138		
Age, mean (SD), years	32.4 (6.6)	29.4 (9.3)	0.120		
Age at diagnosis, mean (SD), years	29.7 (6.0)	25.3 (7.5)	0.010*		
Duration of DM, median (IQR), years	1.0 (0.1 to 5.0)	3.0 (1.0 to 6.0)	0.083		
Family history of DM, n (%)	105 (84.0)	14 (93.3)	0.469		
BMI, mean (SD), kg/m ²	28.1 (4.1)	23.0 (4.0)	<0.0001***		
SBP, mean (SD), mm Hg	125.2 (13.9)	122.2 (14.3)	0.433		
DBP, mean (SD), mm Hg	78.8 (11.2)	76.6 (10.2)	0.474		
FPG, median (IQR), mmol/L	7.32 (6.16 to 9.75)	7.20 (5.10 to 7.83)	0.222		
HbA1c, median (IQR), mmol/mol	76.0 (58.0 to 95.5)	56.0 (42.0 to 67.0)	0.001**		
HbA1c, median (IQR), %	9.10 (7.50 to 10.85)	7.30 (6.00 to 8.30)	0.001**		
TC, mean (SD), mmol/L	4.87 (1.10)	4.21 (0.69)	0.024*		
LDL-c, mean (SD), mmol/L	3.14 (0.87)	2.58 (0.75)	0.019*		
HDL-c, mean (SD), mmol/L					
Male	0.96 (0.19)	1.15 (0.35)	0.016*		
Female	1.02 (0.24)	1.45 (0.46)	0.001**		
Triglycerides, median (IQR), mmol/L	1.72 (1.22 to 2.70)	1.10 (0.74 to 1.53)	0.002**		
hs-CRP, median (IQR), mg/L	2.14 (1.22 to 4.02)	0.41 (0.10 to 2.13)	0.002*		
CRE, mean (SD), µmol/L	60.3 (14.8)	68.7 (21.4)	0.056		
UA, mean (SD), μmol/L					
Male	390.8 (92.7)	313.7 (53.5)	0.033*		
Female	353.2 (92.6)	266.6 (61.0)	0.024*		
ACR, median (IQR), mg/g	8.36 (3.90 to 20.25)	5.00 (3.85 to 6.50)	0.078		
eGFR, median (IQR), mL/min/1.73 m ²	126.0 (113.3 to 139.0)	134.5 (119.0 to 160.2)	0.272		
Comorbidities and complications					
Hypertension, n (%)	19 (15.2)	1 (6.7)	0.696		
Dyslipidemia, n (%)	69 (55.2)	2 (13.3)	0.002**		
Diabetic nephropathy, n (%)	9 (7.2)	1 (6.7)	1.000		
Diabetic retinopathy, n (%)	11 (8.8)	4 (26.7)	0.058		
Coronary heart disease, n (%)	1 (0.8)	1 (6.7)	0.203		
Cerebrovascular disease, n (%)	2 (1.6)	1 (6.7)	0.290		
Obesity, n (%)	62 (49.6)	1 (6.7)	0.002**		
Treatment					
OHA, n (%)	65 (52.0)	10 (66.7)	0.258		
Insulin, n (%)	40 (32.0)	8 (53.3)	0.077		

Data are presented as means (SD) or medians (IQRs); categorical variables are presented as n (%). eGFR (mL/min per 1.73m²) = 175×CRE(mg/dL)^{-1.234}×age (years)^{-0.179}×0.79 (if female).

*p<0.05, **p<0.01 and ***p<0.001.

ACR, urinary albumin/creatinine ratio; BMI, body mass index; CRE, serum creatinine; DBP, diastolic blood pressure; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; EOD, young early-onset type 2 diabetes; FPG, fasting plasma glucose; HbA1c, hemoglobin A1c; HDL-c, high-density lipoprotein cholesterol; HNF1A, hepatocyte nuclear factor-1 alpha; hs-CRP, high-sensitivity C reactive protein; LDL-c, low-density lipoprotein cholesterol; MODY, maturity-onset diabetes of the young; OHA, oral hypoglycemic agent; SBP, systolic blood pressure; TC, total cholesterol; UA, uric acid.

levels of BMI, TGs and hs-CRP, as well as higher levels of HDL-c (table 2).

Evaluation of the effectiveness of the HNF1A-CSS in cohort 3

As shown in figure 2, we tested the effectiveness of HNF1A-CSS on all probands with HNF1A-MODY in cohort 3 and the young relatives with HNF1A-MODY (n=9). Nine individuals met three criteria, eight met four criteria and none met one criterion or two criteria. There were 113 patients including 9 with HNF1A-MODY meeting at least three of four criteria (HNF1A-CSS1), according to

HNF1A-CSS1. This indicates that out of sequencing 12.6 patients only one patient could be screened with HNF1A-MODY. The sensitivity and specificity of HNF1A-CSS1 were 100% and 69.7%, respectively, and the positive and negative prediction values were 6.9% and 100%, respectively. On the other hand, there were 51 patients including eight with HNF1A-MODY meeting all of four criteria (HNF1A-CSS2). Through sequencing only 6.4 patients, we could identify one patient with HNF1A-MODY according to HNF1A-CSS2. The sensitivity and specificity

Table 2 Clinical characteristics of the patients with HNF1A-MODY and young patients with EOD in cohort 3					
Characteristic	EOD (N=402)	HNF1A-MODY (N=9)	P value		
Sex, male/female	273/129	3/6	0.064		
Age, median (IQR), years	33.0 (29.0 to 36.0)	31.0 (21.0 to 33.5)	0.146		
Age at diagnosis, median (IQR), years	30.0 (26.0 to 33.0)	18.0 (13.0 to 30.5)	0.006**		
Duration of DM, median (IQR), years	2.0 (1.0 to 6.0)	4.0 (1.0 to 17.0)	0.077		
Family history of DM, n (%)	307 (76.4)	8 (88.9)	0.692		
Waist circumference†, mean (SD), cm	92.5 (12.5)	78.9 (9.0)	0.002**		
BMI, mean (SD), kg/m ²	26.8 (4.6)	20.9 (1.9)	<0.0001***		
SBP, mean (SD), mm Hg	122.6 (15.7)	113.0 (22.3)	0.073		
DBP, mean (SD), mm Hg	78.8 (12.1)	69.9 (15.2)	0.058		
FPG, median (IQR), mmol/L	8.37 (6.53 to 11.05)	6.86 (5.58 to 8.63)	0.068		
HbA1c, median (IQR), mmol/mol	77.0 (57.3 to 98.0)	60.5 (47.0 to 75.0)	0.095		
HbA1c, median (IQR), %	9.20 (7.43 to 11.10)	7.65 (6.45 to 8.98)	0.095		
TC, mean (SD), mmol/L	4.82 (1.18)	4.73 (1.41)	0.709		
LDL-c, mean (SD), mmol/L	2.82 (0.85)	2.84 (0.90)	0.940		
HDL-c†, mean (SD), mmol/L	1.04 (0.27)	1.41 (0.20)	<0.0001***		
Triglycerides, median (IQR), mmol/L	1.74 (1.16 to 2.74)	1.05 (0.87 to 1.53)	0.033*		
hs-CRP, median (IQR), mg/L	1.47 (0.71 to 3.21)	0.17 (0.11 to 0.32)	<0.0001***		
CRE, median (IQR), µmol/L	62.0 (50.0 to 72.0)	69.0 (46.0 to 89.0)	0.520		
UA†, median (IQR), µmol/L	345.0 (283.0 to 411.0)	350.0 (264.5 to 363.5)	0.796		
ACR, median (IQR), mg/g	8.61 (4.19 to 30.00)	10.87 (6.67 to 1004.42)	0.368		
eGFR, median (IQR), mL/min/1.73 m ²	139.4 (120.6 to 167.9)	126.4 (73.4 to 180.4)	0.518		
Comorbidities and complications					
Hypertension, n (%)	56 (13.9)	1 (11.1)	1.000		
Dyslipidemia, n (%)	375 (93.3)	2 (22.2)	<0.0001***		
Diabetic nephropathy, n (%)	50 (12.4)	3 (33.3)	0.097		
Diabetic retinopathy, n (%)	36 (9.0)	3 (33.3)	0.044*		
Coronary heart disease, n (%)	9 (2.2)	0 (0)	1.000		
Cerebrovascular disease, n (%)	5 (1.2)	0 (0)	1.000		
Obesity, n (%)	156 (38.8)	0 (0)	0.015*		
Treatment					
OHA, n (%)	171 (42.5)	5 (55.6)	0.506		
Insulin, n (%)	174 (43.3)	4 (44.4)	1.000		

Data are presented as the means (SD) or medians (IQR); categorical variables are presented as n (%). eGFR (mL/min per 1.73m²)=175×CRE(mg/dL)^{-1.234}×age (years)^{-0.179}×0.79 (if female).

*p<0.05, **p<0.01 and ***p<0.001.

+Three male patients were diagnosed with HNF1A-MODY, and therefore, the waist circumference, HDL-c level and uric acid level of all patients with HNF1A-MODY (males and females) were calculated in this analysis.

ACR, urinary albumin/creatinine ratio; BMI, body mass index; CRE, serum creatinine; DBP, diastolic blood pressure; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; EOD, early-onset type 2 diabetes; FPG, fasting plasma glucose; HbA1c, hemoglobin A1c; HDL-c, high-density lipoprotein cholesterol; HNF1A, hepatocyte nuclear factor-1 alpha; hs-CRP, high-sensitivity C reactive protein; LDL-c, low-density lipoprotein cholesterol; MODY, maturity-onset diabetes of the young; OHA, oral hypoglycemic agent; SBP, systolic blood pressure; TC, total cholesterol; UA, uric acid.

of CSS2 were 88.9% and 89.6%, respectively. The positive and negative prediction values were 15.7% and 99.7%, respectively. Notably, HNF1A-CSS1 exhibited the highest sensitivity and HNF1A-CSS2 exhibited the highest specificity for clinically diagnosing HNF1A-MODY.

We conducted an analysis by pooling all patients with EOD and HNF1A-MODY in this study (24 with HNF1A-MODY and 527 with EOD). As shown in figure 3 and online supplementary figures S3 and S4, after excluding three patients with missing data for the four criteria from cohort 1, 21 patients with HNF1A-MODY were included in the analysis. Nineteen of these patients (90.5%) met the HNF1A-CSS1 criteria and 15 patients (71.4%) met

the HNF1A-CSS2 criteria, with specificities of 73.6% and 91.3%, respectively.

DISCUSSION

In the present study, we confirmed the utility of several reported biomarkers (hs-CRP, BMI and HDL-c) for differentiating young patients with HNF1A-MODY from EOD in a Chinese population. A new HNF1A-CSS composed of all the aforementioned biomarkers and the Fins level was established for the clinical identification of individuals with a high probability of having HNF1A-MODY among patients with EOD. This study is the first to establish a



Figure 2 Flowchart for the approach used to distinguish between patients with *HNF1A*-MODY and young patients with earlyonset type 2 diabetes in cohort 3. *A family member with *HNF1A*-MODY is included in this analysis. *HNF1A*-CSS0 indicates that patients meet two criteria, one criterion or zero criteria, and *HNF1A*-CSS1 indicates that patients meet three or four criteria, and HNF1A-CSS2 indicates that patients meet four criteria. BMI, body mass index; CSS, clinical screening strategy; EOD, early-onset type 2 diabetes; Fins, fasting insulin; HDL-c, high-density lipoprotein cholesterol; *HNF1A*, hepatocyte nuclear factor-1 alpha; hs-CRP, high-sensitivity C reactive protein; MODY, maturity-onset diabetes of the young.

CSS for *HNF1A*-MODY in East Asian population using the largest sample to date.

The major obstacle for the diagnosis of patients with *HNFIA*-MODY is that the diagnosis rate based on existing



Figure 3 Distribution of patients with HNF1A-MODY among the four groups according to the four criteria. In this figure, patients with HNF1A-MODY were divided into four groups according to the following four criteria: BMI <28 kg/ m², hs-CRP level <0.75 mg/L, insulin treatment or Fins level <102 pmol/L without insulin treatment and an HDL-c level >1.12 mmol/L. Twenty-one patients with HNF1A-MODY were included in our study. The hs-CRP level was not available for two patients with HNF1A-MODY, and the Fins level was not available for one patient; thus, these patients were excluded from this analysis. Fifteen patients met four criteria, two patients met the BMI, Fins and HDL-c criteria, one patient met the BMI, Fins and hs-CRP criteria, and another patient met the Fins, hs-CRP and HDL-c criteria. Only two patients met two criteria (BMI and Fins). BMI, body mass index; CSS, clinical screening strategy; Fins, fasting insulin; HDL-c, highdensity lipoprotein cholesterol; HNF1A, hepatocyte nuclear factor-1 alpha; hs-CRP, high-sensitivity C reactive protein; MODY, maturity-onset diabetes of the young.

clinical screening criteria is relatively low (10%-14%).⁷¹¹ A recent study confirmed that the addition of hs-CRP levels to other clinical parameters used for identifying *HNF1A*-MODY slightly improved the performance of the predictive model.²⁷ Different cut-off values for hs-CRP levels (0.25–0.75 mg/L) have been reported in different study populations, with a sensitivity ranging from 71% to 83% and a specificity ranging from 68% to 86%.^{9 24 25} Our study confirmed that the hs-CRP level is a useful biomarker for the differential diagnosis of *HNF1A*-MODY in Chinese population with EOD using a similar cut-off value (0.72 mg/L).

A higher HDL-c level has also been observed in patients with *HNF1A*-MODY compared with those with T2DM or MODY-like phenotypes.^{7 28} However, only one study reported that the cut-off value for HDL-c (1.12 mmol/L) was able to distinguish patients with *HNF1A*-MODY from patients with T2DM.²⁶ These findings were similar to our results, and with greater effectiveness in diagnosing females (online supplementary figure S3) in our study. In fact, in *HNF1A*^{-/-} mice, increased plasma cholesterol levels are predominantly located in large, buoyant HDL particles, likely due to the reduced activity of the HDLcatabolic enzyme hepatic lipase and increased expression of the HDL-esterifying enzyme lecithin: cholesterol acyl transferase.²⁹ Not surprisingly, HDL-c is a relatively specific biomarker for *HNF1A*-MODY.

HNF1A encodes a transcription factor that increases insulin secretion. Impaired insulin secretion, but not insulin resistance, is the major pathogenic factor associated with *HNF1A*-MODY. Although the clinical parameters associated with impaired insulin secretion or insulin resistance differ between patients with *HNF1A*-MODY and T2DM, their application in the differential diagnosis of *HNF1A*-MODY had not been well assessed. In our study, we adopted the upper limit of the normal reference range for the Fins level as a cut-off and expected to exclude most patients with obvious insulin resistance. Unfortunately, 88/125 patients with EOD in cohort 1 and 336/402 with copyright

EOD in cohort 3 had the Fins levels of <102 pmol/L or insulin treatment, which indicates the Fins cut-off was not a good indicator for distinguishing *HNF1A*-MODY and EOD when it was used as a single marker, but it remained helpful to filter out some patients with both T2DM and insulin resistance, and in turn reduce the number of individuals for genetic screening.

The absence of obesity is one criterion for clinically screening HNF1A-MODY in Caucasians,³⁰ and we confirmed that the obesity-specific cut-off for BMI in the Chinese population is also sufficiently sensitive to identify patients with HNF1A-MODY in a Chinese population. In fact, it was reported that a few of patients with monogenic diabetes had overweight or even obesity,²³ and some patients with HNF1A-MODY would become overweight or obesity when they were treated with insulin. As observed in our study, one patient with HNF1A-MODY (carrier of p.Pro33Leu in cohort 1) was obese, which might attribute to long-term insulin injection because her maximum body weight occurred after insulin treatment. Thus, we recommended 28 kg/m^2 as the cut-off for clinically identifying HNF1A-MODY instead of the values identified in our ROC analysis (online supplementary figures S1 and S3) to avoid as much as possible missing patients with HNF1A-MODY.

Similar to other studies, 13 of the 23 patients with *HNF1A*-MODY (15 patients from cohort 1 and 8 from cohort 3) identified in our study (56.5%) were diagnosed before the age of 25 years old, but the family history of diabetes did not differ significantly between the patients with *HNF1A*-MODY and those with EOD. Thus, many patients with *HNF1A*-MODY would be missed if a stringent criterion^{2 7 11} (eg, onset before 25 years of age or a family history of diabetes) is adopted to clinically screen patients for genetic testing.

Multiple biomarkers have been used in previous studies to improve the sensitivity and specificity of predictive models for HNF1A-MODY. One study, which included the largest sample of patients with *HNF1A*-MODY,⁷ reported that a clinical predictive model for HNF1A-MODY based on age, BMI, number of generations diagnosed with diabetes, presence of diabetes symptoms and geographic origin only exhibited 49% specificity when the sensitivity was set to 90%. However, its performance has not been confirmed in other populations, and this model must be improved by introducing new biomarkers. Our CSS for HNF1A-MODY, which consists of multiple criteria, would be effective and partly make up the shortcoming of single criterion. For example, the individuals with obesity might have opportunity to be picked up through other criteria, such as Fins, HDL-c or hs-CRP. Although insulin treatment could cause weight gain, insulin treatment itself is one of four criteria.

According to the available clinical screening criteria and after considering the cut-off values for several new biomarkers and pathophysiological characteristics of *HNF1A*-MODY,^{2 7 22} we established the *HNF1A*-CSS to identify patients with *HNF1A*-MODY in the clinic. Because a higher sensitivity is necessary to increase the efficiency of screening for *HNF1A*-MODY, when balancing specificity and sensitivity, *HNF1A*-CSS1 exhibited better performance (90.5% sensitivity and 73.6% specificity) in analyzing pooled data than previously reported models (90% sensitivity and 49% specificity)^{7 9 24 25} and single biomarker (figure 3 and online supplementary figure S4). Certainly, *HNF1A*-CSS2 exhibited higher specificity, but it might miss more patients with *HNF1A*-MODY.

In this study, we established an approach to identify candidates for genetic testing of HNF1A-MODY rather than the full range of MODYs. In fact, each subtype MODY has its specific clinical features and proper therapeutic regimen. For example, hs-CRP is a biomarker for HNF1A-MODY⁹ but not for HNF4A-MODY; and GCK-MODY was characterized by stable mild hyperglycemia. American Diabetes Association recommended that SUs were first-line therapy for HNF1A/4A-MODY and the patients with GCK-MODY did not require antihyperglycemic therapy except sometimes during pregnancy, so they are important to establish MODY subtype-specific CSS and correctly determine the subtype of MODY. Certainly, the CSS for the full range of MODY was also useful if only basic clinical characteristics were available. A clinical prediction model to determine the probability of common MODY subtypes (GCK-MODY, HNF1A/4A-MODY) had been developed.⁸ Using that model, candidate patients could be picked up, and then separate gene for subtype of MODY need to be sequenced if they meet CSS for single subtype of MODY or have additional clinical characteristics.

Diabetes due to variants of mitochondrial DNA, *GCK*, *HNF4A* or *HNF1B* was also common among monogenic diabetes. To avoid the potential interference from other subtype monogenic diabetes, we excluded four patients (two mitochondrial 3243A>G, one *GCK-MODY* and one *HNF4A-MODY*) in evaluating the effectiveness of the *HNF1A*-CSS. Of them, only the patient with *GCK*-MODY met our *HNF1A*-CSS, but because he had stable mild hyperglycemia, it was easy to pick him up as *GCK*-MODY.¹⁰

Our study has some limitations. (1) This HNF1A-CSS was established for a selected Chinese population of young patients with EOD and must be tested in other populations. (2) Although the sample size of the patients with HNF1A-MODY in this study is the largest sample of Asians reported to date, it is not large enough to obtain the best predictive model for HNF1A-MODY. Cut-off values with the highest sensitivity reported in previous studies and in our analysis were selected, which was verified feasible in the study, to improve the performance of HNF1A-CSS. (3) Indeed, a fasting C peptide level would be helpful to define the insulin resistance. Unfortunately, we did not detect the serum C peptide in cohort 2 and some patients in cohort 1 and cohort 3. (4) Because only the patients carrying pathogenic or likely pathogenic variants of HNF1A were diagnosed with HNF1A-MODY in this study, it is possible that those patients with uncertain significance variants of HNF1A (eg, p.Gly47Arg and

copyright

p.Glu548Gly in online supplementary table S2) were misdiagnosed as EOD, future functional studies and other information help to further evaluate the pathogenicity of these variants. (5) Due to the limited sensitivity of these autoantibody kits, we are not able to exclude all the patients with T1DM, although GADA, ICA and IA were analyzed in this study. However, basing on the fact of low prevalence of T1DM in China, we thought the effect was ignorable.

In summary, we found a *HNF1A*-CSS which combining hs-CRP, BMI, HDL-c and Fins levels is very useful for distinguishing patients with *HNF1A*-MODY from young patients with EOD in a Chinese population. This screening strategy is practical and easy to use in a wide variety of settings in the real world.

Author affiliations

¹Department of Endocrinology and Metabolism, Peking University People's Hospital, Peking University Diabetes Center, Beijing, China

²Department of Endocrinology and Metabolism, Beijing Airport Hospital, Beijing, China

³Department of Endocrinology, Peking Union Medical College Hospital, Diabetes Research Center of Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China

⁴Department of Endocrinology, Chinese PLA General Hospital, Beijing, China
⁵Department of Endocrinology, Beijing Tong Ren Hospital, Capital Medical University, Beijing, China

⁶Department of Endocrinology, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China

⁷Department of Endocrinology and Metabolism, Beijing Tsinghua Changgung Hospital, Tsinghua University, Beijing, China

⁸Department of Endocrinology and Metabolism, Beijing Pinggu Hospital, Beijing, China

⁹Department of Endocrinology, Beijing Liangxiang Hospital, Capital Medical University, Beijing, China

Acknowledgements We thank all the research staff for collecting data and blood samples. We also thank all study participants for their participation and contributions.

Contributors YM performed a literature review, collected data, analyzed the data, and wrote the manuscript. LJ and XH designed the study, contributed to the discussion, and reviewed and edited the manuscript. SG, XW, XC, XX, WG, JY, LZhon, JX, ML,WL, SZ, XZho, YLi, LZhou, YZ, YLu, QR, XHu, XG, XZha, RZ, LC, FW, QW and MH help collect the data and reviewed the manuscript. YM, XHa and LJ are the guarantors of this work and, as such, had full access to all the data reported in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. All authors approved the final draft.

Funding This research was supported by grants from the National Key Research and Development Program (2016YFC1304901) and the Beijing Municipal Commission of Science and Technology funds (Z141100007414002 and D131100005313008).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study was performed in accordance with the tenets of the Declaration of Helsinki and approved by the ethics committee of Peking University People's Hospital. Written informed consent was obtained from all participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work noncommercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: $\true{http://creativecommons.org/licenses/by-nc/4.0/.}$

ORCID iDs

Yumin Ma http://orcid.org/0000-0002-4992-4365 Meng Li http://orcid.org/0000-0003-3908-4830 Ling Chen http://orcid.org/0000-0001-9639-2691

REFERENCES

- 1 Pihoker C, Gilliam LK, Ellard S, *et al.* Prevalence, characteristics and clinical diagnosis of maturity onset diabetes of the young due to mutations in HNF1A, HNF4A, and glucokinase: results from the search for diabetes in youth. *J Clin Endocrinol Metab* 2013;98:4055–62.
- 2 Ellard SB-C C, Hattersley AT. European molecular genetics quality network, MODY group: best practice guidelines for the molecular genetic diagnosis of maturity-onset diabetes of the young. *Diabetologia* 2008;51:546–53.
- 3 Gardner DS, Tai ES. Clinical features and treatment of maturity onset diabetes of the young (MODY). *Diabetes Metab Syndr Obes* 2012;5:101–8.
- 4 Isomaa B, Henricsson M, Lehto M, et al. Chronic diabetic complications in patients with MODY3 diabetes. *Diabetologia* 1998;41:467–73.
- 5 Pearson ER, Liddell WG, Shepherd M, *et al*. Sensitivity to sulphonylureas in patients with hepatocyte nuclear factor-1alpha gene mutations: evidence for pharmacogenetics in diabetes. *Diabet Med* 2000;17:543–5.
- 6 Shepherd M, Shields B, Ellard S, et al. A genetic diagnosis of HNF1A diabetes alters treatment and improves glycaemic control in the majority of insulin-treated patients. *Diabet Med* 2009;26:437–41.
- 7 Bellanné-Chantelot C, Lévy DJ, Carette C, et al. Clinical characteristics and diagnostic criteria of maturity-onset diabetes of the young (MODY) due to molecular anomalies of the HNF1A gene. J Clin Endocrinol Metab 2011;96:E1346–51.
- 8 Shields BM, McDonald TJ, Ellard S, et al. The development and validation of a clinical prediction model to determine the probability of MODY in patients with young-onset diabetes. *Diabetologia* 2012;55:1265–72.
- 9 McDonald TJ, Shields BM, Lawry J, *et al.* High-sensitivity CRP discriminates HNF1A-MODY from other subtypes of diabetes. *Diabetes Care* 2011;34:1860–2.
- 10 Ma Y, Han X, Zhou X, et al. A new clinical screening strategy and prevalence estimation for glucokinase variant-induced diabetes in an adult Chinese population. Genet Med 2019;21:939–47.
- 11 Shields BM, Hicks S, Shepherd MH, et al. Maturity-Onset diabetes of the young (MODY): how many cases are we missing? *Diabetologia* 2010;53:2504–8.
- 12 Chambers C, Fouts A, Dong F, *et al.* Characteristics of maturity onset diabetes of the young in a large diabetes center. *Pediatr Diabetes* 2016;17:360–7.
- 13 Grzanka M, Matejko B, Szopa M, et al. Assessment of Newly Proposed Clinical Criteria to Identify HNF1A MODY in Patients with an Initial Diagnosis of Type 1 or Type 2 Diabetes Mellitus. Adv Med 2016;2016:1–3.
- 14 IDF. IDF consensus worldwide definition of the metabolic syndrome, 2017.
- 15 Bergman RN, Ader M, Huecking K, et al. Accurate assessment of beta-cell function: the hyperbolic correction. *Diabetes* 2002;51 Suppl 1:S212–20.
- 16 Zhou B-F, Cooperative Meta-Analysis Group of the Working Group on Obesity in China. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults--study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci* 2002;15:83–96.
- 17 Li H, Ji C-Y, Zong X-N, et al. [Height and weight standardized growth charts for Chinese children and adolescents aged 0 to 18 years]. Zhonghua Er Ke Za Zhi 2009;47:487–92.
- 18 Alberti KG, Zimmet PZ, Definition Zimmet PZ: Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a who consultation. *Diabet Med* 1998;15:539–53.
- 19 Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and genomics and the association for molecular pathology. *Genet Med* 2015;17:405–23.

Genetics/Genomes/Proteomics/Metabolomics

- 20 Ellard S, Lango Allen H, De Franco E, *et al.* Improved genetic testing for monogenic diabetes using targeted next-generation sequencing. *Diabetologia* 2013;56:1958–63.
 21 Diabetologia 2013;56:1958–63.
 - 21 Owen KR. Rd Lawrence lecture 2012: assessing aetiology in diabetes: how C-peptide, CRP and fucosylation came to the Party! Diabet Med 2013;30:260–6.
 - 22 Carroll RW, Murphy R. Monogenic diabetes: a diagnostic algorithm for clinicians. *Genes* 2013;4:522–35.
 - 23 Kleinberger JW, Copeland KC, Gandica RG, et al. Monogenic diabetes in overweight and obese youth diagnosed with type 2 diabetes: the today clinical trial. *Genet Med* 2018;20:583–90.
 - 24 Owen KR, Thanabalasingham G, James TJ, et al. Assessment of high-sensitivity C-reactive protein levels as diagnostic discriminator of maturity-onset diabetes of the young due to HNF1A mutations. Diabetes Care 2010;33:1919–24.
 - 25 Thanabalasingham G, Shah N, Vaxillaire M, et al. A large multi-centre European study validates high-sensitivity C-reactive protein (hsCRP) as a clinical biomarker for the diagnosis of diabetes subtypes. *Diabetologia* 2011;54:2801–10.

- 26 McDonald TJ, McEneny J, Pearson ER, et al. Lipoprotein composition in HNF1A-MODY: differentiating between HNF1A-MODY and type 2 diabetes. *Clin Chim Acta* 2012;413:927–32.
- 27 Bellanne-Chantelot C, Coste J, Ciangura C, et al. Monogenic diabetes Study group of the Societe Francophone Du D: highsensitivity C-reactive protein does not improve the differential diagnosis of HNF1A-MODY and familial young-onset type 2 diabetes: a grey zone analysis. *Diabetes & metabolism* 2016;42:33–7.
- 28 Xu JY, Dan QH, Chan V, *et al.* Genetic and clinical characteristics of maturity-onset diabetes of the young in Chinese patients. *Eur J Hum Genet* 2005;13:422–7.
- 29 Shih DQ, Bussen M, Sehayek E, et al. Hepatocyte nuclear factor-1alpha is an essential regulator of bile acid and plasma cholesterol metabolism. Nat Genet 2001;27:375–82.
- 30 Fajans SS, Bell GI, Polonsky KS. Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. N Engl J Med 2001;345:971–80.