

# Weight Management in Phenylketonuria: What Should Be Monitored?

Julio César Rocha<sup>a–c</sup> Margreet van Rijn<sup>d</sup> Esther van Dam<sup>d</sup> Kirsten Ahring<sup>e</sup>  
Amaya Bélanger-Quintana<sup>f</sup> Katharina Dokoupil<sup>g</sup> Hulya Gokmen Ozel<sup>h</sup>  
Anna Maria Lammardo<sup>i</sup> Martine Robert<sup>j</sup> Carina Heidenborg<sup>k</sup>  
Anita MacDonald<sup>l</sup>

<sup>a</sup>Centro de Genética Médica Doutor Jacinto de Magalhães, CHP EPE, <sup>b</sup>Faculdade de Ciências da Saúde, Universidade Fernando Pessoa, and <sup>c</sup>Center for Health Technology and Services Research (CINTESIS), Porto, Portugal; <sup>d</sup>University of Groningen, University Medical Center Groningen, Beatrix Children's Hospital, Groningen, The Netherlands; <sup>e</sup>Department of PKU, Kennedy Centre, Copenhagen University Hospital, Glostrup, Denmark; <sup>f</sup>Unidad de Enfermedades Metabólicas Servicio de Pediatría, Hospital Ramon y Cajal, Madrid, Spain; <sup>g</sup>Department of Metabolism and Nutrition, Dr. von Hauner Children's Hospital, University of Munich, Munich, Germany; <sup>h</sup>Department of Nutrition and Dietetics, Faculty of Health Sciences, Hacettepe University, Ankara, Turkey; <sup>i</sup>Department of Pediatrics, San Paolo Hospital University of Milan, Milan, Italy; <sup>j</sup>Nutrition and Metabolism Unit, Hôpital Universitaire des Enfants Reine Fabiola, Brussels, Belgium; <sup>k</sup>Department of Clinical Nutrition and Dietetics, Karolinska University Hospital, Stockholm, Sweden; <sup>l</sup>The Children's Hospital, Birmingham, UK

## Key Words

Overweight · Obesity · Body composition · Comorbidities · Phenylketonuria · Diet

## Abstract

**Background:** Severe intellectual disability and growth impairment have been overcome by the success of early and continuous treatment of patients with phenylketonuria (PKU). However, there are some reports of obesity, particularly in women, suggesting that this may be an important comorbidity in PKU. It is becoming evident that in addition to acceptable blood phenylalanine control, metabolic dieticians should regard weight management as part of routine clinical practice. **Summary:** It is important for practitioners to differentiate the 3 levels for overweight interpretation: anthropometry, body composition and frequency and severity of associated metabolic comorbidities. The main objectives of this review are to suggest proposals for the mini-

mal standard and gold standard for the assessment of weight management in PKU. While the former aims to underline the importance of nutritional status evaluation in every specialized clinic, the second objective is important in establishing an understanding of the breadth of overweight and obesity in PKU in Europe. **Key Messages:** In PKU, the importance of adopting a European nutritional management strategy on weight management is highlighted in order to optimize long-term health outcomes in patients with PKU.

© 2015 S. Karger AG, Basel

## Introduction

Historically, the main objective of phenylketonuria (PKU) management was to prevent profound and irreversible intellectual disability together with the avoidance of nutritional deficiencies resulting in growth impairment [1]. Traditional low phenylalanine dietary manage-

**Table 1.** International BMI cut-offs for classifying overweight and obesity according to different definitions

| Criteria               | Age group  | Risk of overweight | Overweight                                                         | Obesity                                                            |
|------------------------|------------|--------------------|--------------------------------------------------------------------|--------------------------------------------------------------------|
| WHO growth charts [8]  | <5 years   | 1 SD               | 2 SD                                                               | 3 SD                                                               |
| WHO growth charts [9]  | 5–19 years | –                  | 1 SD                                                               | 2 SD                                                               |
| CDC growth charts [12] | 2–20 years | –                  | 1 SD                                                               | 2 SD                                                               |
| IOTF [11]              | 2–18 years | –                  | Cut-off corresponding to a BMI of 25 kg/m <sup>2</sup> at 18 years | Cut-off corresponding to a BMI of 30 kg/m <sup>2</sup> at 18 years |
| WHO adults [6]         | Adults     | –                  | 25 ≤ BMI <30 kg/m <sup>2</sup>                                     | BMI ≥30 kg/m <sup>2</sup>                                          |

ment, and more recently with pharmacological treatment, has been very successful in achieving these goals. However, there is concern that obesity may be a common comorbidity. Although in PKU there is still limited evidence, some studies highlight an increased obesity risk particularly in females [2–4]. It is therefore essential that PKU health professionals are able to conduct lifestyle ‘risk assessments’ to identify potential obesogenic factors, consider trends in nutritional status by systematically monitoring key clinical and biochemical measures, and provide effective nutritional and lifestyle advice in order to avoid obesity. Also, it is important to standardize procedures for data collection in order to monitor trends between centres and countries. This will help clarify if overweight is related to the disorder, the treatment or to a disrupted energy balance.

Therefore, this paper has 3 objectives: (1) to clarify definition and interpretation of measurements for overweight and obesity; (2) to define minimal standards for monitoring overweight and obesity in the PKU clinic; and (3) to propose a ‘gold standard’ for data collection for assessment of overweight, obesity and lifestyle in PKU.

### Defining and Assessing Overweight and Obesity

Adequate assessment depends on age and gender, and careful interpretation is needed particularly in childhood and adolescence, where the correct diagnosis of overweight and obesity is more challenging than in adulthood. Several methods exist for assessing overweight and obesity, all with their pros and cons.

#### Body Mass Index

Body mass index (BMI) has been used in paediatrics since the 1980s [5] and defines the widely used criteria for overweight and obesity in adults (BMI ≥25 kg/m<sup>2</sup>

corresponds to overweight and BMI ≥30 kg/m<sup>2</sup> corresponds to obesity) [6]. The preference for its use is from the association with body composition and disease risk factors, as well as being calculated by weight and height measurements only [7]. In paediatrics, comparison with a reference population is required. Several countries have their own nationally representative reference population growth charts [7]. However, the World Health Organization (WHO) growth charts (0–5 years of age) represent a standard of physiological growth, since breast-fed children from controlled pregnancies were included [8] and they have also proposed references for children aged 5–19 years [9]. Other international criteria for diagnosing overweight and obesity in paediatrics are available from the International Obesity Task Force (IOTF). Although this can be used for international descriptive and comparative studies, it is not designed for clinical use and so should not replace the national reference data [10]. Nevertheless, IOTF cut-offs expressed as centiles permit comparisons with other international criteria like WHO [11]. Table 1 summarizes internationally available criteria.

Caution is needed when interpreting results from studies using different criteria to classify overweight and obese individuals, for example, IOTF, WHO or other national data, like Centers for Disease Control and Prevention (CDC) [12], since variable results are expected [13]. In addition, more research is needed to better identify universal BMI cut-offs and its association with health outcomes later in life [14].

#### Body Composition

Although BMI is the most commonly used marker for diagnosing overweight and obesity, it should be highlighted that overweight definition is based on the proportion of body fat to the body weight, which results from increased energy intake, reduced energy expendi-

**Table 2.** Minimal standards for monitoring overweight and obesity in the PKU clinic

| Measure                              | Justification                                                          | Practicalities/check-points                                                                                                                 |
|--------------------------------------|------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Dietary intake and life-style</i> |                                                                        |                                                                                                                                             |
| Diet                                 | Diet will modulate the risk of developing overweight and obesity       | Total protein, amino acid and natural protein intakes<br>Total energy intakes<br>BH4-treated patients analysed separately<br>Diet practices |
| Social and life-style                | Life-style will modulate the risk of developing overweight and obesity | Daily main occupation: student, employed or no occupation<br>Practice of structured exercise: yes or no                                     |
| <i>Anthropometry</i>                 |                                                                        |                                                                                                                                             |
| BMI                                  | Good epidemiological marker                                            | Weight and height should be measured with the same equipment and at the same moment of the day (morning or afternoon)                       |
| <i>Clinical biochemistry</i>         |                                                                        |                                                                                                                                             |
| Glucose and lipid metabolism         | Will give a picture about the severity of overweight                   | Fasting blood glucose and blood lipids: total cholesterol, HDL, LDL, triglycerides                                                          |

ture or both [15]. According to WHO, the ‘real or clinic’ overweight is perceived when the adiposity level is sufficiently high to increase morbidity and mortality [15]. Although considered a good screening tool, BMI may not always identify the overfat individuals (increased fat mass percentage), since a stunted linear growth may be responsible for the increased BMI [16]. Even in adults, a rigorous assessment and careful interpretation between BMI and health outcome are needed, since these relationships are not always clear [17]. Considering the wide range of methods available for measuring body composition, it is important to refer that these start with the predictive techniques (skinfold thicknesses and bio-electrical impedance analysis), then reference methods like dual energy X-ray absorptiometry, isotope dilution and air displacement plethysmography, finalizing with the gold standard methods (4-component model for quantifying fat, water, mineral and protein and magnetic resonance imaging for quantifying fat distribution) [18].

#### *Metabolic Comorbidities*

The relationship between BMI and disease is mediated by a metabolically abnormal phenotype mainly characterized by dyslipidemia, hypertension, insulin resistance and inflammation, all coexisting with increased abdominal obesity [19, 20]. So, in addition to ‘overweight’, it is increasingly important to define overfat individuals and particularly those at increased risk of metabolic disturbances. Further, considering that visceral adipose tissue assumes great relevance in terms of the inflamma-

tory impact of the adipose tissue between metabolic syndrome and allied comorbidities [21, 22], understanding the main location of body fat deserves clinical attention. Further study is particularly required to investigate if overweight in PKU is associated with the same abnormal metabolic patterns usually seen in the general population [23].

#### **Assessment of Overweight, Obesity and Lifestyle in Patients with PKU**

Published results on overweight and obesity in PKU patients are contradictory; most of them are only representative of single PKU centres and they mainly include children and adolescents [23]. Not all studies demonstrate overweight in patients with PKU [2, 4, 24–27], but a developing trend suggests that overweight in PKU is following the general population [2, 4, 26, 27]. However, this outcome should not be commended, as PKU health care professionals should also focus on general health and prevention of overweight is essential.

#### *Minimal Standards for Monitoring Overweight and Obesity in the PKU Clinic*

Minimal standards for monitoring overweight and obesity are proposed in PKU (table 2).

Delaying diagnosis and treatment of overweight will reduce the chances of a successful intervention. The concept of diet for life adopted in PKU management [31]

**Table 3.** Proposed ‘gold standard’ assessment for data collection in PKU

| Measure                                   | Justification                                                                                   | Practicalities/check-points                                                                                                                                                                                                                                                                                                                                                          |
|-------------------------------------------|-------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Dietary intake and life-style</i>      |                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                      |
| Diet                                      | Diet will modulate the risk of developing overweight and obesity                                | Total protein, amino acid and natural protein intakes<br>Total energy intakes and % of calories from fat and carbohydrate<br>Carbohydrate intakes, mainly amounts of mono and disaccharide ingestion<br>BH4-treated patients analysed separately<br>BH4-treated patients diet patterns detailed<br>Low protein food intake: types, categories and % of the total daily energy intake |
| Social and life-style                     | Life-style will modulate the risk of developing overweight and obesity                          | Daily main occupation: student, employed or no occupation<br>Frequency, duration and intensity of physical activity and exercise activities<br>Eating patterns behaviour; frequency of snacks<br>Patient/caregiver knowledge about healthy foods in PKU                                                                                                                              |
| <i>Anthropometry and body composition</i> |                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                      |
| BMI                                       | Good epidemiological marker                                                                     | Need to use standard methods and calibrated equipment's<br>Weight and height should be measured with the same equipment and at the same moment of the day (morning or afternoon)                                                                                                                                                                                                     |
| Waist circumference                       | Good association with abdominal adiposity                                                       | Need to use standard technique<br>Need to have trained health care professionals                                                                                                                                                                                                                                                                                                     |
| Body composition                          | To differentiate overweight from overfat and to help nutritional status interpretation          | To select an available method in each clinic<br>To follow standardized procedure in every measurements in order to obtain at least body fat %                                                                                                                                                                                                                                        |
| <i>Clinical biochemistry</i>              |                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                      |
| Glucose metabolism                        | Insulin resistance is on the basis of metabolic syndrome                                        | To measure fasting blood glucose<br>Add A1C haemoglobin and insulin, especially when insulin resistance is suspected to occur (presence of visceral obesity)                                                                                                                                                                                                                         |
| Lipid metabolism                          | Dislipidemia is an important feature of metabolic syndrome being linked with insulin resistance | To measure blood total cholesterol, HDL, LDL, triglycerides<br>Need to control the fasting time period in order to have reliable markers<br>Apolipoproteins measurement would be recommended                                                                                                                                                                                         |
| Inflammation                              | Inflammatory markers are associated with abdominal obesity                                      | To measure c-reactive protein, especially high-sensitivity c-reactive protein<br>When possible, cytokines could be monitored, especially in the presence of visceral obesity                                                                                                                                                                                                         |
| <i>Other clinical data</i>                |                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                      |
| Blood pressure                            | It is a metabolic syndrome component well associated with the global cardiovascular risk        | Use standard techniques and procedures<br>The presence of the ‘white coat effect’ should be recognized<br>Careful interpretation is needed in paediatrics                                                                                                                                                                                                                            |

should guide health care professionals into caring for all aspects of health and not solely focused on phenylalanine levels.

#### *Proposed ‘Gold Standard’ Assessment for Data Collection in PKU*

In PKU, in order to understand the European prevalence of overweight and obesity, it is important to assess and collect data in a standardized manner. Table 3 describes a proposed ‘gold standard assessment’ tool. From the beginning, it is important to differentiate between

cross-sectional and longitudinal data. Cross-sectional data will give an indication of the population under immediate follow-up; longitudinal data will demonstrate the evolution of overweight and obesity and a careful interpretation of data will allow the identification of critical periods, helping to clarify if specific PKU treatment strategies have an obesogenic effect. Some studies have already identified adolescence, especially in females [2, 4, 28], and early infancy [29], as critical periods, thereby requiring particular attention. Preventing an early BMI rebound [29] and an understanding of

**Table 4.** Patients', families' and other longitudinal data to be collected in patients with PKU

| Measure                     | Justification                                                                                  | Practicalities/check-points                                                                                                                                                                                                                                                                                                      |
|-----------------------------|------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Patient-related data</i> |                                                                                                |                                                                                                                                                                                                                                                                                                                                  |
| Disorder                    | Disorder severity may affect metabolism, dietary restrictions, adherence and metabolic control | Classification in terms of HPA, mild PKU or classical PKU<br>Late treated patients should be analysed separately                                                                                                                                                                                                                 |
| Early health and nutrition  | Early health and nutrition are important determinants of overweight development later in life  | Birth weight<br>Breastfeeding length<br>Amount of total protein intake in the first 1–2 years of life<br>Degree of prematurity<br>Weight gain pattern in the first years of life<br>Age at BMI rebound<br>Age at menarche<br>Other disorders/comorbidities and/or exposure to chronic anti-biotherapy in the first years of life |
| <i>Family-related data</i>  |                                                                                                |                                                                                                                                                                                                                                                                                                                                  |
| Nutritional status          | Family overweight is positively associated with patients tendency                              | Identify the presence of overweight or obesity in both parents<br>Are there other siblings with overweight or obesity?<br>Examine dietary pattern of the family<br>Document parental BMI and waist circumference                                                                                                                 |

feeding and lifestyle behaviour during adolescence should be beneficial [30] (see table 4 for patient and family data collection).

### Importance of Data Collection in Clinical Setting

Adopting standard procedures to monitor dietary intake, weight gain, body composition and clinical biochemistry within the clinical setting not only enables systematic monitoring of nutritional status. It will also enable the application of early preventative action to avoid the development of overweight and obesity. This should help in the provision of appropriate dietary counselling on the use of special low-protein foods; type of phenylalanine-free L-amino acid supplement; choice of natural protein sources particularly in sapropterin responsive patients who may have a higher natural protein tolerance; and also any requirement for additional exercise programmes.

### Conclusion

In PKU, nutritional status, particularly being overweight, deserves our attention as it influences long-term health status. Until now, the limited evidence within the field of overweight in PKU does not clarify if this is a serious issue in this special group of individuals. There is an

urgent need to have a systematic and standardized overweight-related data collection in European representative cohorts of PKU patients, so that different treatment interventions and approaches can be compared.

### Disclosure Statement

H.G.O., K.A., K.D., A.M.L., M.R., E.D. and J.C.R. are members of the European Nutrition Expert Panel, which is supported by Merck Serono.

C.H. was a member of the European Nutrition Expert Panel, which is supported by Merck Serono.

A.M. has received research funding and honoraria from Nutricia, Vitaflor International and Merck Serono, Member of European Nutrition Expert Panel (Merck Serono international), Member of Sapropterin Advisory Board (Merck Serono international), Member of the Advisory Board Element (Danone-Nutricia) and Arla Advisory Board.

M.R. is a Danone Research and Development consultant, Member of the Advisory Board Element (Danone-Nutricia), received financial research support of Nutricia Netherlands and received honoraria for lecturing and guideline development (Nutricia, SSIF-Excemed).

A.B.-Q. is a member of the European Nutrition Expert Panel (Merck Serono) and member of Sapropterin Advisory Board (Merck Serono).

### Sources of Funding

No specific funding for each author or for the manuscript preparation was needed.

## References

- Dokoupil K, Gokmen-Ozel H, Lammardo AM, Motzfeldt K, Robert M, Rocha JC, van Rijn M, Ahring K, Belanger-Quintana A, MacDonald A: Optimising growth in phenylketonuria: current state of the clinical evidence base. *Clin Nutr* 2012;31:16–21.
- Burrage LC, McConnell J, Haesler R, O'Riordan MA, Sutton VR, Kerr DS, McCandless SE: High prevalence of overweight and obesity in females with phenylketonuria. *Mol Genet Metab* 2012;107:43–48.
- Aldamiz-Echevarria L, Bueno MA, Couce ML, Lage S, Dalmau J, Vitoria I, Andrade F, Blasco J, Alcalde C, Gil D, Garcia MC, Gonzalez-Lamuno D, Ruiz M, Pena-Quintana L, Ruiz MA, Gonzalez D, Sanchez-Valverde F: Anthropometric characteristics and nutrition in a cohort of PAH-deficient patients. *Clin Nutr* 2014;33:702–717.
- Belanger-Quintana A, Martinez-Pardo M: Physical development in patients with phenylketonuria on dietary treatment: a retrospective study. *Mol Genet Metab* 2011;104:480–484.
- Rolland-Cachera MF, Sempe M, Guillaud-Bataille M, Patois E, Pequignot-Guggenbuhl F, Fautrad V: Adiposity indices in children. *Am J Clin Nutr* 1982;36:178–184.
- Physical status: the use and interpretation of anthropometry. Report of a WHO expert committee. *World Health Organ Tech Rep Ser* 1995;854:1–452.
- Rolland-Cachera MF: Childhood obesity: current definitions and recommendations for their use. *Int J Pediatr Obes* 2011;6:325–331.
- WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatr Suppl* 2006;450:76–85.
- de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J: Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 2007;85:660–667.
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH: Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000;320:1240–1243.
- Cole TJ, Lobstein T: Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 2012;7:284–294.
- Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, Mei Z, Curtin LR, Roche AF, Johnson CL: CDC growth charts: United States. *Adv Data* 2000;314:1–27.
- Shields M, Tremblay MS: Canadian childhood obesity estimates based on WHO, IOTF and CDC cut-points. *Int J Pediatr Obes* 2010;5:265–273.
- Rolland-Cachera MF: Towards a simplified definition of childhood obesity? A focus on the extended IOTF references. *Pediatr Obes* 2012;7:259–260.
- Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000;894:i–xii, 1–253.
- Wells JC, Fewtrell MS: Measuring body composition. *Arch Dis Child* 2006;91:612–617.
- Flegal KM, Kit BK, Orpana H, Graubard BI: Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA* 2013;309:71–82.
- Wells JC: Body composition in infants: evidence for developmental programming and techniques for measurement. *Rev Endocr Metab Disord* 2012;13:93–101.
- Oliveros E, Somers VK, Sochor O, Goel K, Lopez-Jimenez F: The concept of normal weight obesity. *Prog Cardiovasc Dis* 2014;56:426–433.
- Despres JP, Lemieux I: Abdominal obesity and metabolic syndrome. *Nature* 2006;444:881–887.
- Despres JP: Abdominal obesity and cardiovascular disease: is inflammation the missing link? *Can J Cardiol* 2012;28:642–652.
- Ouchi N, Parker JL, Lugus JJ, Walsh K: Adipokines in inflammation and metabolic disease. *Nat Rev Immunol* 2011;11:85–97.
- Rocha JC, MacDonald A, Trefz F: Is overweight an issue in phenylketonuria? *Mol Genet Metab* 2013;110(suppl):S18–S24.
- Dobbelaere D, Michaud L, Debrabander A, Vanderbecken S, Gottrand F, Turck D, Farriaux JP: Evaluation of nutritional status and pathophysiology of growth retardation in patients with phenylketonuria. *J Inherit Metab Dis* 2003;26:1–11.
- Huemer M, Huemer C, Moslinger D, Huter D, Stockler-Ipsiroglu S: Growth and body composition in children with classical phenylketonuria: results in 34 patients and review of the literature. *J Inherit Metab Dis* 2007;30:694–699.
- Rocha JC, van Spronsen FJ, Almeida MF, Soares G, Quelhas D, Ramos E, Guimaraes JT, Borges N: Dietary treatment in phenylketonuria does not lead to increased risk of obesity or metabolic syndrome. *Mol Genet Metab* 2012;107:659–663.
- Robertson LV, McStravick N, Ripley S, Weetch E, Donald S, Adam S, Micciche A, Boocock S, MacDonald A: Body mass index in adult patients with diet-treated phenylketonuria. *J Hum Nutr Diet* 2013;26(suppl 1):1–6.
- Albersen M, Bonthuis M, de Roos NM, van den Hurk DA, Carbasius Weber E, Hendriks MM, de Sain-van der Velden MG, de Koning TJ, Visser G: Whole body composition analysis by the BodPod air-displacement plethysmography method in children with phenylketonuria shows a higher body fat percentage. *J Inherit Metab Dis* 2010;33(suppl 3):S283–S288.
- Scaglioni S, Verduci E, Fiori L, Lammardo AM, Rossi S, Radaelli G, Riva E, Giovannini M: Body mass index rebound and overweight at 8 years of age in hyperphenylalaninaemic children. *Acta Paediatr* 2004;93:1596–1600.
- Alberga AS, Sigal RJ, Goldfield G, Prud'homme D, Kenny GP: Overweight and obese teenagers: why is adolescence a critical period? *Pediatr Obes* 2012;7:261–273.
- Camp KM, Parisi MA, Acosta PB, Berry GT, Bilder DA, Blau N, Bodamer OA, Brosco JP, Brown CS, Burlina AB, Burton BK, Chang CS, Coates PM, Cunningham AC, Dobrowolski SF, Ferguson JH, Franklin TD, Frazier DM, Grange DK, Greene CL, Groft SC, Harding CO, Howell RR, Huntington KL, Hyatt-Knorr HD, Jevaji IP, Levy HL, Lichter-Konecki U, Lindegren ML, Lloyd-Puryear MA, Matalon K, MacDonald A, McPheeters ML, Mitchell JJ, Mofidi S, Moseley KD, Mueller CM, Mulberg AE, Nerurkar LS, Ogata BN, Pariser AR, Prasad S, Pridjian G, Rasmussen SA, Reddy UM, Rohr FJ, Singh RH, Sirrs SM, Stremer SE, Tagle DA, Thompson SM, Urv TK, Utz JR, van Spronsen F, Vockley J, Waisbren SE, Weglicki LS, White DA, Whitley CB, Wilfond BS, Yannicelli S, Young JM: Phenylketonuria scientific review conference: state of the science and future research needs. *Mol Genet Metab* 2014;112:87–122.