

Human pancreatic Beta-cell glucokinase: cDNA sequence and localization of the polymorphic gene to chromosome 7, band p 13

S. Nishi^{1,3}, M. Stoffel¹, K. Xiang¹, T. B. Shows², G. I. Bell¹ and J. Takeda¹

¹ Howard Hughes Medical Institute, and Departments of Biochemistry and Molecular Biology, and Medicine, University of Chicago, Chicago, Illinois, ² Department of Human Genetics, Roswell Park Memorial Institute, New York State Department of Health, Buffalo, New York, USA, and ³ Second Division of Internal Medicine, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan

Summary. The glucose phosphorylating enzyme glucokinase plays an important role in the regulation of glucose homeostasis. Studies in rodents indicate that pancreatic Beta cells and hepatocytes express different isoforms of this protein as a consequence of the presence of tissue-specific promoters and exon 1 sequences which are spliced to a shared group of nine exons which encode most of the mRNA and protein. Here, we report the isolation and characterization of cDNA clones encoding human Beta-cell glucokinase. The sequence of human Beta-cell glucokinase shows 97% amino acid identity with that of the cognate rat protein. We also mapped the human glucokinase gene to the short arm of chromosome 7 by analysing its segregation in a panel of reduced human-

mouse somatic cell hybrids. In situ hybridization to metaphase chromosomes confirmed the localization of the human glucokinase gene to chromosome 7 and indicated that it was in band p 13. A microsatellite DNA polymorphism that can be typed using the polymerase chain reaction was identified upstream of exon 1 a, the Beta-cell specific first exon. The glucokinase cDNA clone and highly informative DNA polymorphism will be useful for examining the role of this gene in the pathogenesis of diabetes mellitus.

Key words: Glucokinase, polymerase chain reaction, microsatellite DNA polymorphism.

Glucokinase (ATP: D-hexose 6-phosphotransferase, EC 2.7.1.1) is expressed by hepatocytes and pancreatic Beta cells and plays a key role in the regulation of glucose homeostasis [1, 2]. In the hepatocyte, the phosphorylation of glucose by glucokinase facilitates the uptake and metabolism of glucose by maintaining a gradient for glucose transport into these cells. In Beta cells, glucokinase is believed to comprise part of the glucose sensing mechanism which regulates insulin secretion. In the rat, glucokinase is encoded by a single-copy gene that has 11 exons [1, 3, 4]. The glucokinase transcripts present in the liver and Beta-cell share sequences encoded by the last nine exons. The sequences of the liver and Beta-cell glucokinase mRNA differ at their 5'-ends because the glucokinase gene has two promoters, one of which functions in the Beta cell and the other which is active in liver. As a consequence, the sequences of the 5'-untranslated regions of the two transcripts differ. In addition, since translation is initiated within the tissue-specific first exon, the sequence of amino acids 1–15 of the liver and Beta-cell isoforms are different. The presence of tissue-specific alternative promoters allows the glucokinase gene to be differentially regulated in these two tissues.

Because of the important role played by glucokinase in the regulation of insulin secretion and the uptake of glucose by the liver, it has been implicated as a candidate gene whose genetic variation or altered regulation could contribute to the development of Type 2 (non-insulin-dependent) diabetes mellitus. Tanizawa et al. [5] have recently described the isolation and characterization of cDNA clones encoding the hepatic isoform of human glucokinase. In this report, we present the sequence of the human Beta-cell isoform. In addition, we have mapped the human glucokinase gene (GCK) to chromosome 7, band p13, and identified a microsatellite DNA polymorphism that will facilitate genetic studies of its role in the development of Type 2 diabetes.

Materials and methods

General methods

Standard methods were carried out as described in Sambrook et al. [6] and as described previously [7]. DNA sequencing was done by the dideoxynucleotide chain-termination procedure after subcloning appropriate DNA fragments into M13mp18 or M13mp19. The sequence was confirmed on both strands.

Fig. 1. Composite nucleotide sequence of human Beta-cell glucokinase cDNA and predicted amino acid sequence of the protein. The sequence encoded by the Beta-cell-specific exon 1a is shown in bold-face type. The number of the nucleotide at the end of each line is noted. The sequence was obtained from the following clones: hGK-p5, nucleotides 1-393; hGK3.1, nucleotides 372-831; and hGK12-1, nucleotides 771-2606. The corresponding amino acid residue of rat Beta-cell glucokinase [4, 14] is indicated above that of the human sequence at those sites at which the sequences differ. There are three differences between the cDNA sequence presented here and the sequence of human liver glucokinase [15] in the region in which these two sequences overlap—Codon 107 is ATG (Met)

here, in the human glucokinase gene (unpublished) and in the rat glucokinase cDNA sequence [14], and ACG (Thr) in Matsutani et al. [15]; and there are two differences in the 3'-untranslated region (underlined) which are G and C in our sequence and GG and CC, respectively, in Reference 15. We show the sequence of codon 74 as being TTC (Phe) which is also the sequence in Reference 15 and in the gene (unpublished). However, in two independent polymerase chain reaction products that were obtained from this region it was TCC (Ser). We assume that this difference is a consequence of misincorporation by either reverse transcriptase or Taq DNA polymerase. It is unknown if the sequence differences noted above represent polymorphisms in the human glucokinase gene




Fig. 2. Ideogram of human chromosome 7 showing silver grain distribution after hybridization with phGK12-1. One hundred metaphase spreads were examined. Of the total number of grains, 27.1 % (65 of 240) were on chromosome 7, and of these, 36.9 % (24 of 65) were at 7p13→p14. Eighteen percent of metaphase spreads examined showed a silver grain at 7p13→p14

Isolation of human glucokinase cDNA clones

A human liver cDNA library was screened using low-stringency hybridization conditions [8] (hybridization conditions – $5 \times$ SSC, 25 % formamide, $2 \times$ Denhardt's solution, 20 mmol/l sodium phosphate buffer, pH 6.5, 0.1 % sodium dodecyl sulphate, 100 μ g/ml of sonicated and denatured salmon testes DNA, 10 % dextran sulphate, and 1×10^6 cpm/ml of probe at 37 °C for 16–20 h; washing conditions – $2 \times$ SSC and 0.1 % sodium dodecyl sulphate, for 1 h each at room temperature and then at 40 °C) with a 32 P-labelled 3248 base pair (bp) EcoRI fragment of the human hexokinase I cDNA clone, λ hHEX-15 [7], which codes for amino acids 92–917 and the 3'-untranslated region of the mRNA. The cDNA clone λ hGK12-1 was obtained using this procedure. The remainder of the cDNA was obtained by reverse transcription-polymerase chain reaction (PCR) amplification of human insulinoma mRNA using specific primers [rGK-13 (sense primer) 5'-GTCGAGCAGATCCTGGCAGAG-3' and ohGK-2r (antisense primer) 5'-TGGTCCAGTTGAGAAG-GAAG-3'; the sequence of rGK-13 was based on the sequence of rat glucokinase mRNA] which gave cDNA clone – hGK3.1, and by the rapid amplification of cDNA ends (RACE) procedure [9] using human insulinoma mRNA and the specific primer (antisense) 5'-CTCTGCCAGGATCTGCTCTAC-3' which generated a cDNA clone, hGK-p5, containing the 5'-end of human Beta-cell glucokinase mRNA. At least two PCR and RACE products obtained from each amplification were sequenced to control for errors that might be introduced in the amplification by Taq DNA polymerase. A

cDNA encoding the human Beta-cell isoform of glucokinase, designated pGEM-hGK20 (vector-pGEM4Z) was generated using the three clones described above.

Gene mapping

The chromosomal location of human GCK was determined by hybridization of 32 P-labelled phGK12-1 (the insert of λ GK12-1 sub-cloned into pBR327) to Southern blots of EcoRI-digested DNA from 36 different reduced human-mouse somatic cell hybrid cell lines [10] as described previously [11]. The regional localization of GCK was determined by in situ hybridization of 3 H-labelled phGK12-1 to normal human prometaphase chromosomes as described by Nakai et al. [12].

Isolation of the human GCK gene and identification of a microsatellite DNA polymorphism

The human GCK gene was isolated from a genomic library (946203, Stratagene, La Jolla, Calif., USA) by hybridization with the insert from the cDNA clone pGEM-hGK20. Five clones [λ hGK-1, -2, -4, -5 and -7] were isolated; these clones contain all the exons of the human GCK gene. The clones containing CA-dinucleotide repeats were identified by hybridization with nick-translated 32 P-labelled poly (dA-dC)-poly (dG-dT) (Pharmacia LKB Biotechnology, Piscataway, NJ, USA) [13] (hybridization conditions – 0.5 mol/l sodium phosphate buffer, pH 7.0, 7 % sodium dodecyl sulphate, 1 % bovine serum albumin, and 1×10^6 cpm/ml of probe at 42 °C for 16–20 h; washing conditions – $1 \times$ SSC and 1 % sodium dodecyl sulphate, for 1 h each at room temperature and then at 50 °C). A 3.0 kb Hind III fragment from λ hGK-7 was isolated, digested with Sau3A I, and ligated into BamHI-digested M13mp19. The inserts in M13 clones hybridizing with 32 P-labelled poly (dA-dC)-poly (dG-dT) were sequenced.

Amplification of the microsatellite DNA polymorphism

Two primers (hGK-CA-1, 5'-AACAGATACGCTTCATCCTG-3'; and hGK-CA-2, 5'-TGTCTGCAACTTACTCTTAC-3') were used to amplify a 127–143 bp TC and AC repeat-rich region upstream of the Beta-cell specific exon, exon 1a, of human GCK. The PCR was performed using 32 P-labelled hGK-CA-1 and unlabelled hGK-CA-2. DNA was initially denatured at 94 °C for 6 min, followed by 30 cycles of denaturation at 94 °C for 1 min, annealing at 60 °C for 2 min, extension at 72 °C for 2 min and a final extension step of 10 min. The PCR products were analysed on a 5 % denaturing polyacrylamide gel. The PCR reactions were carried out in a volume of 25 μ l containing 50 mmol/l KCl, 10 mmol/l Tris-HCl (pH 8.3), 1.5 mmol/l MgCl₂, 100 μ g/ml gelatin, 200 μ mol/l each of dATP, dGTP, dCTP, and dTTP; and 0.1 μ g of DNA, 10 pmol of each primer, and 1.5 U of Taq polymerase (Perkin Elmer Cetus, Norwalk, Conn., USA).

Results

Sequence of human Beta-cell glucokinase cDNA and protein

Human glucokinase cDNA clones were isolated from a liver cDNA library by low-stringency cross-hybridization with a human hexokinase I cDNA probe. The sequence of the insert in one of these clones, now termed λ hGK12-1, did not correspond to hexokinase I. With the publication

Table 1. Characterization of microsatellite DNA polymorphism in the human glucokinase gene

Allele frequencies		Racial group		
Allele	Size (base pairs)	Caucasian (n = 29)	Asian (n = 14)	African- American (n = 13)
1	143	—	—	0.04
2	141	0.32	0.25	0.46
3	139	0.62	0.57	0.23
4	137	0.02	0.18	0.27
5	135	0.02	—	—
6	127	0.02	—	—
Heterozygosity		0.51	0.58	0.66
PIC value		0.43	0.52	0.60

PIC, polymorphic information content

tani et al. [15] reported the mapping of human GCK to chromosome 7p by linkage analysis. Our results are in complete agreement with those of these authors and, in addition, provide a precise physical localization for GCK on the short arm of chromosome 7 within band p13. Matsutani et al. [15] also described a (CA)_n repeat polymorphism in GCK. This microsatellite DNA polymorphism is different from the one that we have reported here. These two microsatellite DNA polymorphisms can be combined to make GCK more informative for linkage studies.

Froguel et al. [16] have recently reported studies showing close linkage of DNA polymorphisms in the glucokinase locus with early-onset Type 2 diabetes. The sequences of human Beta-cell glucokinase mRNA and protein described above will facilitate studies of the role of this candidate diabetes-susceptibility gene in the development of Type 2 diabetes mellitus.

Acknowledgements. This research was supported by the Howard Hughes Medical Institute and NIH Grants DK-20595, GM-20454 and HD-05196. M. Stoffel was supported by a fellowship from the Deutsche Forschungsgemeinschaft. The nucleotide sequences reported in this paper have been deposited in the GenBank data base (accession nos. M86676 and M88011).

References

1. Magnuson MA (1990) Glucokinase gene structure: functional implications of molecular genetic studies. *Diabetes* 39: 523–527
2. Matschinsky FM (1990) Glucokinase as glucose sensor and metabolic signal generator in pancreatic β -cells and hepatocytes. *Diabetes* 39: 647–652

3. Magnuson MA, Andreone TL, Printz RL, Koch S, Granner DK (1989) Rat glucokinase gene: structure and regulation by insulin. *Proc Natl Acad Sci USA* 86: 4838–4842
4. Magnuson MA, Shelton KD (1989) An alternative promoter in the glucokinase gene is active in the pancreatic β cell. *J Biol Chem* 264: 15936–15942
5. Tanizawa Y, Koranyi LI, Welling CM, Permutt MA (1991) Human liver glucokinase gene: cloning and sequence determination of two alternatively spliced cDNAs. *Proc Natl Acad Sci USA* 88: 7294–7297
6. Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning, a laboratory manual*, 2nd Ed., Cold Spring Harbor Laboratory Press Cold Spring Harbor, NY
7. Nishi S, Seino S, Bell GI (1988) Human hexokinase: sequences of amino- and carboxylterminal halves are homologous. *Biochem Biophys Res Commun* 157: 937–943
8. Fukumoto H, Seino S, Imura H et al. (1988) Sequence, tissue distribution, and chromosomal localization of mRNA encoding a human glucose transporter-like protein. *Proc Natl Acad Sci USA* 85: 5434–5438
9. Frohman MA, Dush MK, Martin GR (1988) Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. *Proc Natl Acad Sci USA* 85: 8998–9002
10. Shows T, Eddy R, Haley L et al. (1984) Interleukin-2 (IL2) is assigned to human chromosome 4. *Somatic Cell Mol Genet* 10: 315–318
11. Shows TB, Eddy RL, Byers MG et al. (1987) Polymorphic human glucose transporter gene (GLUT) is on chromosome 1p31.3→p35. *Diabetes* 36: 546–549
12. Nakai H, Byers MG, Shows TB, Taggart RT (1986) Assignment of the pepsinogen gene complex (PGA) to human chromosome region 11q13 by in situ hybridization. *Cytogenet Cell Genet* 43: 215–217
13. Litt M, Luty JA (1989) A hypervariable microsatellite revealed by in vitro amplification of a dinucleotide repeat within the cardiac actin gene. *Am J Hum Genet* 44: 397–401
14. Andreone TL, Printz RL, Pilgis SJ, Magnuson MA, Granner DK (1989) The amino acid sequence of rat liver glucokinase deduced from cloned cDNA. *J Biol Chem* 264: 363–369
15. Matsutani A, Janssen R, Donis-Keller H, Permutt MA (1992) Polymorphic (CA)_n repeat element maps the human glucokinase gene (GCK) to chromosome 7p. *Genomics* 12: 319–325
16. Froguel Ph, Vaxillaire M, Sun F et al. (1992) Close linkage of glucokinase locus on chromosome 7p to early-onset non-insulin-dependent diabetes mellitus. *Nature* 356: 162–164

Received: 28 January 1992
and in revised form: 19 March 1992

Dr. G. Bell
Howard Hughes Medical Institute
University of Chicago
5841 S. Maryland Ave., MC1028
Chicago, IL 60637
USA