

The effects of coenzyme Q₁₀ treatment on maternally inherited diabetes mellitus and deafness, and mitochondrial DNA 3243 (A to G) mutation

S. Suzuki, Y. Hinokio, M. Ohtomo, M. Hirai, A. Hirai, M. Chiba, S. Kasuga, Y. Satoh, H. Akai, T. Toyota

Third Department of Internal Medicine, Tohoku University School of Medicine, Seiryō-machi, Aoba-ku, Sendai, Japan

Summary The characteristic clinical features of diabetes mellitus with mitochondrial DNA (mtDNA) 3243(A-G) mutation are progressive insulin secretory defect, neurosensory deafness and maternal inheritance, referred to as maternally inherited diabetes mellitus and deafness (MIDD). A treatment for MIDD to improve insulin secretory defects and reduce deafness has not been established. The effects of coenzyme Q₁₀ (CoQ₁₀) treatment on insulin secretory response, hearing capacity and clinical symptoms of MIDD were investigated. 28 MIDD patients (CoQ₁₀-DM), 7 mutant subjects with impaired glucose tolerance (IGT), and 15 mutant subjects with normal glucose tolerance (NGT) were treated daily with oral administration of 150 mg of CoQ₁₀ for 3 years. Insulin secretory response, blood lactate after exercise, hearing capacity and other laboratory examinations were investigated every year. In the same way we evaluated 16 MIDD patients (control-

DM), 5 mutant IGT and 5 mutant NGT subjects in yearly examinations. The insulin secretory response assessed by glucagon-induced C-peptide secretion and 24 h urinary C-peptide excretion after 3 years in the CoQ₁₀-DM group was significantly higher than that in the control-DM group. CoQ₁₀ therapy prevented progressive hearing loss and improved blood lactate after exercise in the MIDD patients. CoQ₁₀ treatment did not affect the diabetic complications or other clinical symptoms of MIDD patients. CoQ₁₀ treatment did not affect the insulin secretory capacity of the mutant IGT and NGT subjects. There were no side effects during therapy. This is the first report demonstrating the therapeutic usefulness of CoQ₁₀ on MIDD. [Diabetologia (1998) 41: 584–588]

Keywords Deafness, mitochondrial DNA mutation, insulin, C-peptide, lactate.

Received: 16 October 1997 and in revised form: 21 January 1998

Corresponding author: Dr. S. Suzuki, M.D., Third Department of Internal Medicine, Tohoku University School of Medicine, Seiryō-machi 1–1, Aoba-ku, Sendai 980, Japan.

Abbreviations: CoQ₁₀, Coenzyme Q₁₀; IGT, impaired glucose tolerance; NGT, normal glucose tolerance; OHA, oral hypoglycaemic agents; NDR, without diabetic retinopathy; SDR, diabetic simple retinopathy; PPDR, diabetic preproliferative retinopathy; PDR, diabetic proliferative retinopathy; normo, normoalbuminuria; micro, microalbuminuria; macro, macroalbuminuria; mt DN, mitochondrial DNA; MIDD, maternally inherited diabetes and deafness; ANOVA, analysis of variance; CPR, C-peptide immunoreactivity.

In recent years, mitochondrial DNA mutations have been identified as one cause of diabetes mellitus [1–4]. The most commonly identified mtDNA mutation linked to diabetes is the single base mutation (A-G) at position 3243 within the tRNA^{LEU(UUR)} gene [3, 4]. The clinical features of diabetes mellitus with the 3243 bp mutation are maternal inheritance, progressive insulin secretory defect and a high association with neurosensory deafness. The 3243 bp mutation was identified in about 1 % of Japanese NIDDM patients [3]. Therapeutic trials with CoQ₁₀ have been reported to provide clinical benefits in patients with mitochondrial encephalomyopathy and mitochondrial gene mutations [5–14]. Recent reports have also demonstrated clinical benefits on some neurological symptoms of MIDD patients [15, 16]. However, no

Table 1. Clinical characteristics of the CoQ₁₀-treated or untreated diabetic patients, IGT and NGT subjects with mitochondrial mutation of tRNA^{LEU(UUR)} (3243)

Glucose tolerance CoQ ₁₀ treatment	DM +	IGT +	NGT +	DM –	IGT –	NGT –
Numbers of subjects	28	7	15	16	5	5
Gender (M/F)	13/15	3/4	7/8	8/8	2/3	2/3
Mutant mtDNA (%)	10.7 ± 13.0	8.7 ± 8.8	9.2 ± 9.2	11.4 ± 12.5	10.2 ± 8.7	8.1 ± 7.2
Age at trial entry	43.5 ± 12.3	38.5 ± 12.2	35.4 ± 9.8	44.4 ± 9.8	36.1 ± 8.5	35.6 ± 7.7
Age at onset of hearing loss	36.1 ± 9.4	35.4 ± 7.3	32.3 ± 5.8	37.4 ± 7.8	33.2 ± 4.4	32.5 ± 3.9
Age at onset of diabetes	38.7 ± 10.2	–	–	39.6 ± 8.7	–	–
Treatment (Diet/OHA/Insulin)	6/8/14	–	–	4/4/8	–	–
Neurosensory deafness (%)	89	43	20	81	40	20
Retinopathy (NDR/SDR/PPDR/PDR)	10/11/2/5	–	–	5/7/1/3	–	–
Nephropathy (Norm/Micro/Macro)	7/11/10	–	–	5/6/5	–	–

Abbreviations used are IGT, impaired glucose tolerance; NGT, normal glucose tolerance; OHA, oral hypoglycemic agents; NDR, without diabetic retinopathy; SDR, diabetic sim-

ple retinopathy; PPDR, diabetic preproliferative retinopathy; PDR, diabetic proliferative retinopathy; normo, normoalbuminuria; micro, microalbuminuria; macro, macroalbuminuria

treatment for MIDD to prevent insulin secretory defects and hearing loss has been established.

In this paper, we report the results of a long-term open trial of CoQ₁₀ in MIDD patients, mutant IGT and NGT subjects. We found CoQ₁₀ treatment prevented progressive insulin secretory defects, exercise intolerance and hearing loss in the MIDD patients. The molecular mechanism by which CoQ₁₀ prevents the progression of insulin secretory defects and hearing impairment is discussed.

Patients and methods

Protocol. The study protocol was approved by the Tohoku University Institutional Review Board. Informed consent was obtained from each subject. We divided 44 MIDD patients, 12 IGT subjects and 20 NGT subjects with the 3243 bp mutation into two groups; 28 MIDD patients, 7 IGT subjects (CoQ₁₀-IGT) and 15 NGT subjects (CoQ₁₀-NGT) were treated daily with oral administration of 150 mg of CoQ₁₀ (Neuquinon, Tanabe Pharmaceuticals, Osaka Japan) for 3 years. Insulin secretory response, blood lactate after exercise, hearing capacity and other laboratory examinations were investigated every year. The remaining 16 MIDD patients, 5 IGT (control-IGT) and 5 NGT subjects (control-NGT) matched for age and sex, were not treated but were evaluated in the same way in yearly examinations. The clinical characteristics of the two groups are shown in Table 1.

DNA study. Total DNA was isolated from blood and skeletal muscle. A 427 bp fragment encompassing the tRNA^{LEU(UUR)} mutation site located at nucleotide 3,243 was amplified by modified PCR using [α -³²P]-dATP (1.7 Ci · mmol⁻¹) with forward primer 5'-AAGGTTTCGTTTGTTCACGA (3,029–3,048) and reverse primer 5'-AGCGAAGGGTTGTA-GTAGCC (3,437–3,456) [17]. The radioactive fragment was digested with Apa1 for 1 h at 37°C and then followed by electrophoresis on a 5% non-denaturing polyacrylamide gel. Bands were visualized by autoradiography. The percentage of cleaved fragments was determined by densitometric analysis.

Clinical study. The insulin secretory capacity of pancreatic beta cells was evaluated with plasma C-peptide immunoreactivity

(CPR) 6 min after 1 mg of glucagon administration and urinary CPR excretion for 24 h. At diagnosis, all patients fulfilled the WHO criteria for diabetes mellitus [18]. Plasma glucose was assayed using the glucose oxidase method. Plasma insulin and CPR were assayed using a radioimmunoassay. Exercise tolerance was assessed by the blood lactate concentration after 5 W ergometer exercise for 15 min. Neurosensory hearing loss was diagnosed by expert otopharyngologists. Hearing loss was assessed by pure-tone audiometry. The pure tone average was calculated as the average of 0.5, 1, 2 and 4 kHz as described by Yamasoba [19]. The serum levels of CoQ₁₀ were determined by high-performance liquid chromatography [20].

Statistical analysis. All measurements are presented as mean ± SD. Statistical difference was assessed by a repeated measure analysis of variance (ANOVA). Wilcoxon's test was used for paired comparison, and the Mann-Whitney U-test for unpaired comparison. Statistical significance was accepted at the 95 % confidence level ($p < 0.05$).

Results

The mean serum CoQ₁₀ levels of the MIDD patients before treatment were not significantly reduced as compared with the control subjects (0.84 ± 0.25 µg/ml, vs 1.17 ± 0.29). The levels increased significantly after 3 months of therapy, but did not show any further increase at 1, 2 and 3 years (2 months of therapy, 2.76 ± 1.44 , $p > 0.01$; 1 year, 3.17 ± 2.09 , $p < 0.005$; 2 year, 3.05 ± 2.48 , $p < 0.005$; 3 year, 3.16 ± 1.99 , $p < 0.005$).

Plasma CPR was progressively reduced following glucagon in the control-DM group (before, 2.04 ± 0.72 ng/ml; after 3 years, 1.48 ± 0.48 ng/ml, $p < 0.02$), as shown in Figure 1. Plasma CPR was significantly higher following glucagon in the CoQ₁₀-DM group than that in the control-DM group after 2 years (2.53 ± 0.24 vs 1.78 ± 0.73 , $p < 0.001$) and 3 years (1.48 ± 0.48 vs 2.70 ± 0.38 , $p < 0.0001$). CoQ₁₀ treatment significantly improved the plasma CPR following glucagon in the MIDD patients (before, 2.53 ± 0.24 ; after 3 years, 2.70 ± 0.38 , $p < 0.02$).




Fig. 1. The effects of long-term CoQ₁₀ treatment on plasma C-peptide concentration 6 min after glucagon injection in MIDD patients. Open bars are shown as the control-DM group, closed bars are the CoQ₁₀-DM group. All measurements are presented as mean \pm SD. * p < 0.02, ** p < 0.001, # p < 0.0001




Fig. 2. The effects of long-term CoQ₁₀ treatment on 24 h urinary C-peptide excretion in MIDD patients. Open bars are shown as the control-DM group, closed bars are the CoQ₁₀-DM group. Results are mean \pm S.D. * p < 0.02, ** p < 0.01, # p < 0.002, ## p < 0.001

The 24-h urinary excretion of CPR gradually deteriorated in the control-DM group (before, 31.7 ± 7.0 mg/day; after 3 years, 25.2 ± 6.1 , p < 0.02), as shown in Figure 2. The urinary CPR excretion for 24 h in the CoQ₁₀-DM group was significantly higher than that in the control-DM group after 2 years (38.6 ± 3.4 vs 30.3 ± 11.9 , p < 0.01) and 3 years (49.3 ± 7.4 vs 25.2 ± 6.1 , p < 0.0001). CoQ₁₀ treatment significantly improved the urinary CPR excretion in the MIDD patients (before, 31.1 ± 9.3 ; after 2 years, 36.6 ± 3.4 , p < 0.002; after 3 years, 49.3 ± 7.4 , p < 0.001).

The blood lactate concentration after exercise gradually decreased in the CoQ₁₀-DM group (before, 24.2 ± 7.12 mg/dl; after 3 years, 16.5 ± 3.61 , p < 0.01), but was not changed in the control-DM group, as shown in Figure 3. The blood lactate concentration after exercise in the CoQ₁₀-DM group was significantly lower than that in the control-DM group after




Fig. 3. The effects of long-term CoQ₁₀ treatment on plasma lactate after exercise in the MIDD patients. Open bars are shown as the control-DM group, closed bars are the CoQ₁₀-DM group. Results are mean \pm S.D. * p < 0.05, ** p < 0.01




Fig. 4. The effects of long-term CoQ₁₀ treatment on pure-tone average in MIDD patients. Open bars are shown as the control-DM group, closed bars are the CoQ₁₀-DM group. Results are mean \pm S.D. * p < 0.05, ** p < 0.02

1 year (19.0 ± 4.82 vs 23.6 ± 3.76 , p < 0.05), 2 years (18.7 ± 5.0 vs 24.5 ± 3.27 , p < 0.05) and 3 years (16.5 ± 3.61 vs 24.4 ± 3.63 , p < 0.001).

Bilateral neurosensory deafness was demonstrated in 38 (86%) of 44 mutant diabetic patients. Hearing capacity gradually deteriorated in the control-DM group (Fig. 4). The progression rate of hearing loss ranged from 2.2 to 3.1 dB/year in the control-DM group. CoQ₁₀ treatment significantly prevented the hearing loss in the MIDD patients after 2 years treatment (40.1 ± 10.2 dB vs 50.2 ± 6.9 , p < 0.05) and after 3 years treatment (40.8 ± 9.7 vs 53.3 ± 7.9 , p < 0.02).

After 3 years, one of 5 untreated IGT subjects but none of 7 CoQ₁₀-treated IGT subjects were diagnosed as having diabetes by oral glucose tolerance test. In contrast, one of 5 untreated NGT subjects and one of 15 treated NGT subjects were also diagnosed as IGT. The insulin secretory capacity assessed by the plasma CPR after glucagon and urinary CPR excretion, blood lactate after exercise and hearing capacity were not significantly reduced in the control-IGT group and the control-NGT group after 3 years.

CoQ₁₀ treatment for 3 years did not affect the plasma CPR after glucagon, urinary CPR excretion, blood lactate after exercise and hearing capacity in the mutant IGT and NGT subjects (data not shown).

The examined 8 CoQ₁₀-MIDD, 4 control-MIDD, 4 CoQ₁₀-IGT, 3 control-IGT, 5 CoQ₁₀-NGT and 2 control-NGT had normal insulin sensitivity assessed by minimal model analysis. After 3 year trials, the CoQ₁₀ treatment did not affect insulin sensitivity in either CoQ₁₀- or control-groups (data not shown). CoQ₁₀ treatment did not affect the states of diabetic retinopathy, nephropathy, neuropathy or other diabetic chronic complications. Apart from the improvement of insulin secretory defects, there were no other significant differences in insulin requirement, metabolic control or other clinical parameters between the CoQ₁₀-DM group and the control-DM group. In addition, there were no detectable side effects from CoQ₁₀.

Discussion

In our open trial study, we confirmed that long-term CoQ₁₀ treatment prevents progressive insulin secretory defect, exercise intolerance and hearing loss in MIDD patients. However, short-term CoQ₁₀ treatment for 1–3 months did not affect the insulin secretory capacity or clinical symptoms in MIDD patients (data not shown). We also investigated the effects of long-term CoQ₁₀ treatment on the insulin secretory capacity and hearing capacity in the 12 mutant IGT and 20 mutant NGT subjects. There was only a tendency for the treatment to improve the insulin secretory capacity in the mutant IGT subjects, and we failed to demonstrate an improvement in the glucose tolerance or hearing capacity.

CoQ₁₀ is well known to act as an electron carrier of the respiratory chain in mitochondria and has been shown to improve the mutation-associated dysfunction of the respiratory chain in mitochondria. CoQ₁₀ has been tested with success in some patients with mitochondrial encephalopathy due to mitochondrial gene mutations [5–14]. A multi-centre open trial of CoQ₁₀ treatment showed that sixteen of forty-four patients with mitochondrial myopathies showed at least a 25% decrease in post-exercise lactate levels [12]. However, in the following double-blind trial of three months, the effectiveness of CoQ₁₀ treatment for mitochondrial myopathies could not be demonstrated. Thus, the effectiveness of CoQ₁₀ therapy on mitochondrial encephalopathy has not been confirmed. There are a few reports concerning the therapeutic effects on the clinical symptoms of MIDD patients. Suzuki et al. reported that CoQ₁₀ administration was effective in improving the symptoms in the legs and the residual urine volume in the bladder of a MIDD patient [15]. They also reported that the in-

sulin-induced oedema observed in MIDD patients responded to CoQ₁₀ treatment [16]. Silvestre-Aillaud et al. treated a MIDD patient with a combination of CoQ₁₀ and L-carnitine for 6 months, but could not confirm a significant effect either on insulin secretion or on insulin sensitivity [21]. Some reports described CoQ₁₀ as deficient in some patients with mitochondrial encephalomyopathy and that its replacement improved the clinical symptoms [22]. However, there was no CoQ₁₀ deficiency in the MIDD patient. Long-term CoQ₁₀ replacement increased the serum CoQ₁₀ concentrations in the MIDD patients.

In recent years, CoQ₁₀ has attracted increasing attention with regard to its function in the reduced form (ubiquinol-10) as an antioxidant. Ubiquinol-10 efficiently protects membrane phospholipids and serum low-density lipoprotein from lipid peroxidation and also mitochondrial membrane proteins and DNA from oxidative damage induced by free radicals [23]. Ubiquinol-10 is as effective in preventing oxidative damage to lipids as alpha-tocopherol, and is considered to be the best lipid-soluble antioxidant in humans. Ubiquinol-10 is known to be regenerated from CoQ₁₀ by electron transport carriers present in various biomembranes and by some enzymes. Plasma membrane DT-diaphorase has recently been proven to maintain CoQ₁₀ in its reduced antioxidant state as ubiquinol-10 and to protect membrane components from free radical damage such as lipid peroxidation [24].

Mitochondrial gene mutations are known to enhance the release of reactive oxygen species from mitochondria and induce a dysfunction in the mitochondrial respiratory chain [25], which might result in the progression of insulin secretory defects and neurosensory deafness. We speculate that the effects of CoQ₁₀ on insulin secretion, blood lactate after exercise and hearing capacity in MIDD were due to its anti-oxidation activities. A large-scale double blind cross-over study should be performed to confirm the clinical usefulness of CoQ₁₀ on MIDD.

Acknowledgements. This investigation was supported in part by a Grant for Diabetes Research from the Ministry of Health and Welfare and a grant for the Ministry of Education, Science and Culture of Japan. We gratefully acknowledge the expert assistance of Chitose Suzuki.

References

1. Ballinger SW, Shoffner JM, Hedaya EV et al. (1992) Maternally transmitted diabetes and deafness associated with a 10.4 kb mitochondrial DNA deletion. *Nature Genetics* 1: 11–15
2. Van den Ouweland JMW, Lemkes HHPJ, Ruitenbeek W et al. (1992) Mutation in mitochondrial tRNA^{LEU(UUR)} gene in a large pedigree with maternally transmitted type II diabetes mellitus and deafness. *Nature Genetics* 1: 368–371

3. Kadowaki T, Kadowaki H, Mori Y et al. (1994) A subtype of diabetes mellitus associated with a mutation of mitochondrial DNA. *New Engl J Med* 330: 962–968
4. Maassen JA, Kadowaki T (1996) Maternally inherited diabetes and deafness: a new diabetes subtype. *Diabetologia* 39: 375–382
5. Ogasahara S, Yorifuji S, Nishikawa Y et al. (1985) Improvement of abnormal pyruvate metabolism and cardiac conduction defect with coenzyme Q₁₀ in Kearns-Sayre syndrome. *Neurology* 35: 372–377
6. Ogasahara S, Yorifuji S, Nishikawa Y et al. (1986) Improvement of abnormal pyruvate metabolism and cardiac conduction defect with coenzyme Q₁₀ in Kearns-Sayre syndrome. *Neurology* 36: 45–53
7. Ogasahara S, Nishikawa Y, Yorifuji S et al. (1986) Treatment of Kearns-Sayre syndrome with coenzyme Q₁₀. *Neurology* 36: 45–53
8. Yamamoto M, Sato T, Anno M et al. (1987) Mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episodes with recurrent abdominal symptoms and coenzyme Q₁₀ administration. *J Neurol Neurosurg Psychiatry* 50: 1475–1481
9. Goda S, Hamada T, Ishimoto S et al. (1987) Clinical improvement after administration of coenzyme Q₁₀ in a patient with mitochondrial encephalomyopathy. *J Neurol* 234: 62–63
10. Nishikawa Y, Takahashi M, Yorifuji S et al. (1989) Long-term coenzyme Q₁₀ therapy for a mitochondrial encephalomyopathy with cytochrome c oxidase deficiency: a ³¹P NMR study. *Neurology* 39: 399–403
11. Zierz S, Jahns G, Jerusalem F (1989) Coenzyme Q in serum and muscle of 5 patients with Kearns-Sayre syndrome and 12 patients with ophthalmoplegia plus. *J Neurol* 236: 97–101
12. Bresolin N, Doriguzzi C, Ponzetto C et al. (1990) Ubidecarenone in the treatment of mitochondrial myopathies: a multi-center double-blind trial. *J Neurol Sci* 100: 70–78
13. Abe K, Fujimura H, Nishikawa Y et al. (1991) Marked reduction in CSF lactate and pyruvate levels after CoQ therapy in a patient with mitochondrial myopathy, encephalopathy, lactic acidosis and stroke-like episodes (MELAS). *Acta Neurol Scand* 83: 356–359
14. Chen RS, Huang CC, Chu NS (1997) Coenzyme Q₁₀ treatment in mitochondrial encephalomyopathy. *Eur Neurol* 37: 212–218
15. Suzuki Y, Kadowaki H, Atsumi Y et al. (1995) A case of diabetic amyotrophy associated with 3243 mitochondrial tRNA^{Leu(UUR)} mutation and successful therapy with coenzyme Q₁₀. *Endocr J* 42: 141–145
16. Suzuki Y, Kadowaki H, Taniyama M et al. (1995) Insulin edema in diabetes mellitus associated with the 3243 mitochondrial tRNA^{Leu(UUR)} mutation; Case reports. *Diabetes Res Clin Pract* 29: 137–142
17. Suzuki S, Hinokio Y, Hirai S et al. (1994) Pancreatic beta-cell secretory defect associated with mitochondrial point mutation of the tRNA^{Leu(UUR)} gene: A study in seven families with mitochondrial encephalomyopathy, lactic acidosis and stroke-like episodes (MELAS). *Diabetologia* 37: 818–825
18. American Diabetes Association. (1991) Official guide to diagnosis and classification of diabetes mellitus and other categories of glucose intolerance. *Diabetes Care* 14: 3–4
19. Yamasoba T, Oka Y, Tsukuda K, Nakamura M, Kaga K (1996) Auditory findings in patients with maternal-inherited diabetes mellitus and deafness harboring a point mutation in the mitochondrial transfer RNA^{Leu(UUR)} gene. *Laryngoscope* 106: 49–53
20. Takeda M, Ikenoya S, Yuzuriha T, Katayama K (1982) Studies on reduced and oxidized coenzyme Q in mitochondria, microsome and plasma by high-performance liquid chromatography. *Biochem Biophys Acta* 679: 308–314
21. Silvestre-Aillaud P, Ben-Dahan D, Paquis-Frucking V et al. (1996) Could coenzyme Q₁₀ and L-carnitine be a treatment for diabetes secondary to 3243 mutation of mtDNA? *Diabetologia* 39: 1485–1486
22. Ogasahara S, Engel AG, Frens D et al. (1989) Muscle coenzyme Q deficiency in familial mitochondrial encephalomyopathy. *Proc Natl Acad Sci USA* 86: 2379–2382
23. Frei B, Kim MC, Ames BN (1990) Ubiquinol-10 is an effective lipid-soluble antioxidant at physiological concentrations. *Proc Natl Acad Sci USA* 87: 4879–4883
24. Beyer RE, Segura-Aguilar J, Di Bernardo S et al. (1996) The role of DT-diaphorase in the maintenance of the reduced antioxidant form of coenzyme Q in membrane systems. *Proc Natl Acad Sci USA* 93: 2528–2532
25. Richter C (1995) Oxidative damage to mitochondrial DNA and its relationship to aging. *Int J Biochem Cell Biol* 27: 647–653