

GALOIS POINTS ON VARIETIES

MOSHE JARDEN AND BJORN POONEN

ABSTRACT. A field K is ample if for every geometrically integral K -variety V with a smooth K -point, $V(K)$ is Zariski dense in V . A field K is Galois-potent if every geometrically integral K -variety has a closed point whose residue field is Galois over K . We prove that every ample field is Galois-potent. But we construct also non-ample Galois-potent fields; in fact, every field has a regular extension with these properties.

1. INTRODUCTION

Definition 1.1. Let X be a variety over a field K . By a **Galois point** on X , we mean a closed point whose residue field is Galois over K . We say that K is **Galois-potent** if every geometrically integral K -variety has a Galois point.

Question 1.2. Is every field Galois-potent?

We do not even know if $\mathbb{Q}$ is Galois-potent. On the other hand, we have the following definition of Florian Pop:

Definition 1.3 (cf. [Pop96, p. 2] and [Jar11, Chapter 5]). A field K is called **ample** (or **large** or **anti-Mordellic**) if for every geometrically integral K -variety V with a smooth K -point, $V(K)$ is Zariski dense in V .

Pseudo-algebraically closed (PAC) fields and Henselian fields (e.g., $\mathbb{Q}_p$) are ample; see [Jar11, Chapter 5] for these and many more examples. Our first main theorem is the following:

Theorem 1.4. *Ample fields are Galois-potent.*

Some non-ample fields too are Galois-potent. For example, finite fields are non-ample, but have abelian absolute Galois group and hence trivially are Galois-potent. Less trivially, we can also construct *infinite* non-ample fields that are Galois-potent: in Section 5 we will prove the following.

Theorem 1.5. *Every field admits a regular extension that is Galois-potent but not ample.*

Bary-Soroker and Fehm in [BSF13, Section 2.2] write that “all infinite non-ample fields appearing in the literature are Hilbertian”. The non-ample fields we construct in the proof of Theorem 1.5 turn out to be Hilbertian too (Remark 5.4), and in particular they have non-abelian absolute Galois group.

Date: November 26, 2015.

2010 *Mathematics Subject Classification.* Primary 12E30; Secondary 11G35, 14G99.

The first author was supported by the Minkowski Center for Geometry at Tel Aviv University, established by the Minerva Foundation. The second author was supported in part by National Science Foundation grant DMS-1069236 and a grant from the Simons Foundation (#340694 to Bjorn Poonen). This article has been published in *J. Ramanujan Math. Soc.* **31** (2016), no. 2, 189–194.

Question 1.6. Are there also infinite non-ample fields with abelian absolute Galois group?

If the answer to Question 1.6 were yes, it would immediately yield another example of a non-ample Galois-potent field. But there are several questions and conjectures in the literature that each imply a negative answer to Question 1.6, as we now explain. Let G_K be the absolute Galois group of K .

- Is every infinite non-ample field Hilbertian? (Cf. [BSF13, Section 2.2] again.) A Hilbertian field K cannot have an abelian G_K .
- Is every infinite field K with topologically finitely generated G_K ample? (See the paragraph before Theorem 5.5 of [JK10], or see [BSF13, 4.2, Question III].) By [Koe01], if G_K is abelian, either K is henselian and hence ample, or G_K is procyclic and in particular topologically finitely generated, and hence ample if the question at the start of this bulleted item has a positive answer.
- Is every infinite field with prosolvable G_K ample? (See [BSF13, 4.1, Question II].) If so, then every infinite field with abelian G_K is ample too.

2. NOTATION

Let K be a field. Let $\overline{K}$ be an algebraic closure of K . By a K -variety we mean a separated scheme of finite type over K . Given a K -variety X and a finite extension $L \supset K$, let X_L be the L -variety $X \times_K L$; similarly, if ϕ is a K -morphism, let ϕ_L be its base change. If X is an integral K -variety, let $K(X)$ denote its function field. By a K -curve, we mean a smooth geometrically integral K -variety of dimension 1. By the **absolute genus** g_X of a K -curve X , we mean the genus of the smooth projective model of $X_{\overline{K}}$. Given a curve X and $n \geq 1$, let $X^{(n)}$ denote the n^{th} **symmetric power**, defined as the (variety) quotient of X^n by the symmetric group S_n .

3. RESTRICTION OF SCALARS

Let $K \subseteq L$ be a finite extension of fields. For any quasi-projective L -variety X , the functor sending each K -scheme S to the set $X(S_L)$ is representable by a K -variety $\text{Res}_{L/K} X$ called the **restriction of scalars** (cf. [BLR90, §7.6, Theorem 4]). If, in addition, L is separable over K , then $\text{Res}_{L/K} X$ after base field extension becomes isomorphic to a product of $[L : K]$ conjugates of X ; in particular, if X is geometrically integral and smooth of dimension d over L , then $\text{Res}_{L/K} X$ is geometrically integral and smooth of dimension $[L : K]d$ over K .

4. AMPLE FIELDS ARE GALOIS-POTENT

In this section we prove Theorem 1.4.

Proposition 4.1. *Let K be an ample field. Let X be a geometrically integral K -variety. Let N be a finite separable extension of K . If X has a smooth N -point, then X has a point with residue field N .*

Proof. Replacing X by a Zariski-open subvariety, we may assume that X is smooth and affine. For each finite separable extension L of K , define $X^L := \text{Res}_{L/K}(X_L)$. Thus $X^L(S) = X_L(S_L) = X(S_L)$ for any K -scheme S , and in particular, $X^L(K) = X(L)$. If $L \subseteq L'$ are two such extensions, the natural map $X(S_L) \rightarrow X(S_{L'})$ can be rewritten as $X^L(S) \rightarrow X^{L'}(S)$, and it is functorial in S , so Yoneda's lemma yields a K -morphism $X^L \rightarrow X^{L'}$.

We have $X^N(K) = X(N)$, which is nonempty by assumption. Also, K is ample, and X^N is smooth and geometrically integral, so $X^N(K)$ is Zariski-dense in X^N . There are only finitely many fields L with $K \subseteq L \subsetneq N$, and for each such L , we have $\dim X^L < \dim X^N$. Thus X^N has a K -point outside the image of every morphism $X^L \rightarrow X^N$. In other words, $X(N)$ has a element outside $X(L)$ for every L . For this element, the image of $\text{Spec } N \rightarrow X$ is a closed point with residue field N . $\square$

Proof of Theorem 1.4. Let K be an ample field, and let X be any geometrically integral K -variety. Choose a finite separable extension N of K such that $X(N) \neq \emptyset$. Enlarge N to assume that N is Galois over K . By Proposition 4.1, X has a closed point with residue field N . $\square$

5. INFINITE GALOIS-POTENT FIELDS THAT ARE NOT AMPLE

In this section we prove Theorem 1.5.

Lemma 5.1. *Let K be an algebraically closed field. Let X, Y, C be K -curves with $g_C > 1$. Any rational map $X \times Y \dashrightarrow C$ factors through one of the projections to X or Y .*

Proof. A rational map $\phi: X \times Y \dashrightarrow C$ may be viewed as an algebraic family of rational maps $X \dashrightarrow C$ parametrized by (an open subvariety of) Y . But the de Franchis–Severi theorem [Sam66, Théorème 1] implies that there are no nonconstant algebraic families of nonconstant rational maps $X \dashrightarrow C$. Thus either the rational maps in the family are all the same, in which case ϕ factors through the first projection, or each rational map in the family is constant, in which case ϕ factors through the second projection. $\square$

Lemma 5.2. *Let X be a curve over a field K . Let $F = K(X^{(2)})$. Then X has a point over a quadratic extension of F , and $C(F) = C(K)$ for every K -curve C of absolute genus > 1 .*

Proof. Either projection $X \times X \rightarrow X$ gives a point of X over the quadratic extension $K(X \times X)$ of $K(X^{(2)}) = F$.

Let $c \in C(F)$. Then c corresponds to a rational map $X^{(2)} \dashrightarrow C$. Composing $X \times X \rightarrow X^{(2)}$ with such a rational map yields a rational map $\phi: X \times X \dashrightarrow C$. By Lemma 5.1, $\phi_{\bar{K}}$ is constant on all vertical copies of $X_{\bar{K}}$ or constant on all horizontal copies of $X_{\bar{K}}$. Since $\phi_{\bar{K}}$ is S_2 -invariant, it is constant on all vertical and horizontal copies of $X_{\bar{K}}$. Thus $\phi_{\bar{K}}$ is constant. Hence ϕ is constant. Equivalently, $c \in C(K)$. $\square$

Lemma 5.3. *Every field K admits a regular extension K' such that*

- (i) *Every K -curve X has a point over an at most quadratic extension of K' (possibly depending on X).*
- (ii) *For every K -curve C of absolute genus > 1 , we have $C(K') = C(K)$.*

Proof. Let $(X_\alpha)_{\alpha < \tau}$ be a well-ordering of the set of K -curves up to isomorphism, indexed by an ordinal τ . For $\alpha \leq \tau$, define K_α by transfinite induction as follows:

- Let $K_0 := K$.
- For each $\alpha < \tau$, define $K_{\alpha+1} := K_\alpha(X_\alpha^{(2)})$;
- If α is a limit ordinal, define $K_\alpha := \varinjlim_{\beta < \alpha} K_\beta$.

Let $K' := K_\tau$. Then K' is regular over K by [FJ08, Corollary 2.6.5(d)], whose proof remains valid when the cardinal m is replaced by an ordinal τ .

- (i) Any K -curve X is X_α for some $\alpha < \tau$. Lemma 5.2 shows that X has a point over a quadratic extension of $K_{\alpha+1}$, and hence also a point over an at most quadratic extension of K' .
- (ii) Let C be a K -curve of absolute genus > 1 . By Lemma 5.2 and induction, $C(K_\alpha) = C(K)$ for all $\alpha \leq \tau$. In particular, $C(K') = C(K)$. $\square$

Proof of Theorem 1.5. Let K be the given field. Construct a sequence of fields $L_0, L_1, \dots$ inductively as follows: let $L_0 := K(t)$ with t transcendental over K , and let L_{i+1} be the regular extension of L_i given by Lemma 5.3. Let $L_\infty := \varinjlim L_i$.

Let X be an L_∞ -curve. Then X is definable over some $\tilde{L}_i$. The conditions in Lemma 5.3 imply that X has a point over an at most quadratic extension of L_{i+1} , and hence a point over an at most quadratic extension of L_∞ . Every geometrically integral L_∞ -variety other than a point contains an L_∞ -curve, so L_∞ is Galois-potent.

Choose a non-isotrivial curve C of absolute genus > 1 over $L_0 = K(t)$ such that C has a smooth L_0 -point; then $C(L_0)$ is finite [Sam66, Théorème 4]. The conditions in Lemma 5.3 imply that $C(L_0) = C(L_1) = \dots$, so $C(L_\infty) = C(L_0)$, which is finite. Thus L_∞ is not ample. $\square$

Remark 5.4. The field L_∞ constructed in the proof of Theorem 1.5 is Hilbertian, because of the following two facts:

- (a) Any finitely generated transcendental extension of a field is Hilbertian [FJ08, Theorem 13.4.2].
- (b) If $(K_\alpha)_{\alpha \leq \tau}$ is an ascending tower of fields indexed by an ordinal τ , and K_α is Hilbertian for each $\alpha < \tau$, and $K_{\alpha+1}$ is a regular extension of K_α for each $\alpha < \tau$, and $K_\alpha = \varinjlim_{\beta < \alpha} K_\beta$ for each limit ordinal $\alpha \leq \tau$, then K_τ is Hilbertian.

(Fact (b) appears as [FJ08, Chapter 12, Exercise 2], but the hypothesis $K_\alpha = \varinjlim_{\beta < \alpha} K_\beta$ is missing there.)

Proof of (b). Given irreducible polynomials $f_i(\vec{t}, x)$ over K_τ for $i = 1, \dots, r$, each separable in x , we must find infinitely many $\vec{a}$ over K_τ such that the specializations $f_i(\vec{a}, x)$ for $i = 1, \dots, r$ are irreducible over K_τ . There exists $\alpha < \tau$ such that all the f_i are defined over K_α . Since K_α is Hilbertian, there exist infinitely many $\vec{a}$ over K_α such that $f_i(\vec{a}, x)$ for each i is irreducible over K_α . These specializations are irreducible over K_τ as well, since K_τ is a regular extension of K_α by [FJ08, Corollary 2.6.5(d)]. $\square$

Remark 5.5. Padmavathi Srinivasan has constructed a field K with stronger properties than the L_∞ in the proof of Theorem 1.5, namely that $C(K)$ is finite for every K -curve C of absolute genus > 1 , while there is a degree 2 field extension $L \supset K$ that is ample [Sri15].

ACKNOWLEDGEMENTS

We thank the referee for many insightful comments and corrections.

REFERENCES

- [BSF13] Lior Bary-Soroker and Arno Fehm, *Open problems in the theory of ample fields*, Geometric and differential Galois theories, Sémin. Congr., vol. 27, Soc. Math. France, Paris, 2013, pp. 1–11 (English, with English and French summaries). MR3203546

- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 21, Springer-Verlag, Berlin, 1990. MR1045822 (91i:14034)
- [FJ08] Michael D. Fried and Moshe Jarden, *Field arithmetic*, 3rd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 11, Springer-Verlag, Berlin, 2008. Revised by Jarden. MR2445111 (2009j:12007)
- [Jar11] Moshe Jarden, *Algebraic patching*, Springer Monographs in Mathematics, Springer, Heidelberg, 2011. MR2768285 (2012e:12007)
- [JK10] Markus Junker and Jochen Koenigsmann, *Schlanke Körper (slim fields)*, J. Symbolic Logic **75** (2010), no. 2, 481–500, DOI 10.2178/jsl/1268917491. MR2648152 (2011f:03047)
- [Koe01] Jochen Koenigsmann, *Solvable absolute Galois groups are metabelian*, Invent. Math. **144** (2001), no. 1, 1–22, DOI 10.1007/s002220000117. MR1821143 (2002a:12006)
- [Pop96] Florian Pop, *Embedding problems over large fields*, Ann. of Math. (2) **144** (1996), no. 1, 1–34. MR1405941 (97h:12013)
- [Sam66] Pierre Samuel, *Compléments à un article de Hans Grauert sur la conjecture de Mordell*, Inst. Hautes Études Sci. Publ. Math. **29** (1966), 55–62 (French). MR0204430 (34 #4272)
- [Sri15] Padmavathi Srinivasan, *A virtually ample field that is not ample*, December 31, 2015. Preprint, available at http://math.mit.edu/~padma_sk/virtually_ample.pdf.

SCHOOL OF MATHEMATICS, TEL AVIV UNIVERSITY, RAMAT AVIV, TEL AVIV 6139001, ISRAEL
E-mail address: jarden@post.tau.ac.il

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139-4307, USA
E-mail address: poonen@math.mit.edu
URL: <http://math.mit.edu/~poonen/>