

A COMPUTABLE FUNCTOR FROM GRAPHS TO FIELDS

RUSSELL MILLER, BJORN POONEN, HANS SCHOUTENS, AND ALEXANDRA SHLAPENTOKH

Abstract. Fried and Kollár constructed a fully faithful functor from the category of graphs to the category of fields. We give a new construction of such a functor and use it to resolve a longstanding open problem in computable model theory, by showing that for every nontrivial countable structure $\mathcal{S}$, there exists a countable field $\mathcal{F}$ of arbitrary characteristic with the same essential computable-model-theoretic properties as $\mathcal{S}$. Along the way, we develop a new “computable category theory”, and prove that our functor and its partially-defined inverse (restricted to the categories of countable graphs and countable fields) are computable functors.

§1. Introduction.

1.A. A functor from graphs to fields. Let **Graphs** be the category of symmetric irreflexive graphs in which morphisms are isomorphisms onto induced subgraphs (see Section 2.A). Let **Fields** be the category of fields, with field homomorphisms as the morphisms.

Fried and Kollár [FK82, Theorem 2.1]¹ proved the following:

THEOREM 1.1. *There exists a fully faithful functor $\mathcal{F}: \mathbf{Graphs} \rightarrow \mathbf{Fields}$.*

The definitions of “full” and “faithful” are reviewed in Section 2.B. Using Theorem 1.1, Fried and Kollár showed that every group arises as the automorphism group of a field [FK82, Theorem 1.1].

In the present paper, we

- give a new construction of such a functor, using the arithmetic of curves instead of towers of fields obtained by iteratively adjoining indeterminates and adjoining p th roots as in [FK82];
- construct computable versions of the functor and its partially-defined inverse, computable in the sense of a “computable category theory” that we create in Section 3; and
- derive consequences in computable model theory, and in particular show that for every nontrivial countable structure $\mathcal{S}$, there exists a countable field $\mathcal{F}$ with the same essential computable-model-theoretic properties as $\mathcal{S}$. This answers a question left open in [HKSS02].

1.B. Computable functors. For applications to computable model theory, we are interested in graphs and fields whose underlying set is $\omega := \{0, 1, 2, \dots\}$. These form full subcategories **Graphs** $_{\omega}$ and **Fields** $_{\omega}$. The functor $\mathcal{F}$ of Theorem 1.1 will be constructed so that it restricts to a functor $\mathcal{F}_{\omega}: \mathbf{Graphs}_{\omega} \rightarrow \mathbf{Fields}_{\omega}$. Let **E** $_{\omega}$ denote the essential image of $\mathcal{F}_{\omega}$ (see Section 2.B for definitions). Then we may view $\mathcal{F}_{\omega}$ as a functor from **Graphs** $_{\omega}$ to **E** $_{\omega}$. We will also define a functor $\mathcal{G}_{\omega}: \mathbf{E}_{\omega} \rightarrow \mathbf{Graphs}_{\omega}$, and we would like to say that $\mathcal{F}_{\omega}$ and $\mathcal{G}_{\omega}$ are computable and are inverse to each other in some computable way.

To guide us to the correct formulation of such statements, we create a new “type-2 computable category theory”; see Section 3. The adjective “type-2”, borrowed from computable analysis (see Remark 3.2), indicates

Received by the editors June 11, 2017.

1991 *Mathematics Subject Classification.* MSC 03C57 (primary), 03D45, 12L12, 18A15, 08A35 (secondary).

Key words and phrases. completeness, computability, computable model theory, fields, functor, graphs.

This article is to appear in *J. Symbolic Logic*. This project was initiated at a workshop held at the American Institute of Mathematics. The first author was partially supported by National Science Foundation grants DMS-1001306 and DMS-1362206, and by several PSC-CUNY Research Awards.

The second author was partially supported by National Science Foundation grants DMS-1069236 and DMS-1601946 and grants from the Simons Foundation (340694 and 402472 to Bjorn Poonen).

The fourth author was partially supported by National Science Foundation grant DMS-1161456.

¹The category of graphs in [FK82] was slightly different: morphisms were required to be injective on vertices and to map edges to edges, but were not required to map non-edges to non-edges. This is not a significant difference, however.

that we work with noncomputable objects; indeed, $\mathbf{Graphs}_\omega$ and $\mathbf{Fields}_\omega$ each contain uncountably many noncomputable objects. The effectiveness in our definitions really arises in the concept of a *computable functor*, a functor in which the processes of transforming objects to objects and morphisms to morphisms are given by Turing functionals: roughly speaking, each output should be computable given an oracle for the input (see Definition 3.1). Our computable category theory includes also a notion of *computable isomorphism of functors* (see Definition 3.3). Using this lexicon, we can now state our main result on the computability of the functors:

THEOREM 1.2.

- (a) The functors $\mathcal{F}_\omega$ and $\mathcal{G}_\omega$ are computable in the sense of Definition 3.1.
- (b) The composition $\mathcal{G}_\omega \mathcal{F}_\omega$ is computably isomorphic to $1_{\mathbf{Graphs}_\omega}$, and $\mathcal{F}_\omega \mathcal{G}_\omega$ is computably isomorphic to $1_{\mathbf{E}_\omega}$.

We summarize Theorem 1.2 by saying that $\mathcal{F}_\omega$ and $\mathcal{G}_\omega$ give a computable equivalence of categories between $\mathbf{Graphs}_\omega$ and $\mathbf{E}_\omega$ (see Definition 3.4). In fact, we will define $\mathcal{G}_\omega$ so that $\mathcal{G}_\omega \mathcal{F}_\omega$ equals $1_{\mathbf{Graphs}_\omega}$, but computable isomorphism in place of equality here would suffice for the applications in Section 9.

Here is one concrete consequence of Theorem 1.2:

COROLLARY 1.3. *If a field $F \in \mathbf{Fields}_\omega$ is isomorphic to a field in the image of $\mathcal{F}_\omega$, then one can compute from F a graph $G \in \mathbf{Graphs}_\omega$ and an isomorphism $F \rightarrow \mathcal{F}_\omega(G)$.*

PROOF. Apply the isomorphism $\mathcal{G}_\omega \mathcal{F}_\omega \simeq 1_{\mathbf{Graphs}_\omega}$ of Theorem 1.2(b) to $G := \mathcal{G}_\omega(F)$. $\dashv$

Specializing Corollary 1.3 to the case in which F is computable yields the following:

COROLLARY 1.4. *Every computable field isomorphic to a field in the image of $\mathcal{F}_\omega$ is computably isomorphic to $\mathcal{F}_\omega(G)$ for some computable $G \in \mathbf{Graphs}_\omega$.*

Following [Kni86, §4], we call a structure **automorphically trivial** if there exists a finite subset S_0 of its domain S such that every permutation of S fixing S_0 pointwise is an automorphism.

EXAMPLE 1.5. The automorphically trivial graphs $G = (S, E)$ are exactly those obtained as follows: choose a partition $S = S_0 \amalg S_1$ with S_0 finite, choose a subset $S'_0 \subseteq S_0$, and let E be the union of the edge sets of a graph on S_0 , the complete bipartite graph on S'_0 and S_1 , and optionally the complete graph on S_1 .

PROPOSITION 1.6. *Let $F \in \mathbf{E}_\omega$; that is, F is isomorphic to a field in the image of $\mathcal{F}_\omega$. Let $\mathcal{I} := \{G \in \mathbf{Graphs}_\omega : \mathcal{F}_\omega(G) \simeq F\}$, so $\mathcal{I} \neq \emptyset$.*

- (a) The set $\mathcal{I}$ is an isomorphism class in $\mathbf{Graphs}_\omega$.
- (b) Either every graph in $\mathcal{I}$ is automorphically trivial and computable, or there exists a graph in $\mathcal{I}$ of the same Turing degree as F .

Proofs of Theorem 1.2 and Proposition 1.6 appear in Section 7.

Remark 1.7. As will be explained in Section 3.C, there have been other attempts to give effective versions of category theory (and, more successfully, to give a categorical underpinning to computability theory). To our knowledge, however, ours is the first attempt to define effectiveness for functors using type-2 Turing computation.

1.C. Computable model theory. One of the goals of computable model theory is to help distinguish various classes of countable structures according to the algorithmic complexity of those structures. The class of algebraically closed fields of characteristic 0, for example, is viewed throughout model theory as a particularly simple class, and its computability-theoretic properties confirm this view: every countable algebraically closed field can be computably presented, all of them are relatively Δ_2^0 -categorical, and the only one that is not relatively computably categorical has infinite computable dimension. (All these terms are defined in Section 9.) In contrast, the theory of linear orders is a good deal more complex: there do exist countable linear orders with no computable presentation, and linear orders with much higher degrees of categoricity than $\mathbf{0}'$. In this view, the theory of graphs is even more complicated: for example, every computable linear order has computable dimension either 1 or ω , whereas for computable (symmetric irreflexive) graphs all computable dimensions $\leq \omega$ are known to occur.

We will discuss specific properties such as computable presentability and computable dimension when we come to prove results about them, in Section 9. For now, we simply note that a substantial body of results has been established on the possibility of transferring these properties from one class of countable structures to another. After much piecemeal work by assorted authors, most of these results were gathered together and brought to completion in the work [HKSS02], by Hirschfeldt, Khoussainov, Shore, and Slinko. There it was proven that the class of symmetric irreflexive graphs is *complete*, in the very strong sense of their Definition 1.21. The authors gave a coding procedure that, given any countable structure S with domain ω (in an arbitrary computable language, with one quite trivial restriction) as its input, produced a countable graph G on the same domain with the same computable-model-theoretic properties as S . Several other natural properties have been introduced since then (the automorphism spectrum, in [HMM10, Definition 1.1], and the categoricity spectrum, in [FKM10, Definition 1.2], for example), and each of these has also turned out to be preserved under the construction from [HKSS02]. The method they gave was quite robust, in this sense, and one may expect that it will also be found to preserve other properties that are yet to be defined.

Having established the completeness of the class of countable graphs in this sense, the authors went on to consider many other everyday classes of countable first-order structures. By doing a similar coding from graphs into other classes, they succeeded in proving the completeness (in this same sense) of the following classes:

- countable directed graphs;
- countable partial orderings;
- countable lattices;
- countable rings (with zero-divisors);
- countable integral domains of arbitrary characteristic;
- countable commutative semigroups;
- countable 2-step nilpotent groups.

(We note again that in some cases, these results had been established in earlier work by other authors. In certain of these cases, the language must be augmented by a finite number of constant symbols.) On the other hand, various existing and subsequent results demonstrated that none of the following classes is complete in this sense:

- countable linear orders (e.g., by results in [GD80, Theorem 2], [Rem81, Corollary 1], and [Ric81, Theorem 3.3]);
- countable Boolean algebras (see [DJ94, Theorem 1], or use [Ric81, Theorem 3.1]);
- countable trees, either as partial orders or under the meet relation $\wedge$ (by [Ric81, Theorem 3.4] or [LMMS05, Theorem 1.8]);
- algebraic field extensions of $\mathbb{Q}$ or of $\mathbb{F}_p$ (see [Mil09, Corollary 5.5], for example);
- field extensions of $\mathbb{Q}$ of finite transcendence degree (see [Mil09, §6]);
- countable archimedean ordered fields (by Theorem 3.3.1 in Levin’s thesis [Lev09]).

In addition, Goncharov announces in [Gon81] that every computable abelian group has computable dimension 1 or ω , though it seems that a detailed proof does not exist in print. This result would add the class of countable abelian groups to the list of non-complete classes.

Conspicuously absent from both of the lists above is the class of countable fields. The question whether this class possesses the completeness property has remained open, despite substantial interest since the publication of [HKSS02]: see the introduction to [LMMS05]. In this paper we resolve the question, using the computable functors of Section 1.B:

THEOREM 1.8. *The class of countable fields has the completeness property of [HKSS02, Definition 1.21].*

Proving the many specific aspects of this theorem will take most of Section 9. Our proof of Theorem 1.8 shows more specifically that the class of countable fields of characteristic 0 has the completeness property. In Section 10, we will prove the same for characteristic p for each prime p .

Remark 1.9. The constructions in [HKSS02] too can be expressed in terms of our computable category theory.

1.D. Structure of the paper. Section 2 introduces notation and definitions. Section 3 introduces the key definitions of our computable category theory, and discusses its relation to other work in the literature. Section 4 defines some algebraic curves used in Section 5 to construct $\mathcal{F}$, and proves arithmetic properties of these curves that are used in Section 6 to construct $\mathcal{G}$ and prove enough properties of $\mathcal{F}$ and $\mathcal{G}$ to prove Theorem 1.1. Section 7 constructs the computable analogues $\mathcal{F}_\omega$ and $\mathcal{G}_\omega$, and proves Theorem 1.2. Section 8 proves that for every $G \in \mathbf{Graphs}_\omega$, the field $\mathcal{F}_\omega(G)$ is isomorphic to a subfield of $\mathbb{R}$; but $\mathcal{F}_\omega$ cannot be viewed as a functor to the category of ordered fields if morphisms in the latter are required to respect the orderings. Section 9 explains the implications of our functors for computable model theory. Many of the results there follow formally from [HKSS02, Theorem 3.1] and Theorem 1.2, so they could be stated more generally in terms of any computable equivalence of categories of structures, but for concreteness, we state them specifically for the categories of graphs and fields. Finally, Section 10 explains how the Fried–Kollár approach can be combined with Markerization to yield analogous computable-model-theoretic results for the category of fields of prescribed characteristic.

§2. Notation and definitions.

2.A. Graphs. Given a set I , let $\binom{I}{2}$ be the set of 2-element subsets of I . A (symmetric, irreflexive) graph is a set V equipped with a subset $E \subseteq \binom{V}{2}$; then $\{(i, j) \in V \times V : \{i, j\} \in E\}$ is a symmetric irreflexive relation on V . If G is a graph (V, E) , then $\#G := \#V$; call G finite if V is finite. Define a **morphism of graphs** $(V, E) \rightarrow (V', E')$ to be an injection $f: V \hookrightarrow V'$ such that for each $\{i, j\} \in \binom{V}{2}$, we have $\{i, j\} \in E$ if and only if $\{f(i), f(j)\} \in E'$. In other words, a morphism of graphs $(V, E) \rightarrow (V', E')$ is an isomorphism from (V, E) onto an induced subgraph of (V', E') . These notions define a category **Graphs**.

2.B. Category theory. A full subcategory of a category $\mathbf{C}$ is a category consisting of some the objects of $\mathbf{C}$ but all of the morphisms between pairs of these chosen objects. Let $\mathcal{F}: \mathbf{C} \rightarrow \mathbf{D}$ be a functor. Then $\mathcal{F}$ is full (respectively, faithful) if for any two objects $C_1, C_2 \in \mathbf{C}$, the map $\text{Hom}(C_1, C_2) \rightarrow \text{Hom}(\mathcal{F}(C_1), \mathcal{F}(C_2))$ is surjective (respectively, injective). The **essential image** of $\mathcal{F}$ is the full subcategory of $\mathbf{D}$ consisting of objects $D \in \mathbf{D}$ that are isomorphic to $\mathcal{F}(C)$ for some $C \in \mathbf{C}$.

2.C. Fields and arithmetic geometry. Let **Fields** be the category of fields, with field homomorphisms as the morphisms. If X is a k -variety, and $L \supseteq k$ is a field extension, let X_L denote the base extension; concretely, X_L is the L -variety defined by the same equations as X , but with coefficients considered to be in L . A k -variety is **integral** if it is irreducible and the ring of functions on any Zariski open subset has no nilpotent elements. A k -variety is **geometrically integral** if it remains integral after any extension of the ground field; for example, the affine curve $x^2 - 2y^2 = 0$ over $\mathbb{Q}$ is integral but not geometrically integral, because the polynomial $x^2 - 2y^2$ is irreducible over $\mathbb{Q}$ but not irreducible over $\mathbb{Q}(\sqrt{2})$. If X is an integral k -variety, let $k(X)$ denote its function field, defined as the fraction field of the affine coordinate ring of any nonempty affine open subvariety of X . If $(X_i)_{i \in I}$ is a collection of geometrically integral k -varieties indexed by a set I , let $k(\prod X_i)$ denote the direct limit of the function fields $k(\prod_{i \in J} X_i)$ over all finite subsets $J \subseteq I$; this is a compositum of the function fields of the X_i . A **birational map** between integral varieties $X \dashrightarrow Y$ is a rational map $f: X \dashrightarrow Y$ for which there exists a rational map $g: Y \dashrightarrow X$ such that gf and fg are defined and equal to the identity on some Zariski dense open subvarieties of X and Y , respectively; then say that X and Y are birational. A **birational automorphism** of X is a birational map $X \dashrightarrow X$; let $\text{Bir } X$ be the group of birational automorphisms of X . Every geometrically integral curve X over a perfect field k is birational to a unique smooth projective geometrically integral curve $\tilde{X}$; let $g_X \in \mathbb{Z}_{\geq 0}$ be the **geometric genus** of X , defined as the dimension of the space of global 1-forms on $\tilde{X}$.

2.D. Computable model theory. Given a structure M , let $\text{dom}(M)$ be its domain (underlying set), and let $\Delta(M)$ be its **atomic diagram** (the set of atomic sentences true in M). If M is a countable structure with $\text{dom}(M) = \omega$, let $\deg M$ be the Turing degree of $\Delta(M)$, and define the **spectrum** of M as $\text{Spec } M := \{\deg N : N \simeq M \text{ and } \text{dom}(N) = \omega\}$. Given two such structures M and N , we say that M is **computable** from N , or write $M \leq_T N$, if $\Delta(M)$ is computable under a $\Delta(N)$ -oracle, or equivalently $\deg M \leq_T \deg N$.

§3. Computable category theory.

3.A. Computable functors. By a category of structures on ω , we mean a subcategory $\mathbf{C}$ of the category of all first-order $\mathcal{L}$ -structures with domain $\omega := \{0, 1, 2, \dots\}$, for some computable language $\mathcal{L}$. (In some cases, one might allow also finite subsets of ω as domains.) Here are two examples:

- Let $\mathbf{Graphs}_\omega$ be the category whose objects are the (symmetric irreflexive) graphs having underlying set ω , and whose morphisms are isomorphisms onto induced subgraphs.
- Let $\mathbf{Fields}_\omega$ be the category whose objects are the fields having underlying set ω , and whose morphisms are field homomorphisms.

There is nothing particularly effective about these categories. The requirement that the domain equal ω gives us the opportunity to consider computability questions about the structures in a category, but a graph G on ω with a noncomputable edge relation would be an object of $\mathbf{Graphs}_\omega$ in perfectly good standing. Also, even for computable sets $E \subseteq \omega^2$, there is no general way to determine whether E actually is the edge relation of a symmetric irreflexive graph.

Effectiveness considerations arise when we consider functors between these categories. A functor maps objects to objects and morphisms to morphisms, and we would like this process to be effective.

DEFINITION 3.1. Let $\mathbf{C}$ and $\mathbf{C}'$ be categories of structures on ω (with respect to possibly different languages $\mathcal{L}$ and $\mathcal{L}'$). A **computable functor** is a functor $\mathcal{F}: \mathbf{C} \rightarrow \mathbf{C}'$ for which there exist Turing functionals Φ and Ψ such that

- (i) for every $S \in \mathbf{C}$, the function Φ^S computes (the atomic diagram of) the structure $\mathcal{F}(S)$; and
- (ii) for every morphism $g: S \rightarrow T$ in $\mathbf{C}$, we have $\Psi^{S \oplus g \oplus T} = \mathcal{F}(g)$ in $\mathbf{C}'$.

Computing the atomic diagram of $\mathcal{F}(S)$ is equivalent to computing every function and relation from $\mathcal{L}'$ on $\mathcal{F}(S)$ (and, in case $\mathcal{L}'$ is infinite, doing so uniformly and also computing the value of each constant symbol). One could state (i) by saying that, for every $n \in \omega$, the function $\Phi^S(\langle n, \vec{x} \rangle)$ is the value on input $\vec{x}$ of the n th symbol of the language $\mathcal{L}'$, as interpreted in $\mathcal{F}(S)$. For relation symbols, this value is Boolean, while for constants and function symbols, it is an element of $\mathcal{F}(S)$, i.e., a natural number. Condition (ii) states that $\Psi^{S \oplus g \oplus T}$ computes the morphism $\mathcal{F}(g)$ in $\mathbf{C}'$, viewed as a map from $\text{dom}(S)$ into $\text{dom}(T)$. Note that one is allowed to use the objects S and T , in addition to the morphism g between them, to compute $\mathcal{F}(g)$; this is natural if one thinks of the source and target as being part of the data describing a function. Our proofs in Section 7 that $\mathcal{F}_\omega$ and $\mathcal{G}_\omega$ are computable also illustrate why including S and T is appropriate.

Remark 3.2. Our computable functors may also be called **type-2 computable functors**. The terminology “type-2” has been used by researchers as far back as Kleene, for functions F whose domain and range are subsets of Cantor space 2^ω or Baire space ω^ω , and which are given by a Turing functional Φ such that Φ^x is equal to $F(x)$. Today it is primarily used in computable analysis, where this concept of computability is common; the book [PER89] is a standard reference for this topic. Likewise, here in category theory, we are computing functions not between natural numbers, but between sets of natural numbers. This is clear in the case of the edge relation on a graph, but in fact all countable structures (in countable first-order languages) turn out to be represented by subsets of ω , just as all real numbers are.

3.B. Computable morphisms of functors.

DEFINITION 3.3. Let $\mathcal{F}_1, \mathcal{F}_2: \mathbf{C} \rightarrow \mathbf{C}'$ be computable functors. A **computable morphism of functors** (or **computable natural transformation**) from $\mathcal{F}_1$ to $\mathcal{F}_2$ is a morphism of functors $\tau: \mathcal{F}_1 \rightarrow \mathcal{F}_2$ such that there exists a Turing functional that on input $S \in \mathbf{C}$ computes the morphism $\tau(S): \mathcal{F}_1(S) \rightarrow \mathcal{F}_2(S)$.

This notion leads in the usual way to the notion of **computable isomorphism of functors**: this is a computable morphism of functors having a two-sided inverse that is again a computable morphism of functors. Then, adapting the definition of equivalence of categories leads to the following.

DEFINITION 3.4. Let $\mathbf{C}$ and $\mathbf{C}'$ be categories of structures on ω . A **computable equivalence of categories** from $\mathbf{C}$ to $\mathbf{C}'$ is a pair of computable functors $\mathcal{F}: \mathbf{C} \rightarrow \mathbf{C}'$ and $\mathcal{G}: \mathbf{C}' \rightarrow \mathbf{C}$ together with computable isomorphisms $\mathcal{G}\mathcal{F} \rightarrow 1_{\mathbf{C}}$ and $\mathcal{F}\mathcal{G} \rightarrow 1_{\mathbf{C}'}$.

In Definition 3.4, we may refer to $\mathcal{F}$ alone as a computable equivalence of categories if the other functors and isomorphisms exist.

3.C. Related work. To a certain extent, the concept of computable functor has appeared before in computable model theory. Turing-computable reducibility (as defined, e.g., in [KMVB07, §1], by Knight, S. Miller, and Vanden Boom) can be viewed as a version of it. In that definition, one class $\mathfrak{C}$ of computable structures is Turing-computably reducible (or TC-reducible) to another such class $\mathfrak{D}$ if there exists a Turing functional Φ that accepts as input the atomic diagram $\Delta(S)$ of a structure $S \in \mathfrak{C}$ and outputs the atomic diagram $\Phi^{\Delta(S)} = \Delta(T)$ of a structure $T \in \mathfrak{D}$. Writing $\mathcal{F}(S)$ for T , the further requirement is that $S \cong S'$ if and only if $\mathcal{F}(S) \cong \mathcal{F}(S')$. So this definition essentially includes the first half of Definition 3.1 above, although stated only for computable structures, not for arbitrary structures with domain ω . The second half is related to the preservation of the isomorphism relation, but here Definition 3.1 is far stronger, requiring the actual computation of an isomorphism $\mathcal{F}(g)$ from $\mathcal{F}(S)$ onto $\mathcal{F}(S')$, given an isomorphism g from S onto S' . It would be reasonable to investigate Definition 3.1 more fully, especially in light of the work in [KMVB07].

In parallel with our introduction of computable category theory, Montalbán examined a notion that he calls **effective interpretation**. This is detailed in [Mon14, §5], and is very much in the vein of the traditional model-theoretic notion of interpretation, with effectiveness conditions added. The functor we build in Section 5 allows for an effective interpretation of each graph G in the field $\mathcal{F}(G)$, since the domain of each graph is definable uniformly by an existential formula in the corresponding field and both the edge relation of the graph and its complement are likewise uniformly existentially definable on that domain within that field. In the other direction, $\mathcal{F}$ has a computable left inverse functor, and it turns out that $\mathcal{F}(G)$ has an effective interpretation in G . In fact, in the wake of the present article, Harrison-Trainor, Melnikov, Miller, and Montalbán proved in [HTMMM17, Theorems 5 and 12] that for categories of structures $\mathbf{C}$ and $\mathbf{C}'$ in which all morphisms are isomorphisms, the existence of a computable functor $\mathbf{C} \rightarrow \mathbf{C}'$ is equivalent to the uniform effective interpretability of all elements of $\mathbf{C}'$ in $\mathbf{C}$; applying this to $\mathcal{F}$ and $\mathcal{G}$ (restricted to the subcategories of $\mathbf{Graphs}_\omega$ and $\mathbf{E}_\omega$ containing all objects but only the isomorphisms) explains the effective interpretations of the previous two sentences.

The results in [HKSS02] are proven largely by the construction of computable functors, although not described in that way. However, one could also ask the same questions about categories known not to be complete. For example, there is a natural construction of a Boolean algebra $\mathcal{F}(L)$ from a linear order L , simply by taking the interval algebra of L , where the morphisms in each category are simply homomorphisms of the structures. On its face, this functor appears to be neither full nor faithful, based on known results, and it does not have a precise computable inverse functor on its image, although it may come close to doing so. It cannot have all of these properties, because there does exist a linear order whose spectrum is not realized by any Boolean algebra, as shown by Jockusch and Soare in [JS91, Theorem 1]. (Here we use a generalization of the result in this article, namely, that a computable equivalence of categories onto a strictly full subcategory allows one to transfer spectra from objects of the first category to objects of the second. This generalization appears to have a straightforward direct proof, and in any case it follows from effective bi-interpretability, hence from [HTMMM17, Theorem 12], using [Mon14, Lemma 5.3].) We suspect that similar results distinguishing the properties of various everyday classes of countable structures may yield further insights into effectiveness, fullness, faithfulness, and other properties of functors among these classes, especially the incomplete ones listed in Section 1.C.

§4. Curves. Define polynomials

$$p(u, v) := u^4 + 16uv^3 + 10v^4 + 16v - 4 \in \mathbb{Q}[u, v], \text{ and} \\ q(T, x, y) := x^4 + y^4 + 1 + T(x^4 + xy^3 + y + 1) \in \mathbb{Q}[T, x, y].$$

Let X be the affine plane curve $p(u, v) = 0$ over $\mathbb{Q}$. For any field F and $t \in F$, let Y_t be the affine plane curve $q(t, x, y) = 0$ over F . If we take $F = \mathbb{Q}(T)$ (the field of rational functions in one indeterminate) and $t = T$, then we obtain a curve Y_T over $\mathbb{Q}(T)$; let $Y = Y_T$. More generally, if $F \supseteq \mathbb{Q}$ and t is transcendental over $\mathbb{Q}$, then Y_t is the base change of Y by the field homomorphism $\mathbb{Q}(T) \rightarrow F$ sending T to t , so Y_t inherits many properties from Y .

The properties we need of these curves to construct the functor $\mathcal{F}$ are contained in parts (1)–(7) of the following lemma. Later, to prove that for every $G \in \mathbf{Graphs}_\omega$ the field $\mathcal{F}(G)$ is isomorphic to a subfield of $\mathbb{R}$, we will also need (8).

LEMMA 4.1.

- (1) Both X and Y are geometrically integral and smooth.
- (2) We have $g_X = g_Y > 1$.
- (3) We have $\text{Bir } X_L = \{1\}$ and $\text{Bir } Y_L = \{1\}$ for every field $L \supseteq \mathbb{Q}$.
- (4) Even after base field extension, there is no birational map from Y to any curve definable over a finite extension of $\mathbb{Q}$.
- (5) We have $X(\mathbb{Q}) = \emptyset$.
- (6) The first coordinate projection $\mathbb{R}^2 \rightarrow \mathbb{R}$ maps $X(\mathbb{R})$ onto $\mathbb{R}$; that is, $u(X(\mathbb{R})) = \mathbb{R}$.
- (7) There exists an open neighborhood U of 0 in $\mathbb{R}$ such that for each $t \in U$ we have $Y_t(\mathbb{R}) = \emptyset$.
- (8) If $t \in [20, \infty)$, then Y_t has a real point with x -coordinate in $[1, 2]$.

PROOF. The projective closure $\tilde{X}$ of X reduces modulo 5 to the curve

$$\tilde{X}_5: u^4 + uv^3 + vw^3 + w^4 = 0$$

in $\mathbb{P}_{\mathbb{F}_5}^2$. The projective closure of Y specializes at $T = 0$ and $T = \infty$ to the curves

$$\tilde{Y}_0: x^4 + y^4 + z^4 = 0 \quad \text{and} \quad \tilde{Y}_\infty: x^4 + xy^3 + yz^3 + z^4 = 0$$

in $\mathbb{P}_{\mathbb{Q}}^2$, respectively (to see the latter, divide the equation $q(T, x, y) = 0$ by T before setting $1/T$ equal to 0). By [Poo05, Case I with $n = 2$, $d = 4$, $c = 1$], $\tilde{X}_5$ and $\tilde{Y}_\infty$ are smooth, projective, geometrically integral plane curves of genus 3 with no nontrivial birational automorphisms even after base extension. It follows that X and Y have the same properties, except for being projective. (Proof: The projective curve $\tilde{X}$ must be smooth, since a singularity on $\tilde{X}$ would reduce to a singularity on $\tilde{X}_5$. If $\tilde{X}$ were not geometrically integral, its equation would factor nontrivially over some extension of $\mathbb{Q}$, and then reducing coefficients would show that the equation of $\tilde{X}_5$ would factor nontrivially over some extension of $\mathbb{F}_5$, contradicting the fact that $\tilde{X}_5$ is geometrically integral. The open subvariety X inherits the properties of being smooth and geometrically integral from $\tilde{X}$. The curves X and $\tilde{X}$ have the same genus as $\tilde{X}_5$. Lemma A.5 yields an injection $\text{Bir } X \simeq \text{Bir } \tilde{X} \hookrightarrow \text{Bir } \tilde{X}_5 = \{1\}$, so $\text{Bir } X = \{1\}$. The same arguments apply to Y and $\tilde{Y}_\infty$.)

- (1) Explained above.
- (2) By the above, $g_X = g_Y = 3$.
- (3) Explained above.
- (4) If there were such a birational map, then the specializations $\tilde{Y}_0$ and $\tilde{Y}_\infty$ would be birational over $\overline{\mathbb{Q}}$. But the former has a nontrivial automorphism $(x, y, z) \mapsto (-x, y, z)$.
- (5) If $\tilde{X}$ had a $\mathbb{Q}$ -point, its homogeneous coordinates could be scaled to relatively prime integers, and reducing the equation modulo 8 would show that $u^4 + 2v^4 + 4w^4 \equiv 0 \pmod{8}$ has a solution with at least one of u, v, w odd. But 4th powers are 0 or 1 modulo 8, so this is a contradiction. Thus $\tilde{X}(\mathbb{Q}) = \emptyset$, so $X(\mathbb{Q}) = \emptyset$.
- (6) Fix $u \in \mathbb{R}$. Then $\lim_{v \rightarrow +\infty} p(u, v) = +\infty$, so by the intermediate value theorem it suffices to find $v \in \mathbb{R}$ such that $p(u, v) < 0$. If $|u| < \sqrt{2}$, then $v = 0$ works. If $|u| \geq \sqrt{2}$, then $p(u, -u) = -5u^4 - 16u - 4 < 0$ by calculus.
- (7) This follows from $\tilde{Y}_0(\mathbb{R}) = \emptyset$ and compactness.
- (8) Suppose that $t \geq 20$ and $x \in [1, 2]$. Then $q(t, x, -3) \leq 2^4 + 3^4 + 1 + 20(2^4 - 27 - 3 + 1) < 0$ but $q(t, x, 0) > 0$, so there exists $y \in \mathbb{R}$ with $q(t, x, y) = 0$.

—

§5. Construction of the functor.

5.A. Construction. We now define a functor $\mathcal{F}: \mathbf{Graphs} \rightarrow \mathbf{Fields}$. Suppose that we are given a graph $G = (V, E)$. Let $K = \mathbb{Q}(\prod_{i \in V} X)$. Let $u_i, v_i \in K$ correspond to the rational functions u, v on the i th copy of X . Thus $(u_i)_{i \in V}$ is a transcendence basis for $K/\mathbb{Q}$. For $\{i, j\} \in \binom{V}{2}$, define the K -curve $Z_{\{i, j\}}$ as $Y_{u_i u_j}$ if $\{i, j\} \in E$, and $Y_{u_i + u_j}$ if $\{i, j\} \notin E$. Define $\mathcal{F}(G) := K(\prod Z_{\{i, j\}})$, where the product is over $\{i, j\} \in \binom{V}{2}$. A morphism $G \rightarrow G'$ induces an obvious field homomorphism $\mathcal{F}(G) \rightarrow \mathcal{F}(G')$. We obtain a functor $\mathcal{F}$.

Remark 5.1. If G is finite, then $\#\mathcal{F}(G) = \aleph_0$. If G is infinite, then $\#\mathcal{F}(G) = \#G$.

5.B. Properties. Here we prove properties of $\mathcal{F}(G)$ that will enable us to recover G from $\mathcal{F}(G)$, or more precisely, to prove that $\mathcal{F}$ is fully faithful. In the proofs in this section, labels like (2) refer to the parts of Lemma 4.1.

LEMMA 5.2. *Fix G . Let L be any field extension of K . Consider the base changes to L of X and all the curves $Y_{u_i u_j}$ and $Y_{u_i + u_j}$. The only nonconstant rational maps between these curves over L are the identity maps from one of them to itself.*

PROOF. By (2), all the curves have the same genus. By (3) and Lemma A.2, it suffices to show that no two distinct curves in this list are birational even after base field extension. By (4), this is already true for X and Y_t for any transcendental t . If t, t' are algebraically independent over $\mathbb{Q}$, and Y_t and $Y_{t'}$ become birational after base field extension, then we can specialize t' to an element of $\overline{\mathbb{Q}}$ while leaving t transcendental, contradicting (4). The previous two sentences apply in particular to any t and t' taken from the $u_i u_j$ and the $u_i + u_j$. $\dashv$

LEMMA 5.3. *Let $G = (V, E)$ be a graph. Let $x_{ij}, y_{ij} \in \mathcal{F}(G)$ correspond to the rational functions x, y on $Z_{\{i, j\}}$.*

- (i) *We have $X(\mathcal{F}(G)) = \{(u_i, v_i) : i \in V\}$.*
- (ii) *If $\{i, j\} \in E$, then $Y_{u_i u_j}(\mathcal{F}(G)) = \{(x_{ij}, y_{ij})\}$ and $Y_{u_i + u_j}(\mathcal{F}(G)) = \emptyset$.*
- (iii) *If $\{i, j\} \notin E$, then $Y_{u_i u_j}(\mathcal{F}(G)) = \emptyset$ and $Y_{u_i + u_j}(\mathcal{F}(G)) = \{(x_{ij}, y_{ij})\}$.*

PROOF.

- (i) By definition, $\mathcal{F}(G)$ is the direct limit of $K(Z)$, where Z ranges over finite products of the $Z_{\{i, j\}}$. Thus, by Lemma A.1, each point in $X(\mathcal{F}(G))$ corresponds to a rational map from some Z to the base change X_K . By Lemmas A.3 and 5.2 every such rational map is constant. In other words, $X(\mathcal{F}(G)) = X(K)$.

Similarly, by Lemma A.1, each point in $X(K)$ corresponds to a rational map from some finite power of X to X . By (5), the rational map is nonconstant. By Lemmas A.3 and 5.2 it is the i th projection for some i . The corresponding point in $X(K)$ is (u_i, v_i) .

- (ii) Suppose that $\{i, j\} \in E$. The same argument as for (i) shows that $Y_{u_i u_j}(\mathcal{F}(G)) = Y_{u_i u_j}(K) \cup \{(x_{ij}, y_{ij})\}$, the last point coming from the identity $Z_{\{i, j\}} \rightarrow Y_{u_i u_j}$. By (6), we may embed K in $\mathbb{R}$ so that the u_i are mapped to algebraically independent real numbers so close to zero that $Y_{u_i u_j}(\mathbb{R}) = \emptyset$ by (7). Thus $Y_{u_i u_j}(K) = \emptyset$. So $Y_{u_i u_j}(\mathcal{F}(G)) = \{(x_{ij}, y_{ij})\}$.

The argument for $Y_{u_i + u_j}(\mathcal{F}(G)) = \emptyset$ is the same, except now that $Z_{\{i, j\}}$ is not birational to $Y_{u_i u_j}$.

- (iii) The argument is the same as for (ii). $\dashv$

§6. Construction of the inverse functor.

6.A. Construction. Let $\mathbf{E}$ be the essential image of $\mathcal{F}$. We may view $\mathcal{F}$ as a functor $\mathbf{Graphs} \rightarrow \mathbf{E}$. We now construct an essential inverse $\mathcal{G} : \mathbf{E} \rightarrow \mathbf{Graphs}$ of $\mathcal{F}$.

Suppose that F is an object of $\mathbf{E}$. Define $V := X(F)$. For $i \in V = X(F)$, let u_i be the first coordinate of i . For $\{i, j\} \in \binom{V}{2}$, Lemma 5.3(ii,iii) shows that exactly one of $Y_{u_i u_j}$ and $Y_{u_i + u_j}$ has an F -point. Let E be the set of $\{i, j\} \in \binom{V}{2}$ for which it is $Y_{u_i u_j}$ that has a F -point. Let $\mathcal{G}(F)$ be the graph (V, E) .

Suppose that $f : F \rightarrow F'$ is a morphism of $\mathbf{E}$. We want to define a morphism of graphs $g : \mathcal{G}(F) \rightarrow \mathcal{G}(F')$. On vertices, g is the map $X(F) \rightarrow X(F')$ induced by f . If $\{i, j\}$ is an edge of $\mathcal{G}(F)$, then $Y_{u_i u_j}$ has an F -point, and applying f shows that $Y_{f(u_i) f(u_j)}$ has an F' -point, so $\{g(i), g(j)\}$ is an edge of $\mathcal{G}(F')$. If $\{i, j\}$ is *not* an edge of $\mathcal{G}(F)$, then the analogous argument using $Y_{u_i + u_j}$ shows that $\{g(i), g(j)\}$ is *not* an edge of $\mathcal{G}(F')$. Thus g is a morphism of graphs. Define $\mathcal{G}(f) := g$. This completes the specification of a functor $\mathcal{G}$.

6.B. Properties. For each graph G , let $\epsilon_G : G \rightarrow X(\mathcal{F}(G)) = \mathcal{G}(\mathcal{F}(G))$ be the map sending i to (u_i, v_i) .

PROPOSITION 6.1. *We have $\mathcal{G}\mathcal{F} \simeq 1_{\mathbf{Graphs}}$.*

PROOF. By Lemma 5.3, ϵ_G is a bijection, and in fact a graph isomorphism since the following are equivalent for a pair $\{i, j\}$:

- $\{i, j\}$ is an edge of G ;

- $Y_{u_i u_j}(\mathcal{F}(G)) \neq \emptyset$;
- There is an edge between the vertices (u_i, v_i) and (u_j, v_j) of $\mathcal{G}(\mathcal{F}(G))$.

The isomorphism ϵ_G varies functorially with G . $\dashv$

PROPOSITION 6.2. *The functor $\mathcal{G}$ is faithful.*

PROOF. We must show how to recover a morphism $f: F \rightarrow F'$ in $\mathbf{E}$ given F , F' , and $\mathcal{G}(f)$. Without loss of generality, replace F by an isomorphic field to assume that $F = \mathcal{F}(G)$, which is generated by elements u_i, v_i, x_{ij}, y_{ij} . Since $\mathcal{G}(f)$ is the map $X(F) \rightarrow X(F')$ induced by f , we recover $f(u_i)$ and $f(v_i)$ for all i . For each edge $\{i, j\}$ of G , the homomorphism f maps $(x_{ij}, y_{ij}) \in Y_{u_i u_j}(F)$ to a point of $Y_{u_i u_j}(F')$, but by Lemma 5.3(ii) that point is unique, so we recover $f(x_{ij})$ and $f(y_{ij})$. Similarly, for each non-edge $\{i, j\}$ of G , Lemma 5.3(iii) lets us recover $f(x_{ij})$ and $f(y_{ij})$. Together, these determine f . $\dashv$

PROOF OF THEOREM 1.1. Propositions 6.1 and 6.2 formally imply that $\mathcal{F}$ is fully faithful. $\dashv$

§7. Computability. The specification of $\mathcal{F}$ in Section 5.A sufficed for Theorem 1.1. But for the applications to computable model theory, for $G \in \mathbf{Graphs}_\omega$ we need to modify the definition of $\mathcal{F}(G)$ to ensure in particular that its domain is ω and not some other countable set. To do this, we will iteratively build a standard presentation of a function field starting from a presentation of its constant field.

Let $\pi: \omega \times \omega \xrightarrow{\sim} \omega$ be the standard pairing function. We will repeatedly use the following: Given a field k with $\text{dom}(k) \subseteq \omega$, and given an irreducible polynomial $f(x, y) \in k[x, y]$ of y -degree $d \in \mathbb{Z}_{>0}$, the function field of the integral curve $f(x, y) = 0$ is

$$k(x)[y]/(f(x, y)) \simeq k(x) \oplus k(x)y \oplus \cdots \oplus k(x)y^{d-1},$$

and the presentation of k can be extended to a presentation of this function field on the disjoint union $\text{dom}(k) \sqcup \omega$.

Let $G = (\omega, E) \in \mathbf{Graphs}_\omega$. Partition the ω that is to be $\text{dom}(\mathcal{F}_\omega(G))$ into countably many infinite columns $\omega^{[n]} := \pi(\{n\} \times \omega)$. Define field operations on $K_0 := \omega^{[0]}$ so that K_0 is computably isomorphic to $\mathbb{Q}$. For each $i \in \omega$, use the elements of $\omega^{[2i+2]}$ to extend the field K_i to $K_{i+1} := K_i(X) = K_i(u_{i+1}, v_{i+1})$ in a uniform way such that $\pi(u_{i+1}, v_{i+1}) > \pi(u_i, v_i)$. Then $K \simeq \bigcup K_i$, which has domain $\bigcup_{n \text{ even}} \omega^{[n]}$. Let $F_0 := K$. Next, order the pairs (i, j) with $i > j$ in lexicographic order, hence in order type ω . If (i, j) is the k th such pair (so $k = \frac{i(i-1)}{2} + j + 1$), inductively define F_k by adjoining the elements of $\omega^{[2k-1]}$ to F_{k-1} in a uniform way to form the function field $F_{k-1}(Z_{\{i,j\}})$. Then define $\mathcal{F}_\omega(G) := \bigcup F_k$, which has domain $\bigcup \omega^{[n]} = \omega$. The action of $\mathcal{F}_\omega$ on morphisms is defined as for $\mathcal{F}$.

Really all we have done is to identify $\text{dom}(\mathcal{F}(G))$ with ω . In fact, we might as well modify the definition of $\mathcal{F}$ so that $\text{dom}(\mathcal{F}(G)) = \omega$. Then $\mathcal{F}$ restricts to $\mathcal{F}_\omega$.

We record a property of the construction that will be useful in the proof of Theorem 9.7:

LEMMA 7.1. *Fix $G \in \mathbf{Graphs}_\omega$. Let $\mu: G \rightarrow \mathcal{F}_\omega(G)$ be the injection sending i to u_i . Then μ is computable, its range $\mu(G)$ is a computable subset of $\mathcal{F}_\omega(G) = \omega$, and μ^{-1} (defined on $\mu(G)$) is computable. Moreover, μ varies functorially with G .*

PROOF. The uniformity in the construction of the K_i in $\mathcal{F}_\omega(G)$ ensures that μ is computable even when G is not. (In fact, μ is independent of G .) Given $j \in \mathcal{F}_\omega(G)$, we can determine if $j \in \omega^{[2i+2]}$ for some $i \in \omega$; if so, then compute u_i , and check if $j = u_i$. This lets us check if $j \in \mu(G)$, and if so computes i such that $\mu(i) = j$. $\dashv$

For $F \in \mathbf{Fields}_\omega$, the injection

$$X(F) \subseteq F \times F = \omega \times \omega \xrightarrow{\pi} \omega$$

defines a well-ordering on the set $X(F)$.

LEMMA 7.2. *If $F \in \mathbf{E}_\omega$, then there is an order-preserving bijection $\delta_F: \omega \rightarrow X(F)$, and δ_F and δ_F^{-1} are computable uniformly from an F -oracle.*

PROOF. We have $F \simeq \mathcal{F}_\omega(G)$ for some $G \in \mathbf{Graphs}_\omega$. By Lemma 5.3(i), $\#X(F) = \#X(\mathcal{F}_\omega(G)) = \#G = \aleph_0$. Given F , the elements of $X(F)$ can be enumerated by searching in order; call the i th element $\delta_F(i)$ (starting with $i = 0$). This defines δ_F and shows that δ_F and δ_F^{-1} are computable. $\dashv$

LEMMA 7.3. *If $G \in \mathbf{Graphs}_\omega$, then the bijection $\epsilon_G: G \rightarrow X(\mathcal{F}_\omega(G))$ is order-preserving, and $\epsilon_G = \delta_{\mathcal{F}_\omega(G)}$.*

PROOF. The bijection ϵ_G is order-preserving by the condition $\pi(u_{i+1}, v_{i+1}) > \pi(u_i, v_i)$. Since ϵ_G and $\delta_{\mathcal{F}_\omega(G)}$ are both order-preserving bijections $\omega \rightarrow X(\mathcal{F}_\omega(G))$, they are equal. $\dashv$

Let us now define the promised “inverse” functor $\mathcal{G}_\omega: \mathbf{E}_\omega \rightarrow \mathbf{Graphs}_\omega$. If F is an object of $\mathbf{E}_\omega$, then by transport of structure across the bijection $\delta_F: \omega \rightarrow X(F)$, the graph $\mathcal{G}(F)$ with vertex set $X(F)$ becomes a graph $\mathcal{G}_\omega(F)$ with vertex set ω . If $f: F \rightarrow F'$ is a morphism of $\mathbf{E}_\omega$, then again by transport of structure, the morphism $\mathcal{G}(f): \mathcal{G}(F) \rightarrow \mathcal{G}(F')$ becomes $\mathcal{G}_\omega(f): \mathcal{G}_\omega(F) \rightarrow \mathcal{G}_\omega(F')$.

PROPOSITION 7.4. *We have $\mathcal{G}_\omega \mathcal{F}_\omega = 1_{\mathbf{Graphs}_\omega}$.*

PROOF. If G is an object of $\mathbf{Graphs}_\omega$, the composition of sets

$$G \xrightarrow{\epsilon_G} X(\mathcal{F}_\omega(G)) \xrightarrow{\delta_{\mathcal{F}_\omega(G)}^{-1}} \mathcal{G}_\omega(\mathcal{F}_\omega(G))$$

is the identity $\omega \rightarrow \omega$ by Lemma 7.3. Each step is functorial in G by construction. $\dashv$

PROPOSITION 7.5. *The functor $\mathcal{G}_\omega: \mathbf{E}_\omega \rightarrow \mathbf{Graphs}$ is computable.*

PROOF. Given $F \in \mathbf{E}_\omega$, the construction of $\mathcal{G}_\omega(F)$ is effective, as we now explain. To compute whether a given pair $\{i, j\}$ is an edge of $\mathcal{G}_\omega(F)$, first use Lemma 7.2 to compute $(u_i, v_i) := \delta_F(i)$ and $(u_j, v_j) := \delta_F(j)$. By Lemma 5.3(ii,iii), exactly one of $Y_{u_i u_j}$ and $Y_{u_i + u_j}$ has an F -point. To find out which, search both curves in parallel. When a point is found, which curve it is on tells us whether $\{i, j\}$ is an edge.

Given fields F, F' and a morphism $f: F \rightarrow F'$ in $\mathbf{E}_\omega$, the morphism $\mathcal{G}_\omega(f)$ is the composition

$$\omega \xrightarrow{\delta_F} X(F) \xrightarrow{f} X(F') \xrightarrow{\delta_{F'}^{-1}} \omega,$$

which is computable by Lemma 7.2. $\dashv$

Next is an effective version of Proposition 6.2.

PROPOSITION 7.6. *There exists a Turing functional that given $F, F' \in \mathbf{E}$ and $g: \mathcal{G}_\omega(F) \rightarrow \mathcal{G}_\omega(F')$ computes the unique morphism $f: F \rightarrow F'$ of $\mathbf{E}_\omega$ such that $\mathcal{G}_\omega(f) = g$.*

PROOF. Existence and uniqueness of f follow from Theorem 1.1. Define $u_i, v_i \in F$ by $(u_i, v_i) = \delta_F(i) \in X(F)$. Since $\mathcal{G}_\omega(f) = g$, the map $X(F) \rightarrow X(F')$ induced by f is the composition

$$X(F) \xrightarrow{\delta_F^{-1}} \mathcal{G}_\omega(F) \xrightarrow{g} \mathcal{G}_\omega(F') \xrightarrow{\delta_{F'}^{-1}} X(F').$$

Thus we may compute $f(u_i)$ and $f(v_i)$ for any given $i \in \omega$. For each $\{i, j\}$, let (x_{ij}, y_{ij}) be the point of $Y_{u_i u_j}(F)$ or $Y_{u_i + u_j}(F)$, according to whether $\{i, j\}$ is an edge of $\mathcal{G}_\omega(F)$ or not. Then f maps (x_{ij}, y_{ij}) to the unique point of $Y_{f(u_i)f(u_j)}(F')$ or $Y_{f(u_i)+f(u_j)}(F')$, and that point can be found by a search, so we can compute $f(x_{ij})$ and $f(y_{ij})$. Finally, given any $z \in F$, search for u 's, v 's, x 's, y 's as above and a rational function expressing z in terms of them; evaluate the same rational function on their images under f to compute $f(z)$ in F' . $\dashv$

PROPOSITION 7.7. *The functor $\mathcal{F}_\omega: \mathbf{Graphs}_\omega \rightarrow \mathbf{Fields}_\omega$ is computable.*

PROOF. The constructions of the fields K_i and F_i in Section 5 are done by a uniform process, so $\mathcal{F}_\omega(G)$ has been defined uniformly effectively below a $\Delta(G)$ -oracle (specifically, below the set E of edges in G). This provides the Turing functional Φ in Definition 3.1) for $\mathcal{F}_\omega$.

Now suppose that we are given graphs G and G' (or rather, their atomic diagrams) and a morphism $g: G \rightarrow G'$. By the previous paragraph, $\mathcal{F}_\omega(G)$ and $\mathcal{F}_\omega(G')$ are computable from these. Also, $\mathcal{G}_\omega(\mathcal{F}_\omega(g))$ is computable, since by Proposition 7.4 it equals g . By Proposition 7.6, we can compute $\mathcal{F}_\omega(g)$. $\dashv$

PROPOSITION 7.8. *The composition $\mathcal{F}_\omega \mathcal{G}_\omega$ is computably isomorphic to $1_{\mathbf{E}_\omega}$.*

PROOF. For each $F \in \mathbf{E}_\omega$, Proposition 7.4 yields $\mathcal{G}_\omega(\mathcal{F}_\omega(\mathcal{G}_\omega(F))) = \mathcal{G}_\omega(F)$. Applying Proposition 7.6 to the equality in each direction yields an isomorphism $\mathcal{F}_\omega(\mathcal{G}_\omega(F)) \rightarrow F$ and its inverse, and shows that they are computable from F . Moreover, the construction is functorial in F . $\dashv$

PROOF OF THEOREM 1.2. Combine Propositions 7.7, 7.5, 7.4, and 7.8. $\dashv$

PROOF OF PROPOSITION 1.6.

- (a) Theorem 1.2(b) shows that $\mathcal{F}_\omega: \mathbf{Graphs}_\omega \rightarrow \mathbf{Fields}_\omega$ is fully faithful; from this, the result follows formally.
- (b) By Corollary 1.3, there exists $G \in \mathcal{J}$ with $G \leq_T F$.

First suppose that G is automorphically trivial. Then so is every other $G' \in \mathcal{J}$. For any automorphically trivial G' , the presence of $\{i, j\}$ as an edge in G' is determined by the answers to the questions of the form “Is $i = m$?” or “Is $j = m$?” for m in some finite set, so G' is computable.

Now suppose instead that G is not automorphically trivial. Trivially, $\text{Spec } G$ contains $\deg G$. A theorem of Knight, namely [Kni86, Theorem 4.1], states that if a structure is not automorphically trivial, then its spectrum is upwards closed. Thus $\text{Spec } G$ contains $\deg F$ too. In other words, there exists $G' \in \mathcal{J}$ with $\deg G' = \deg F$. $\dashv$

§8. Ordered fields.

PROPOSITION 8.1. *For each $G \in \mathbf{Graphs}_\omega$, the field $\mathcal{F}_\omega(G)$ is isomorphic to a subfield of $\mathbb{R}$. Moreover, the field homomorphism $\mathcal{F}_\omega(G) \hookrightarrow \mathbb{R}$ may be chosen so that the induced ordering on $\mathcal{F}_\omega(G)$ is computable from G .*

PROOF. Given a subfield $k \subseteq \mathbb{R}$, an integral curve Z over k , and a point $z \in Z(\mathbb{R})$ having at least one coordinate transcendental over k , we obtain a k -embedding $k(Z) \hookrightarrow \mathbb{R}$ by sending f to $f(z)$. We will use this observation inductively to show that all the fields K_i and F_i in Section 5 embed into $\mathbb{R}$.

First, $K_0 := \mathbb{Q}$ is a subfield of $\mathbb{R}$. By the Lindemann–Weierstrass theorem, the numbers $e_i := \exp(2^{1/i})$ for $i \in \{2, 3, \dots\}$ are algebraically independent over $\mathbb{Q}$. By Lemma 4.1(6), X has a real point with u -coordinate $10e_{2j+2}$; choose the one with least v -coordinate. This lets us inductively extend $K_{j-1} \hookrightarrow \mathbb{R}$ to $K_j = K_{j-1}(X) \hookrightarrow \mathbb{R}$ so that u_j maps to $10e_{2j+2}$. Taking the union embeds $F_0 = K$ in $\mathbb{R}$. For the k th pair $i > j$, we have $u_i u_j, u_i + u_j \in [20, \infty)$ and $e_{2k+3} \in [1, 2]$, so by Lemma 4.1(8), the curve $Z_{\{i, j\}}$ has a real point with x -coordinate e_{2k+3} ; choose the one with least y -coordinate. Thus for each $k \geq 1$ we may extend the embedding $F_{k-1} \hookrightarrow \mathbb{R}$ to $F_k \hookrightarrow \mathbb{R}$. Taking the union yields an embedding $\mathcal{F}_\omega(G) \hookrightarrow \mathbb{R}$.

Given two distinct elements of $\mathcal{F}_\omega(G)$, expressed in terms of the generators of $\mathcal{F}_\omega(G)$, we may compute them numerically until we determine which is greater. $\dashv$

Unfortunately, Proposition 8.1 implies neither that $\mathcal{F}_\omega$ is a functor to the category of ordered fields (with order-preserving field homomorphisms), nor that our completeness results hold for ordered fields. The problem is that isomorphisms in the category of ordered fields are much more restricted than isomorphisms in the category of fields. Indeed, the following result was noted by Fried and Kollár in [FK82].

PROPOSITION 8.2. *There is no faithful functor from $\mathbf{Graphs}_\omega$ to the category of ordered fields, or even to the category of ordered sets.*

PROOF. Every (order-preserving) automorphism of a finite totally ordered set is trivial. Therefore every finite-order automorphism of a totally ordered set acts trivially on every orbit, and hence is trivial. On the other hand, there exist countable graphs with nontrivial finite-order automorphisms. $\dashv$

Fried and Kollár also proved the following:

THEOREM 8.3 ([FK82, Theorem 2.4]). *Fix a group G and a subgroup $H \subseteq G$. Then H is right orderable (as a group) if and only if there is an ordered field F and an isomorphism ϕ mapping G onto the group of field automorphisms of F such that the image $\phi(H)$ contains precisely the order-preserving automorphisms of F .*

§9. Consequences in computable model theory. Many of the results below rely on [HKSS02, Theorem 3.1], which shows that for every countable structure $\mathcal{A}$ that is not automorphically trivial, there exists a graph $G \in \mathbf{Graphs}_\omega$ with the same spectrum and the same $\mathbf{d}$ -computable dimension for each Turing degree $\mathbf{d}$, and with the property that for every relation R on $\mathcal{A}$, there exists a relation on G with the same degree spectrum as R on $\mathcal{A}$.

9.A. Turing degree spectrum. Recall from Section 2.D the definition of the spectrum:

$$\text{Spec } M := \{\deg N : N \simeq M \text{ and } \text{dom}(N) = \omega\}.$$

PROPOSITION 9.1.

- (a) If $G \in \mathbf{Graphs}_\omega$, then $G \equiv_T \mathcal{F}_\omega(G)$.
- (b) If $g: G \rightarrow G'$ is a morphism between computable graphs in $\mathbf{Graphs}_\omega$, then $g \equiv_T \mathcal{F}_\omega(g)$.

PROOF. Theorem 1.2 showed that $\mathcal{F}_\omega$ and $\mathcal{G}_\omega$ are computable functors. ⊢

THEOREM 9.2. Let $G \in \mathbf{Graphs}_\omega$.

- (a) If G is automorphically trivial, then $\text{Spec } G = \{\mathbf{0}\}$, and $\text{Spec } \mathcal{F}_\omega(G)$ contains all Turing degrees.
- (b) If G is not automorphically trivial, then $\text{Spec } G = \text{Spec } \mathcal{F}_\omega(G)$.

PROOF.

- (a) Every automorphically trivial graph is computable, so $\text{Spec } G = \{\mathbf{0}\}$. The field $\mathcal{F}_\omega(G)$ is computable but not automorphically trivial; Knight's theorem [Kni86, Theorem 4.1] applied to $\mathcal{F}_\omega(G)$ yields the result.
- (b) Applying Proposition 9.1(a) to every G' isomorphic to G yields $\text{Spec } G \subseteq \text{Spec } \mathcal{F}_\omega(G)$. On the other hand, since G is not automorphically trivial, Proposition 1.6(b) yields $\text{Spec } \mathcal{F}_\omega(G) \subseteq \text{Spec } G$. ⊢

COROLLARY 9.3. For every countable structure $\mathcal{A}$ that is not automorphically trivial, there exists $F \in \mathbf{Fields}_\omega$ with $\text{Spec } F = \text{Spec } \mathcal{A}$.

PROOF. Since $\mathcal{A}$ is not automorphically trivial, [HKSS02, Theorem 3.1] yields some $G \in \mathbf{Graphs}_\omega$ such that $\text{Spec } G = \text{Spec } \mathcal{A} \neq \{\mathbf{0}\}$. By Knight's theorem, G is not automorphically trivial, so Theorem 9.2 shows $\text{Spec } G = \text{Spec } \mathcal{F}_\omega(G)$. Let $F := \mathcal{F}_\omega(G)$. ⊢

9.B. Computable categoricity. Let $\mathcal{A}$ be a computable structure, let $\mathbf{d}$ be a Turing degree, and let α be a computable ordinal. If $\mathcal{B}$ is any structure on ω , let $\mathcal{B}^{(\alpha)}$ denote the α -jump of the atomic diagram of $\mathcal{B}$. Let $\text{Isom}_{\mathbf{d}}(\mathcal{A})$ be the set of $\mathbf{d}$ -computable isomorphism classes in the set of computable structures isomorphic to $\mathcal{A}$; then the cardinal $\#\text{Isom}_{\mathbf{d}}(\mathcal{A}) \in \{1, 2, \dots, \aleph_0\}$ is called the $\mathbf{d}$ -computable dimension of $\mathcal{A}$ [HKSS02, Definition 1.2]. The categoricity spectrum of $\mathcal{A}$ is the set of Turing degrees $\mathbf{d}$ such that the $\mathbf{d}$ -computable dimension of $\mathcal{A}$ is 1. Finally, $\mathcal{A}$ is relatively computably categorical if every structure $\mathcal{B} \cong \mathcal{A}$ with domain ω is $\mathcal{B}$ -computably isomorphic to $\mathcal{A}$, and relatively Δ_α^0 -categorical if every structure $\mathcal{B} \cong \mathcal{A}$ with domain ω is $\mathcal{B}^{(\alpha)}$ -computably isomorphic to $\mathcal{A}$.

THEOREM 9.4. For every computable structure $\mathcal{A}$, there exists a computable field F such that

- (i) for each Turing degree $\mathbf{d}$, the field F has the same $\mathbf{d}$ -computable dimension as $\mathcal{A}$;
- (ii) F has the same categoricity spectrum as $\mathcal{A}$; and
- (iii) for every computable ordinal α , the field F is relatively Δ_α^0 -categorical if and only if $\mathcal{A}$ is.

PROOF. First suppose that $\mathcal{A}$ is automorphically trivial. Let $F = \mathbb{Q}$. Then $\mathcal{A}$ has $\mathbf{d}$ -computable dimension 1 for every $\mathbf{d}$, and $\mathcal{A}$ is relatively Δ_α^0 -categorical for all α , and the field $\mathbb{Q}$ has the same properties.

From now on, suppose that $\mathcal{A}$ is not automorphically trivial. Use [HKSS02, Theorem 3.1] to replace $\mathcal{A}$ by a computable graph G on domain ω , and let $F = \mathcal{F}_\omega(G)$.

(i) The functors $\mathcal{F}_\omega$ and $\mathcal{G}_\omega$ are computable (Theorem 1.2), so they map computable objects of $\mathbf{Graphs}_\omega$ to computable objects of $\mathbf{E}_\omega$ and vice versa, and they respect isomorphism and $\mathbf{d}$ -computable isomorphism of such objects (it will be OK to work in $\mathbf{E}_\omega$ instead of $\mathbf{Fields}_\omega$ since all fields in $\mathbf{Fields}_\omega$ isomorphic to F are in $\mathbf{E}_\omega$). Thus they induce maps between the sets $\text{Isom}_{\mathbf{d}}(G)$ and $\text{Isom}_{\mathbf{d}}(F)$. The composition of these maps in either order is the identity since $\mathcal{G}_\omega(\mathcal{F}_\omega(G'))$ is computably isomorphic (in fact, equal) to G' for every

$G' \in \mathbf{Graphs}_\omega$ and $\mathcal{F}_\omega(\mathcal{G}_\omega(F'))$ is computably isomorphic to F' for every $F' \in \mathbf{E}_\omega$ (Theorem 1.2(b)). Thus $\# \text{Isom}_d(G) = \# \text{Isom}_d(F)$.

(ii) This follows from (i), since the categoricity spectrum is defined as the set of d such that the d -computable dimension equals 1.

(iii) Fix α . If G is not relatively Δ_α^0 -categorical, then some graph G' violates the condition in the definition above, and $\mathcal{F}_\omega(G)$ and $\mathcal{F}_\omega(G')$ violate the same condition, so $\mathcal{F}_\omega(G)$ is not relatively Δ_α^0 -categorical either. On the other hand, if it holds of G , then it holds immediately of $\mathcal{F}_\omega(G)$ and every other field $\mathcal{F}_\omega(G')$ with $G' \simeq G$. But for a field $F \simeq \mathcal{F}_\omega(G)$, Corollary 1.3 shows that there exists an F -computable isomorphism $f: F \rightarrow \mathcal{F}_\omega(H)$ for some graph H that is isomorphic to G and Turing-computable from F . Therefore, there is a $(\deg H)^{(\alpha)}$ -computable (hence $(\deg F)^{(\alpha)}$ -computable) isomorphism g from H onto G , whose image $\mathcal{F}_\omega(g)$ is a $(\deg F)^{(\alpha)}$ -computable isomorphism from $\mathcal{F}_\omega(H)$ onto $\mathcal{F}_\omega(G)$. So the composition $\mathcal{F}_\omega(g) \circ f$ is a $(\deg F)^{(\alpha)}$ -computable isomorphism from F onto $\mathcal{F}_\omega(G)$, proving that $\mathcal{F}_\omega(G)$ is also relatively Δ_α^0 -categorical. $\dashv$

COROLLARY 9.5. *Fields realize all computable dimensions $\leq \omega$.*

PROOF. Theorems of Goncharov [Gon77, Gon80] and others have shown long since that computable structures (in fact, graphs) can have every computable dimension from 1 up to $\aleph_0$. (See [AK00, §12.6] for a summary of these results using the terminology of this article.) Theorem 9.4 allows us to carry this over to fields. $\dashv$

The basic definition of computable categoricity of a computable structure $\mathcal{A}$ shows computable categoricity to be a Π_1^1 property: it is defined by a statement using a universal quantifier over *sets* of natural numbers. One can view it as saying that, for all subsets f of ω^2 and all $e \in \omega$, either the e th computable function φ_e fails to define a computable structure, or the function (if any) defined on ω by f fails to be an isomorphism from $\mathcal{A}$ onto the structure defined by φ_e , or else there exists a Turing program which computes an isomorphism from that structure onto $\mathcal{A}$. All quantifiers over natural numbers can be absorbed into the universal quantifier over sets, so we view this formula as universal over sets, i.e., Π_1^1 , with the superscript 1 signifying quantification over sets.

Sometimes a Π_1^1 property is expressible also by a simpler formula. In [DKL⁺15, Theorem 1], however, Downey, Kach, Lempp, Lewis, Montalbán, and Turetsky give a proof that computable categoricity for trees is Π_1^1 -complete. Theorem 1.2 allows us to transfer this result to fields.

THEOREM 9.6. *The property of computable categoricity for computable fields is Π_1^1 -complete.*

That is,

$$\{e \in \omega : \varphi_e \text{ computes the atomic diagram of a computably categorical field}\}$$

is a Π_1^1 set, and every Π_1^1 set is 1-reducible to this set.

In contrast, the property of computable categoricity for algebraic fields is just Π_4^0 -complete (see [HKMS15, Theorem 5.4]). For fields of infinite transcendence degree, it was shown only recently, in [MS13, Theorem 3.4], that they could be computably categorical at all, and it was not known whether they could be computably categorical without being relatively computably categorical, which is a Σ_3^0 property. So Theorem 9.6 represents a significant step forward.

PROOF. Theorem 3.1 of [HKSS02] enables us to build a computable graph corresponding to an arbitrary computable tree, effectively in the index of the tree, and then Theorem 1.2 builds a computable field that is computably categorical if and only if the graph (and hence the original tree) was. Thus we have a 1-reduction from a Π_1^1 -complete set (the indices of computably categorical trees) to the set of indices of computably categorical fields. $\dashv$

9.C. Degree spectrum. The degree spectrum $\text{DgSp}_\mathcal{A}(R)$ of a relation R on a computable structure $\mathcal{A}$ is the set of all Turing degrees of images of R under isomorphisms f from $\mathcal{A}$ onto computable structures $\mathcal{B}$. (Understand that here the relation R is generally not in the language; if it were, then this image would always be computable, just by the definition of a computable structure. The relation R may be *definable* in the language of the structure, however, in which case its definition places an upper bound on the complexity of these images.) If f is computable, then $f(R)$ is Turing-equivalent to R itself; thus it is the noncomputable

f that give this definition its teeth. Here we show that all degree spectra of relations on automorphically nontrivial computable structures can be realized as degree spectra of relations on fields:

THEOREM 9.7. *Let $\mathcal{A}$ be any computable structure that is not automorphically trivial, and let R be an n -ary relation on $\mathcal{A}$. Then there exists a field F and an n -ary relation S on F such that*

$$\text{DgSp}_{\mathcal{A}}(R) = \text{DgSp}_F(S).$$

PROOF. As usual, we appeal to [HKSS02, Theorem 3.1] to assume that $\mathcal{A}$ is in fact a graph, hereafter named G . For this proof, Theorem 1.2 is not quite enough; to transfer relations we need also Lemma 7.1, about the map $\mu: G \rightarrow \mathcal{F}_{\omega}(G)$ and its inverse. Let $F = \mathcal{F}_{\omega}(G)$, and let $S = \mu(R) \subseteq F^n$ (apply μ coordinate-wise).

Suppose that $g: G \rightarrow G'$ is an isomorphism of computable graphs in **Graphs** $_{\omega}$. In the diagram of sets

$$\begin{array}{ccc} G & \xrightarrow{\mu} & \mathcal{F}_{\omega}(G) \\ g \downarrow & & \downarrow \mathcal{F}_{\omega}(g) \\ G' & \xrightarrow{\mu} & \mathcal{F}_{\omega}(G'), \end{array}$$

R and S map downwards to relations R' on G' and S' on $\mathcal{F}_{\omega}(G')$, and $\text{DgSp}_G(R)$ consists of the degrees $\deg R'$ arising in this way from all possible g . By functoriality of μ , the diagram commutes, so the bottom horizontal μ maps R' to S' . Since μ and μ^{-1} are computable, $R' \equiv_T S'$. Thus $\deg R' = \deg S' \in \text{DgSp}_F(S)$. Hence $\text{DgSp}_G(R) \subseteq \text{DgSp}_F(S)$.

Now suppose that $f: F \rightarrow F'$ is an isomorphism of computable fields in **Fields** $_{\omega}$; it maps S to some S' ; then $\text{DgSp}_F(S)$ consists of the degrees $\deg S'$ arising in this way. By Corollary 1.4, there is a computable graph $G' \simeq G$ and a computable isomorphism $i: F' \rightarrow \mathcal{F}_{\omega}(G')$. Composing f with a computable isomorphism does not change the resulting Turing degree $\deg S'$, so we may assume that F' equals $\mathcal{F}_{\omega}(G')$. Since $\mathcal{F}_{\omega}$ is fully faithful, f is $\mathcal{F}_{\omega}(g)$ for some isomorphism $g: G \rightarrow G'$. This g maps R to some R' , and the previous paragraph shows that $\deg S' = \deg R' \in \text{DgSp}_G(R)$. Hence $\text{DgSp}_F(S) \subseteq \text{DgSp}_G(R)$. $\dashv$

Remark 9.8. It is not true that for every automorphically nontrivial computable structure $\mathcal{A}$, there exists a computable field F of characteristic 0 such that for every relation S on $\mathcal{A}$, there exists a relation R on $\mathcal{A}$ with the same degree spectrum. For example, suppose that $\mathcal{A}$ is the random graph or the countable dense linear order; then the degree spectrum of every relation on $\mathcal{A}$ either is $\{0\}$ or is upwards-closed under $\leq_T$; see [HM07, Corollary 2.11 and Proposition 3.6]. On the other hand, in every computable field F of characteristic 0, one can effectively locate each positive integer n (meaning the sum $1 + \dots + 1$ of the multiplicative identity with itself n times — not the element n of the domain ω), and therefore the unary relation S consisting of those n that lie in the Halting Problem will have degree spectrum exactly $\{0'\}$.

The restriction to automorphically nontrivial structures $\mathcal{A}$ in Theorem 9.7 can be bypassed for unary relations, since on an automorphically trivial computable structure $\mathcal{A}$, each unary relation has degree spectrum either $\{0\}$ (if the relation is either finite or cofinite) or else the set of all Turing degrees. Both of these can easily be realized as spectra of unary relations on fields. We leave the analysis of n -ary relations on such structures for another time.

Functoriality also allows one to show that a relation R on a countable graph G is relatively intrinsically Σ_{α}^0 if and only if its image $h(R)$ (defined exactly as in the proof of Theorem 9.7) is relatively intrinsically Σ_{α}^0 on the field $\mathcal{F}_{\omega}(G)$; and similarly for relations that are relatively intrinsically Π_{α}^0 , Σ_m^1 , etc. Likewise, every definable relation on G is mapped to a definable relation on $\mathcal{F}_{\omega}(G)$: this is immediate if one views our construction as a bi-interpretation between the graph and the field.

9.D. Automorphism spectrum. In [HMM10, Definition 1.1], Harizanov, Miller, and Morozov defined the automorphism spectrum of a computable structure to be the set of all Turing degrees of nontrivial automorphisms of the structure. They used [HKSS02, Theorem 3.1] to show that every automorphism spectrum is the automorphism spectrum of a computable graph. This sets up another application of Theorem 1.2.

THEOREM 9.9. *For every computable structure $\mathcal{A}$, there is a computable field F with the same automorphism spectrum as $\mathcal{A}$.*

PROOF. The theorem follows from the full faithfulness of $\mathcal{F}_\omega$, along with its preservation of Turing degrees of automorphisms (Proposition 9.1(b)). $\dashv$

§10. Alternative approach using the construction of Fried and Kollár.

10.A. Comparison of approaches. In our construction of $\mathcal{F}_\omega(G)$, we adjoined a function field for each edge *and* a function field for each non-edge (the function field of $Y_{u_i u_j}$ or $Y_{u_i + u_j}$, respectively). Doing both gave existential criteria for the existence of an edge and for non-existence of an edge, so that searching for both in parallel (see the proof of Proposition 7.5) let us computably decide whether a particular edge belonged to G . Thus we could show that an inverse functor was computable.

The construction of Fried and Kollár, on the other hand, comes with an existential criterion only for existence of each edge. Thus, although their functor can be made computable, their inverse functor is *not* computable, so it does not yield consequences in computable model theory such as Theorem 1.8. Perhaps it would not be hard to adapt their construction to yield a computable equivalence of categories from $\mathbf{Graphs}_\omega$ to its essential image, but no one has done this.

10.B. Markerization. Instead we propose a workaround based on *Markerization*. In analogy with [Mar89] (see also [FKM10, Section 5]), define the Marker $\exists$ -extension functor $\mathcal{H}: \mathbf{Graphs}_\omega \rightarrow \mathbf{Graphs}_\omega$ as follows. For $n \geq 3$, let C_n denote a rooted graph consisting of a rooted n -cycle and one more vertex with an edge to a neighbor of the root. The automorphism group of C_n as a *rooted* graph is trivial. When we speak of “attaching a copy of C_n to a vertex v of a graph G ”, we mean taking the disjoint union of C_n and G and adding an edge from the root of C_n to v . Given $G \in \mathbf{Graphs}_\omega$, attach a copy of C_3 to each vertex of G , and replace each edge (respectively, non-edge) $\{i, j\}$ of the original graph G by a 2-edge chain with a copy of C_5 (respectively, C_7) attached to the midpoint of the chain; identify the new vertex set with ω in some computable way, and call the resulting graph $\mathcal{H}(G)$. Each morphism $g: G \rightarrow G'$ in $\mathbf{Graphs}_\omega$ extends to a morphism $\mathcal{H}(g): \mathcal{H}(G) \rightarrow \mathcal{H}(G')$ that maps each attached C_n to the corresponding one.

We next define an inverse functor $\mathcal{H}^{-1}$ on the essential image of $\mathcal{H}$. Given H isomorphic to $\mathcal{H}(G)$ for some G , let V be the set of infinite-degree vertices of H that are connected by an edge to a copy of C_3 , let $E \subseteq \binom{V}{2}$ be the set of $\{i, j\}$ such that the midpoint of the 2-edge chain joining i and j is connected by an edge to a copy of C_5 , and let $\mathcal{H}^{-1}(H) = (V, E) \in \mathbf{Graphs}_\omega$ (again, identify V with ω in a computable way). Each morphism $H \rightarrow H'$ between graphs in the essential image of $\mathcal{H}$ restricts to a morphism $\mathcal{H}^{-1}(H) \rightarrow \mathcal{H}^{-1}(H')$.

Then $\mathcal{H}$ defines a fully faithful functor with inverse $\mathcal{H}^{-1}$, and $\mathcal{H}$ is a computable equivalence of categories to its essential image. Moreover, $\mathcal{H}^{-1}$ has a stronger property: given only an oracle enumerating the edge set of H (instead of an oracle capable of deciding whether a given pair is an edge), one can *compute* $\mathcal{H}^{-1}(H)$: given vertices, search for their midpoint and search for a 5-cycle or 7-cycle connected to the midpoint; eventually one or the other will be found.

10.C. Completeness properties of fields of specified characteristic. Let p be a prime or 0. Let $\mathbf{Fields}_{\omega, p}$ be the full subcategory of $\mathbf{Fields}_\omega$ consisting of fields of characteristic p with domain ω . Fried and Kollár [FK82] constructed a fully faithful functor $\mathcal{F}_p: \mathbf{Graphs}_\omega \rightarrow \mathbf{Fields}_{\omega, p}$ (for $p = 2$, this was done later, implicitly in [Prö84]). In retrospect, $\mathcal{F}_p$ is computable. Then $\mathcal{F}_p \mathcal{H}: \mathbf{Graphs}_\omega \rightarrow \mathbf{Fields}_{\omega, p}$ is a fully faithful computable functor with inverse $\mathcal{H}^{-1} \mathcal{F}_p^{-1}$ defined on the essential image $\mathbf{E}_{\omega, p}$ of $\mathcal{F}_p \mathcal{H}$. Given a field $F \in \mathbf{E}_{\omega, p}$, it is unclear whether we can compute the edge relation in the graph $\mathcal{F}_p^{-1}(F)$, but we can *enumerate* this relation, and that is sufficient for us to compute the edge relation of $\mathcal{H}^{-1}(\mathcal{F}_p^{-1}(F))$, thanks to the argument of Section 10.B. In fact, $\mathcal{F}_p \mathcal{H}: \mathbf{Graphs}_\omega \rightarrow \mathbf{E}_{\omega, p}$ is a computable equivalence of categories. In particular, we obtain the following:

THEOREM 10.1. *Let p be a prime or 0. The class of countable fields of characteristic p has the completeness property of [HKSS02, Definition 1.21].*

§Appendix A. Algebraic geometry facts.

LEMMA A.1. *If V and W are varieties over a field k , and W is integral, then $V(k(W))$ is in bijection with the set of rational maps $W \dashrightarrow V$.*

PROOF. The description of a point in $V(k(W))$ involves only finitely many elements of $k(W)$, and there is a dense open subvariety $U \subseteq W$ on which they are all regular. $\dashv$

LEMMA A.2. *Let k be a field of characteristic 0. Let C and D be geometrically integral curves over k such that $g_C = g_D > 1$. Every nonconstant rational map $C \dashrightarrow D$ is birational.*

PROOF. This is a well known consequence of Hurwitz's formula. $\dashv$

LEMMA A.3. *Let $V_1, \dots, V_n$ be geometrically integral varieties over a field k of characteristic 0. Let C be a geometrically integral curve over k such that $g_C > 1$. Then each rational map $V_1 \times \dots \times V_n \dashrightarrow C$ factors through the projection $V_1 \times \dots \times V_n \rightarrow V_i$ for at least one i .*

PROOF. By induction, we may assume that $n = 2$. We may assume that k is algebraically closed. A rational map $\phi: V_1 \times V_2 \dashrightarrow C$ may be viewed as an algebraic family of rational maps $V_1 \dashrightarrow C$ parametrized by (an open subvariety of) V_2 . By the de Franchis–Severi theorem [Lan60, pp. 29–30] there are only finitely many nonconstant rational maps $V_1 \dashrightarrow C$, so they do not vary in algebraic families. Thus either the rational maps in the family are all the same, in which case ϕ factors through the first projection, or each rational map in the family is constant, in which case ϕ factors through the second projection. $\dashv$

Remark A.4. Lemma A.3 holds even if $\text{char } k = p$. This can be deduced from the characteristic p analogue of the de Franchis–Severi theorem [Sam66, Théorème 2]: even though the set of nonconstant rational maps $V_1 \dashrightarrow C$ can now be infinite (because of Frobenius morphisms), they still do not vary in algebraic families.

LEMMA A.5. *Let R be a discrete valuation ring, with fraction field K and residue field k . Let X be a smooth projective geometrically integral curve of genus $g > 1$ over K with good reduction. Let $\text{Bir } X$ be the group of birational automorphisms of X , which is the same as the group of automorphisms. Let $\text{Bir } X_k$ be the corresponding group for the reduction X_k . Then the natural homomorphism $\text{Bir } X \rightarrow \text{Bir } X_k$ is injective.*

PROOF. Suppose that $f \in \text{Bir } X$ reduces to the identity 1_{X_k} . Let Γ be the graph of f , and let Δ be the graph of the identity 1_X , so Δ is the diagonal in $X \times X$. Then Γ and Δ both reduce to the diagonal $\Delta_k \subseteq X_k \times X_k$. Thus the intersection number $\Gamma \cdot \Delta$ equals the self-intersection number $\Delta_k \cdot \Delta_k$, which equals $2 - 2g$ [Har77, Exercise V.1.6(a)], which is negative. On the other hand, if the irreducible curves Γ and Δ were distinct, then $\Gamma \cdot \Delta \geq 0$. Hence $\Gamma = \Delta$, so $f = 1_X$. $\dashv$

For a different proof of Lemma A.5, see [Gre13, Theorem 2.1(a)] and the dissertation of Knaf cited there.

REFERENCES.

- [AK00] C.J. Ash and J.F. Knight, *Computable Structures and the Hyperarithmetical Hierarchy*, Elsevier, Amsterdam, 2000.
- [DJ94] R.G. Downey and C.G. Jockusch Jr., *Every low Boolean algebra is isomorphic to a recursive one*, Proceedings of the American Mathematical Society **122** (1994), 871–880.
- [DKL⁺15] R.G. Downey, A.M. Kach, S. Lempp, A.E.M. Lewis-Pye, A. Montalbán, and D.D. Turetsky, *The complexity of computable categoricity*, Advances in Mathematics **268** (2015), 423–466.
- [FKM10] Ekaterina Fokina, Iskander Kalimullin, and Russell Miller, *Degrees of categoricity of computable structures*, Archive for Mathematical Logic **49** (2010), 51–67.
- [FK82] E. Fried and J. Kollár, *Automorphism groups of fields*, Universal algebra (Esztergom, 1977), Colloq. Math. Soc. János Bolyai, vol. 29, North-Holland, Amsterdam-New York, 1982, pp. 293–303. MR660867
- [Gon77] S.S. Goncharov, *The quantity of nonautoequivalent constructivizations*, Algebra and Logic **16** (1977), 169–185 (English translation).
- [Gon80] S.S. Goncharov, *The problem of the number of nonautoequivalent constructivizations*, Algebra and Logic **19** (1980), 401–414 (English translation).
- [Gon81] S.S. Goncharov, *Groups with a finite number of constructivizations*, Soviet Math. Dokl. **23** (1981), 58–61.
- [GD80] S.S. Goncharov and V.D. Dzgoev, *Autostability of models*, Algebra and Logic **19** (1980), 45–58 (Russian), 28–37 (English translation).
- [Gre13] Barry Green, *Bounds on the number of automorphisms of curves over algebraically closed fields*, Israel J. Math. **194** (2013), no. 1, 69–76, DOI 10.1007/s11856-012-0078-0. MR3047062
- [HM07] Valentina Harizanov and Russell Miller, *Spectra of structures and relations*, The Journal of Symbolic Logic **72** (2007), 324–348.
- [HMM10] Valentina Harizanov, Russell Miller, and Andrei Morozov, *Simple structures with complex symmetry*, Algebra and Logic **49** (2010), 68–90.

- [HTMMM17] Matthew Harrison-Trainer, Alexander Melnikov, Russell Miller, and Antonio Montalbán, *Computable functors and effective interpretability*, Journal of Symbolic Logic **82** (2017), no. 1, 77–97. Preprint, [arXiv:1506.02737v1](#).
- [Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52. MR0463157 (57 #3116)
- [HKSS02] Denis R. Hirschfeldt, Bakhadyr Khoussainov, Richard A. Shore, and Arkadii M. Slinko, *Degree spectra and computable dimensions in algebraic structures*, Annals of Pure and Applied Logic **115** (2002), 71–113.
- [HKMS15] Denis R. Hirschfeldt, Ken Kramer, Russell Miller, and Alexandra Shlapentokh, *Categoricity properties for computable algebraic fields*, Transactions of the American Mathematical Society **367** (2015), 3955–3980.
- [JS91] Carl G. Jockusch and Robert Soare, *Degrees of orderings not isomorphic to recursive linear orderings*, Annals of Pure and Applied Logic **52** (1991), 39–64.
- [Kni86] Julia F. Knight, *Degrees coded in jumps of orderings*, J. Symb. Log. **51** (1986), 1034–1042.
- [KMVB07] Julia F. Knight, Sara Miller, and Michael Vanden Boom, *Turing computable embeddings*, J. Symb. Log. **72** (2007), no. 3, 901–918, DOI 10.2178/jsl/1191333847.
- [Lan60] Serge Lang, *Integral points on curves*, Inst. Hautes Études Sci. Publ. Math. **6** (1960), 27–43. MR0130219 (24 #A86)
- [LMMS05] Steffen Lempp, Charles McCoy, Russell Miller, and Reed Solomon, *Computable categoricity of trees of finite height*, Journal of Symbolic Logic **70** (2005), 151–215.
- [Lev09] Oscar Levin, *Computability Theory, Reverse Mathematics and Ordered Fields*, 2009. Ph.D. thesis, University of Connecticut.
- [Mar89] David Marker, *Non- Σ_n -axiomatizable almost strongly minimal theories*, Journal of Symbolic Logic **54** (1989), 921–927.
- [Mil09] Russell Miller, *$\mathbf{d}$ -Computable categoricity for algebraic fields*, Journal of Symbolic Logic **74** (2009), 1325–1351.
- [MS13] Russell Miller and Hans Schoutens, *Computably categorical fields via Fermat’s Last Theorem*, Computability **2** (2013), 51–65.
- [Mon14] Antonio Montalbán, *Computability theoretic classifications for classes of structures*, Proceedings of the ICM 2014 **2** (2014), 79–101.
- [Poo05] Bjorn Poonen, *Varieties without extra automorphisms. III. Hypersurfaces*, Finite Fields Appl. **11** (2005), no. 2, 230–268, DOI 10.1016/j.ffa.2004.12.001. MR2129679 (2006e:14060)
- [PER89] Marian B. Pour-El and J. Ian Richards, *Computability in analysis and physics*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1989. MR1005942 (90k:03062)
- [Prö84] Péter Pröhle, *Does the Frobenius endomorphism always generate a direct summand in the endomorphism monoids of fields of prime characteristic?*, Bull. Austral. Math. Soc. **30** (1984), no. 3, 335–356, DOI 10.1017/S0004972700002070. MR766793
- [Rem81] J.B. Remmel, *Recursively categorical linear orderings*, Proceedings of the American Mathematical Society **83** (1981), 387–391.
- [Ric81] Linda Jean Richter, *Degrees of structures*, Journal of Symbolic Logic **46** (1981), 723–731.
- [Sam66] Pierre Samuel, *Compléments à un article de Hans Grauert sur la conjecture de Mordell*, Inst. Hautes Études Sci. Publ. Math. **29** (1966), 55–62 (French). MR0204430 (34 #4272)

DEPARTMENT OF MATHEMATICS

QUEENS COLLEGE

65-30 KISSENA BLVD.

QUEENS, NY 11367, USA

AND

PH.D. PROGRAMS IN MATHEMATICS AND COMPUTER SCIENCE

CUNY GRADUATE CENTER

365 FIFTH AVENUE

NEW YORK, NY 10016, USA

E-mail: Russell.Miller@qc.cuny.edu

URL: <http://qcpages.qc.cuny.edu/~rmiller>

DEPARTMENT OF MATHEMATICS

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

CAMBRIDGE, MA 02139-4307, USA

E-mail: poonen@math.mit.edu

URL: <http://math.mit.edu/~poonen>

DEPARTMENT OF MATHEMATICS

NEW YORK CITY COLLEGE OF TECHNOLOGY

300 JAY STREET

BROOKLYN, NY 11201, USA

AND

PH.D. PROGRAM IN MATHEMATICS

CUNY GRADUATE CENTER

365 FIFTH AVENUE

NEW YORK, NY 10016, USA

E-mail: hschoutens@citytech.cuny.edu

DEPARTMENT OF MATHEMATICS

EAST CAROLINA UNIVERSITY

GREENVILLE, NC 27858, USA

E-mail: shlapentokha@ecu.edu

URL: <http://myweb.ecu.edu/shlapentokha/>