

A0142
A0165
A1813
A0085
A0898

$$S_m = 1 + \binom{m}{2} + \frac{1}{2!} \binom{m}{2} \binom{m-2}{2} + \frac{1}{3!} \binom{m}{2} \binom{m-2}{2} \binom{m-4}{2} + \dots$$

It will be important later to know how many of the $n!$ rook arrangements are symmetric about both diagonals S and S' . Denote this number by $(S'')_n$ for the $n \times n$ board. Then

$$(S'')_{2m+1} = (S'')_{2m},$$

and

$$(S'')_{2m} = 2(S'')_{2m-2} + (2m-2)(S'')_{2m-4}.$$

(To see this, note that the rook in the first column may or may not occupy one of the two corner squares in that first column. Consider both cases.) See Table 1.

$$2^n n! \quad \text{Table 1} \quad A_n = A_{n+1} \quad 2A_{n-1} \quad \rightarrow (A0899)$$

n	$n!$	$(R^2)_n$	R_n	S_n	$(S'')_n$	T_n
1	1	1		1	898	1
2	2	2	1813	85	2	1
3	6	2		4	2	2
4	24	8	2	10	6	7
5	120	8	2	26	6	23
6	720	48		76	20	115
7	5,040	48		232	20	694
8	40,320	384	12	764	76	5,282
9	362,880	384	12	2,620	76	46,066
10	3,628,800	3,840		9,496	312	456,454
11	39,916,800	3,840		35,696	312	4,999,004
12	479,001,600	46,080	120	140,152	1,384	59,916,028
13	6,227,020,800	46,080	120	568,504	1,384	778,525,516
14	87,178,291,200	645,120		2,390,480	6,512	10,897,964,660
15	1,307,674,368,000	645,120		10,349,536	6,512	163,461,964,024

~~M1624 M1824 M1986 M1185 M1599 M1708~~

In view of previous difficulties with this problem, it is interesting to look back to see why previous approaches did not work out so well. Kraitichik [2] and Madachy [4] indicate that the basic strategy heretofore was to divide the *essentially different solutions* into five mutually disjoint classes depending upon which *subgroups* of the group of symmetries of the square left them invariant. These classes were defined by:

- A : Those invariant only under I ,
- B : Those invariant under R , (R^2 and R^3),
- C : Those invariant under R^2 , but not invariant under any other rotation or reflection,
- D : Those invariant under S , S' , and R^2 , and
- E : Those invariant under S or S' , but not both.

If we let A_n , B_n , C_n , D_n , and E_n denote the number of elements in these sets, respectively, for the $n \times n$ case, we can see that

$$8A_n + 4C_n + 4E_n + 2B_n + 2D_n = n!$$

and

$$4C_n + 2B_n + 2D_n = 2^k k!, \text{ where } n = 2k \text{ or } 2k + 1$$

It was hoped that by playing with identities like this, one could simplify the problem. But such was not the case.

However, it is possible to work backward to obtain these numbers. It is easy to see that the following identities must hold:

$$E_n = [S_n - (S'')_n]/2; D_n = (S'')_n/2;$$

$$C_n = [(R^2)_n - R_n - (S'')_n]/4; B_n = R_n/2;$$

$$A_n = T_n - (B_n + C_n + D_n + E_n).$$

Notice in Table 2 that as n becomes large, the ratio of A_n to T_n approaches one; that is to say, only a tiny proportion of the arrangements will possess symmetry. This, of course, is to be expected. Thus for large n , $T_n \approx n!/8$.

A0902

Table 2

n	E_n	D_n	C_n	B_n	A_n	$\approx A_n/T_n$
1	A0900	✓	A0901	407	899	
2	✓	1	✓	1	1	.142857
3	1	1	7	1	9	.391304
4	2	3	7	6	4,820	.608696
5	10	3	74	6	44,676	.822767
6	28	10	74	60	59,834,748	.912533
7	106	10	11,144	60	778,230,060	.969826
8	344	38	159,652		10,896,609,768	.987666
9	1,272	38	159,652		163,456,629,604	.996253
10	4,592	156				.998643
11	17,692	156				.999620
12	69,384	692				.999876
13	283,560	692				.999967
14	1,191,984	3,256				
15	5,171,512	3,256				

M 1910

M 2772

M 4335

M 4160

M 4531

Finally, consider the corresponding placement problem for bishops (note that giving the board a 45° turn transforms this problem into one of rooks on a diamond-shaped board). It is easily shown that a maximum of $2n - 2$ non-attacking bishops can be placed on an $n \times n$ board and the total number of ways this can be done is 2^n . To find how many of these are essentially different we may again apply the Burnside Theorem. For this problem (see Reference 4, pp. 43-46, for more detail),

$$F(I) = 2^n,$$

$$F(S) = F(S') = F(R) = F(R^2) = F(R^3) = 0,$$