


RESEARCH ARTICLE

Plasma trimethylamine N-oxide and its metabolic precursors and risk of mortality, cardiovascular and renal disease in individuals with type 2-diabetes and albuminuria

Signe Abitz Winther^{1,2*}, Jens Christian Øllgaard², Tine Willum Hansen¹, Bernt Johan von Scholten², Henrik Reinhard¹, Tarunveer Singh Ahluwalia¹, Zeneng Wang³, Peter Gæde^{4,5}, Hans-Henrik Parving⁶, Stanley Hazen³, Oluf Pedersen⁷, Peter Rossing^{1,8}

1 Steno Diabetes Center, Copenhagen, Denmark, **2** Novo Nordisk A/S, Bagsvaerd, Denmark, **3** Department of Cardiovascular and Metabolic Sciences, Lerner Research Institute, Cleveland Clinic, Cleveland, OH, United States of America, **4** Slagelse Hospital, Slagelse, Denmark, **5** University of Southern Denmark, Odense, Denmark, **6** Rigshospitalet, Copenhagen, Denmark, **7** Novo Nordisk Foundation Center for Basic Metabolic Research, Copenhagen, Denmark, **8** University of Copenhagen, Copenhagen, Denmark

* signe.abitz.winther@regionh.dk



OPEN ACCESS

Citation: Winther SA, Øllgaard JC, Hansen TW, von Scholten BJ, Reinhard H, Ahluwalia TS, et al. (2021) Plasma trimethylamine N-oxide and its metabolic precursors and risk of mortality, cardiovascular and renal disease in individuals with type 2-diabetes and albuminuria. PLoS ONE 16(3): e0244402. <https://doi.org/10.1371/journal.pone.0244402>

Editor: M. Faadiel Essop, Stellenbosch University, SOUTH AFRICA

Received: July 20, 2020

Accepted: December 8, 2020

Published: March 3, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0244402>

Copyright: © 2021 Winther et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: As data contains potentially identifying patient information, access to

Abstract

Aims

The trimethylamine N-oxide (TMAO) pathway is related to intestinal microbiota and has been associated to risk of cardiovascular disease (CVD). We investigated associations between four plasma metabolites in the TMAO pathway and risk of all-cause mortality, CVD and deterioration in renal function in individuals with type 2-diabetes (T2D) and albuminuria.

Materials and methods

Plasma concentrations of TMAO, choline, carnitine, and betaine were measured by liquid chromatography-tandem mass spectrometry at baseline in 311 individuals with T2D and albuminuria. Information on all-cause mortality and fatal/non-fatal CVD during follow-up was obtained from registries. The association of each metabolite, and a weighted sum score of all four metabolites, with the endpoints were examined. Serum creatinine was measured at follow-up visits and the renal endpoint was defined as eGFR-decline of $\geq 30\%$. Associations were analysed using proportional hazards models adjusted for traditional risk factors.

Results

Baseline mean(SD) age was 57.2(8.2) years and 75% were males. Follow-up was up to 21.9 years (median (IQR) follow-up 6.8 (6.1–15.5) years for mortality and 6.5 (5.5–8.1) years for CVD events). The individual metabolites and the weighted sum score were not associated with all-cause mortality ($n = 106$) or CVD ($n = 116$) (adjusted $p \geq 0.09$). Higher choline, carnitine and the weighted sum score of the four metabolites were associated with

data requires approved protocol from the the Capital Region Committee on Health Research Ethics (vek-anmeldelse.center-for-regional-udvikling@regionh.dk) and approved data sharing agreement with the Capital Region (cru-fp-vid@regionh.dk). Thereafter data can be provided from Steno Diabetes Center Copenhagen (stenoinfo.stenodiabetescentercopenhagen@regionh.dk, attention "Complications Research").

Funding: This work was funded by the Novo Nordisk Foundation, grant no. NNF 14OC0013659 as part of the PROTON (Personalizing Treatment of diabetic Nephropathy) study. The Novo Nordisk Foundation Center for Basic Metabolomics Research is an independent research center at the University of Copenhagen partially funded by an unrestricted donation from the Novo Nordisk Foundation (www.cbmr.ku.dk) (grant number NNF18CC0034900). SLH reports funding by NIH grant P01 HL147823 and the Leducq Foundation. SAW was funded by The Innovation Fund Denmark (grant number 5016-00150B) and Novo Nordisk A/S provided support in the form of salary during the PhD fellowship for author SAW. JO and BJvS are after data collection employed by Novo Nordisk A/S. The specific roles of these authors are articulated in the 'author contributions' section. The funding sources did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: I have read the journal's policy and the authors of this manuscript have the following competing interests: PR reports having given lectures for Astra Zeneca, Bayer, Novo Nordisk and Boehringer Ingelheim, and has served as a consultant for AbbVie, Astra Zeneca, Bayer, Eli Lilly, Boehringer Ingelheim, Astellas, Gilead, Mundipharma, Vifor, and Novo Nordisk, all fees given to Steno Diabetes Center Copenhagen. SAW, TWH, BJvS, TSA and PR own stocks in Novo Nordisk A/S and TSA in Zealand Pharma A/S. Z.W. and S.L.H. are named as coinventors on pending and issued patents held by the Cleveland Clinic relating to cardiovascular diagnostics and therapeutics, and have the right to receive royalty payment for inventions or discoveries related to cardiovascular diagnostics or therapeutics from Cleveland Heart Lab, Quest Diagnostics and Proctor & Gamble. S.L.H. also reports having been paid as a consultant from Proctor & Gamble, and having received research funds from Proctor & Gamble and Roche. Novo Nordisk A/S provided support in the form of salaries for authors SAW, JO and BJvS. This does not alter our adherence to

higher risk of decline in eGFR ($n = 106$) (adjusted $p = 0.001$, $p = 0.03$ and $p < 0.001$, respectively).

Conclusions

In individuals with T2D and albuminuria, higher choline, carnitine and a weighted sum of four metabolites from the TMAO pathway were risk markers for deterioration in renal function during long-term follow-up. Metabolites from the TMAO pathway were not independently related to risk of all-cause mortality or CVD.

Introduction

The prevalence of type 2-diabetes (T2D) is increasing world-wide, accompanied by a cumulative burden of micro- and macrovascular complications. Even though prevention and treatment of cardiovascular disease (CVD) and renal complications of T2D have improved, the incidence is still high and the outcome often remains lethal. Moreover, an improved survival with increasing incidence of diabetes is resulting in increasing prevalence of renal complications [1].

Metabolites produced by the intestinal microbiota through digestion of certain dietary products are suggested to be involved in the development of metabolic diseases and risk of CVD, progression of kidney disease and mortality [2, 3]. Recently, there has been particular focus on the trimethylamine N-oxide (TMAO) pathway [4, 5]. A fraction of bacteria in the gut digest choline and L-carnitine from diet sources such as red meat, eggs and cheese and produce trimethylamine (TMA). In the liver TMA is oxidized to TMAO by flavin monooxygenases, and thereafter circulated to various tissues or excreted through the kidneys.

Previous studies have linked circulating concentrations of TMAO and its metabolic precursors (choline, L-carnitine and betaine) to mortality and adverse outcomes of CVD including cohorts of subjects with T2D or pre-existing CVD [6–8]. Plasma TMAO concentrations have also been associated with development of renal insufficiency both in subjects with normal kidney function and in chronic kidney disease [3, 9]. TMAO modulates lipid metabolism and increases vascular cell activation and endogenous inflammation thereby promoting atherogenesis [2, 10, 11]. However, it has been suggested that the adverse effect of TMAO may be mediated by impaired renal function or other mediators [9, 12] and it is still discussed whether the precursors of TMAO are associated with risk of mortality, CVD and renal outcomes independently of TMAO [13–15]. Hence, the role of TMAO and its precursors as risk markers for mortality, CVD and renal outcomes need further clarification. Therefore, in the present follow-up study we investigated associations between TMAO and its metabolic precursors (choline, L-carnitine and betaine) and risk of mortality, CVD and renal function decline in individuals with T2D and microalbuminuria.

Material and methods

Study subjects

We combined two Danish cohorts originating from the outpatient clinic at Steno Diabetes Center Copenhagen. Both cohorts included individuals with T2D and albuminuria. Design, method and follow-up for the original studies have been described in detail previously [16–19]. Briefly, the first cohort included individuals from the Steno-2 trial (ClinicalTrials.gov

PLOS ONE policies on sharing data and materials.
The other authors report no conflicts of interest.

registration no. NCT00320008). This was a randomized controlled trial of 160 T2D individuals with microalbuminuria ($n = 138$ with available metabolite measures for the present study) receiving conventional versus intensified multifactorial treatment [16]. Inclusion criteria included individuals with T2D (defined according to the 1985 WHO criteria), age 40–65 years and persistent microalbuminuria. Microalbuminuria was defined as albumin excretion rate (UAER) of 30–300 mg in four of six 24-h urine samples. The trial was initiated in 1993 with end of intervention after a mean of 7.8 years, and thereafter continued as an observational study. The second cohort comprised 200 T2D individuals with persistent albuminuria (two out of three consecutive measured UAER > 30 mg/24h). Other inclusion criteria included age between 20 and 70 years, no known coronary artery disease and normal plasma creatinine. Of the 200 participants included in the original observational study, 173 had available metabolite measures for the present study. These participants were examined in 2007–08 with the original purpose to investigate presence of asymptomatic coronary artery disease in T2D individuals with albuminuria [18]. All participants received intensive multifactorial treatment constituting of glycaemic, lipid and blood pressure control, as well as antithrombotic therapy and lifestyle modification. Individuals from the original two studies with available metabolite measures from stored blood samples were included in the present study. Fifteen individuals participated in both studies and are in the present analyses included with baseline measures of the metabolites and clinical variables from the participation in the Steno-2 trial. The present study complies with the Declaration of Helsinki and was approved by the local ethics committee (the Capital Region Committee on Health Research Ethics, approval number H-17004111). All individuals gave written informed content at the original recruitment.

Measurements of biochemical variables

Plasma concentrations of TMAO, choline, carnitine and betaine were measured in samples collected at baseline. All samples were centrifuged, immediately stored in freezers at -80°C until analysis and subsequently they were analysed at the same time. Plasma concentrations of TMAO, choline, carnitine and betaine were measured using stable isotope dilution high-performance liquid-chromatography with online electrospray ionization-tandem mass spectrometry (LC/MS/MS) on an AB-SCIEX5000 triple quadrupole mass spectrometer using d9-(trimethyl)-labelled internal standards as described previously [20] and performed in the Lerner Research Institute, Cleveland Clinic, by investigators blinded to any participants' characteristics. Quantification of plasma concentrations of TMAO, choline, carnitine and betaine each had a limit of quantification of 20 nmol/L. Quality control samples were run with each batch, and the inter-day coefficient-of-variation for each analyte was less than 10% over a span of 20 days.

UAER was measured in 24-h urine collections by an enzyme immunoassay. Until September 1st 2004, serum creatinine levels were measured by a modified Jaffe method and thereafter by an isotope dilution-mass spectrometry method. Therefore, all creatinine measurements determined by Jaffe were corrected to standardize measurements [21]. Estimated glomerular filtration rate (eGFR) was calculated using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation [22]. Serum cholesterol concentrations were determined by standard methods. HbA1c was measured by high-performance liquid-chromatography and current smoking was defined as one or more cigarettes/cigars/pipes a day.

Endpoints

Through the Danish nationwide registers, all participants were tracked by their unique and personal identification number to obtain information on all hospital admissions and cause of

death. Reporting to these registries is mandatory and has high validity [23]. One participant was lost to follow-up after 13 years due to emigration. Extracted diagnoses for the two cohorts have previously been specified [17, 19]. We considered four endpoints; All-cause mortality, CV mortality, CVD and eGFR decline $\geq 30\%$. All deaths were classified as CVD related unless an unequivocal non-CV cause was reported, a previously recognized approach [24]. CVD events were defined as a composite of cardiovascular mortality, non-fatal myocardial infarction, ischaemic CVD, non-fatal stroke, amputation due to ischemia and cardiac or peripheral revascularization. Individuals experiencing multiple events stopped their follow-up at time of the first event in the analyses.

The renal endpoint was defined as time to the first occurrence of $\geq 30\%$ decrease in eGFR from baseline, an approach previously proposed [25]. Measurements of renal function (serum creatinine) were obtained from follow-up visits after approximately 2, 4, 8, 13 and 21 years in the Steno-2 trial [26] and in the other observational cohort at follow-up visits after approximately 2, 3, 4 and 5 years. Furthermore, we considered additional renal endpoints and applied linear regression analysis to calculate rate of eGFR decline (eGFR slope) based on measurements from baseline, in individuals with at least two measurements and a minimum of follow-up of 3 years ($n = 273$). Participants from the Steno-2 trial ($n = 138$) had ^{51}Cr -ethylenediamine tetra-acetic acid (^{51}Cr -EDTA) clearance glomerular filtration rate (mGFR) measured by standard methods [27] at baseline and at follow-up visits. For the subset of individuals with mGFR, a subanalysis was performed for the endpoint of $\geq 30\%$ decline in mGFR. In addition, an endpoint of $\geq 40\%$ decline in eGFR from baseline was computed for a sensitivity analysis.

Statistics

Baseline data are given as mean \pm standard derivation (SD) for normal distributed continuous variables and as median and interquartile range (IQR) for non-normal distributed. Categorical variables are reported as numbers (%). The non-normal distributed variables, including the four metabolites, were \log_2 -transformed before correlation and regression analyses. We applied analyses of variance for normal distributed continuous variables and the Kruskal-Wallis analysis of variance for the non-normal distributed continuous variables; and the chi-square test for categorical variables to compare differences between quartiles of TMAO.

Correlations between the \log_2 -transformed four metabolites at baseline were examined using Pearson correlation and reported as the coefficient of determination (R^2).

Survival analyses with proportional hazard were performed to calculate hazard ratios (HRs) with 95% confidence intervals (CIs) per 1 SD increase in \log_2 -transformed concentrations of the metabolites for all endpoints. We considered all-cause mortality as a competing event for CVD and eGFR decline $\geq 30\%$, and non-CV mortality as a competing event for CV mortality. No other competing events were considered. We applied cause-specific hazards models when competing events were present. First, we performed unadjusted analyses. Next, we adjusted for baseline age, sex, HbA_{1c}, systolic blood pressure, body mass index, total cholesterol, smoking, UAER and eGFR by adding each variable solely and, thereafter, stepwise to the model. In addition to the analysis of the four metabolites separately, we calculated a weighted metabolite sum score as the weighted sum of concentrations of the four metabolites—a previously used method applied to reduce variability [28]. The weight for each metabolite was estimated as the parameter estimates for a 1 SD increase in the plasma concentrations of the metabolites, calculated from the proportional hazards model including all four metabolites and covariates. The weighted sum score was thereafter included as an exposure variable in the survival analyses. Moreover, we tested for heterogeneity in the HRs for the influence of age and sex by

introduction of the appropriate interaction terms in the proportional hazards model. We tested the proportional hazards model assumption for linearity of the logarithm of the metabolites for the endpoints by plotting parameter estimates of quintiles versus means of each quintile. Moreover, the assumption for proportional hazard was tested and fulfilled for all endpoints.

A two-tailed p-value < 0.05 was considered significant. Statistical analyses were performed using SAS 9.4 software (SAS Institute, Cary, NC) and R (version 3.5.2; <http://www.r-project.org/>).

Results

Characteristics at baseline

Plasma TMAO and related metabolite concentrations were available from 311 individuals. Baseline clinical characteristics of the participants are summarized by quartiles of TMAO in [Table 1](#). Overall, the mean \pm SD age was 57.2 \pm 8.2 years, known diabetes duration was median

Table 1. Clinical characteristics at baseline.

	Overall	Trimethylamine N-oxide				p
		Quartile 1 $\leq 3.8 \mu\text{M}$	Quartile 2 $> 3.8 \text{ to } \leq 5.9 \mu\text{M}$	Quartile 3 $> 5.9 \text{ to } \leq 9.0 \mu\text{M}$	Quartile 4 $> 9.0 \mu\text{M}$	
Number of participants	311	78	78	79	76	
From the Steno-2 trial, n (%)	138	48 (61.5)	42 (53.8)	28 (35.4)	20 (26.3)	0.003
Age (years)	57.2 \pm 8.2	54.3 \pm 8.1	56.6 \pm 8.0	58.2 \pm 8.2	59.8 \pm 7.7	<0.001
Female, n (%)	77 (24.8)	20 (25.6)	23 (29.5)	15 (19.0)	19 (25.0)	0.50
Known diabetes duration (years)	8 (4–14)	5 (3–10)	8 (4–13)	9 (5–14)	10 (6–17)	0.001
Systolic blood pressure (mmHg)	138 \pm 20	141 \pm 18	140 \pm 22	133 \pm 21	138 \pm 20	0.038
Diastolic blood pressure (mmHg)	79 \pm 12	83 \pm 12	80 \pm 13	76 \pm 13	77 \pm 10	0.002
HbA _{1c} (mmol/mol)	64 \pm 18.2	64 \pm 18.8	65 \pm 18.6	66 \pm 19.0	62 \pm 16.2	0.44
HbA _{1c} (%)	8.0 \pm 3.8	8.0 \pm 3.9	8.1 \pm 3.9	8.2 \pm 3.9	7.8 \pm 3.6	0.44
Body mass index (kg/m ²)	31.4 \pm 5.5	30.6 \pm 4.9	31.8 \pm 5.7	31.9 \pm 5.9	31.4 \pm 5.5	0.45
eGFR (ml/min/1.73m ²)	93 \pm 16.3	100 \pm 14.8	96 \pm 14.3	91 \pm 17.0	86 \pm 17.4	<0.001
UAER (mg/d)	86 (51–180)	90 (55–190)	88 (54–177)	90 (51–156)	74 (37–173)	0.33
Serum LDL cholesterol (mM)	2.5 \pm 1.2	2.8 \pm 1.2	2.8 \pm 1.2	2.4 \pm 1.2	2.2 \pm 1.1	0.006
Serum total cholesterol (mM)	4.7 \pm 1.4	4.9 \pm 1.3	4.9 \pm 1.5	4.6 \pm 1.4	4.4 \pm 1.4	0.10
Current smoker, n (%)	103 (33)	32 (41)	25 (32)	23 (29)	23 (30)	0.38
Plasma choline (μM)	15.2 (12.4–19.2)	12.7 (10.6–16.8)	14.7 (12.5–18.2)	16.3 (13.1–18.6)	17.8 (14.0–21.7)	<0.001
Plasma carnitine (μM)	39.1 (31.0–49.9)	35.3 (28.2–42.6)	38.6 (30–48.2)	39.8 (30.2–49.0)	45.4 (35.9–57.0)	<0.001
Plasma betaine (μM)	35.0 (22.3–53.4)	26.9 (18.7–43.2)	31.8 (22.3–47.6)	36.2 (25.0–58.9)	41.5 (28.7–62.6)	<0.001
Treatment with:						
Oral antidiabetics, n (%)	256 (83)	60 (77)	61 (78)	68 (86)	67 (88)	0.17
Insulin, n (%)	118 (38)	18 (23)	28 (36)	33 (42)	39 (51)	0.003
RAAS blocking agents, n (%)	197 (63)	39 (50)	44 (56)	57 (72)	57 (75)	0.002
Acetylsalicylic acid, n (%)	178 (57)	32 (41)	45 (58)	46 (58)	55 (72)	0.001
Lipid lowering agents, n (%)	165 (53)	30 (39)	34 (44)	47 (60)	54 (71)	<0.001

Data are shown as numbers (%), mean \pm SD and median (IQR). eGFR: estimated glomerular filtrations rate; UAER: urinary albumin excretion rate; RAAS: renin-angiotensin-aldosterone system. We applied analyses of variance for normal distributed continuous variables and the Kruskal-Wallis analysis of variance for the non-normal distributed continuous variables; and the chi-square test for categorical variables to compare differences between quartiles of TMAO. The weighted metabolite sum score was calculated as the weighted sum of concentrations of the four metabolites.

<https://doi.org/10.1371/journal.pone.0244402.t001>

(IQR) 8.0 (4.0–14.0) years and 24.8% were females. The plasma concentrations of TMAO were 5.9 (3.8–9.0) μM , choline 15.2 (12.4–19.2) μM , carnitine 39.1 (31.0–49.9) μM and betaine 35.0 (22.3–53.4) μM , respectively. Individuals with higher TMAO levels were older, had longer diabetes duration, lower eGFR, and lower LDL and total cholesterol levels than subjects with lower TMAO concentrations. In addition, at baseline individuals with higher TMAO were more frequently treated with insulin, renin-angiotensin-aldosterone-system blocking agents, acetylsalicylic acid and cholesterol lowering agents.

Baseline concentrations of all four plasma compounds related to the TMAO pathway were correlated, and the strongest positive correlation was demonstrated between choline and carnitine; and between choline and betaine ($R^2 = 0.33$, $p < 0.0001$ for both) (S1 Fig). TMAO was weakly positively correlated with the other metabolites (TMAO and choline: $R^2 = 0.10$; TMAO and carnitine: $R^2 = 0.07$; TMAO and betaine: $R^2 = 0.08$, $p < 0.001$ for all).

Mortality and CVD events

In the total population, follow-up was up to 21.9 years. Median (IQR) follow-up in years was 6.8 (6.1–15.5) for mortality and 6.5 (5.5–8.1) for CVD events. During follow-up a total of 106 persons died. Of 116 persons that suffered CVD events, 44 of these events were fatal.

The unadjusted and adjusted HRs with 95% CI for all endpoints are presented in Tables 2 and 3 and visualised in Fig 1. In our study cohort, plasma TMAO was not associated with any of the endpoints. Higher plasma concentrations of choline and the weighted sum score of metabolites were in unadjusted analyses associated with all-cause mortality (HR [95% CI] 1.26 [1.01–1.57] and 1.09 [1.02–1.16], respectively). However, these associations were lost after adjustment. A higher weighted sum score was associated with cardiovascular mortality both before and after adjustment (adjusted HR 1.14 [1.00–1.29]). None of the metabolites were associated with risk of CVD.

Progression of kidney disease

For the renal endpoint of an eGFR decline $\geq 30\%$, 289 subjects were available for analyses and 106 reached the endpoint over a median follow-up of 4.9 (3.7–8.0) years. Higher plasma

Table 2. Plasma concentrations of TMAO and related compounds in relation to all-cause mortality, cardiovascular mortality and the combined cardiovascular endpoint.

	All-cause mortality	p	CV mortality	p	CVD	p
Events, n (%)	106 (34.1)		44 (14.1)		116 (37.3)	
TMAO—unadjusted	1.06 (0.88–1.28)	0.53	1.09 (0.82–1.45)	0.54	1.14 (0.96–1.35)	0.14
TMAO—adjusted	1.02 (0.83–1.26)	0.85	0.98 (0.70–1.37)	0.92	1.11 (0.93–1.33)	0.26
Choline—unadjusted	1.26 (1.01–1.57)	0.04	1.33 (0.95–1.87)	0.10	0.98 (0.81–1.18)	0.81
Choline—adjusted	1.17 (0.93–1.48)	0.18	1.29 (0.89–1.86)	0.18	0.98 (0.79–1.21)	0.82
Carnitine—unadjusted	1.04 (0.85–1.28)	0.70	1.28 (0.92–1.78)	0.14	1.00 (0.83–1.21)	0.96
Carnitine—adjusted	1.02 (0.82–1.27)	0.86	1.26 (0.88–1.81)	0.21	1.04 (0.84–1.27)	0.75
Betaine—unadjusted	0.96 (0.77–1.19)	0.70	0.92 (0.66–1.28)	0.63	0.92 (0.75–1.12)	0.38
Betaine—adjusted	0.97 (0.75–1.24)	0.79	0.95 (0.63–1.41)	0.78	1.02 (0.81–1.29)	0.88
Weighted sum score—unadjusted	1.09 (1.02–1.16)	0.01	1.15 (1.03–1.29)	0.01	1.08 (0.98–1.20)	0.12
Weighted sum score—adjusted	1.06 (0.99–1.14)	0.09	1.14 (1.00–1.29)	0.04	1.08 (0.97–1.20)	0.17

Proportional hazards models presented as hazard ratios (HR) with 95% confidence intervals for all-cause mortality, CV mortality and CVD. Cause-specific hazards models were applied for CV mortality and CVD. HRs express risk per 1 standard deviation increase in the log2 transformed plasma concentrations of the metabolites. Weighted sum score was calculated as a weighted score of the four metabolites. Adjustment included age, sex, HbA_{1c}, systolic blood pressure, body mass index, total cholesterol, smoking, urinary albumin excretion rate and eGFR at baseline. CVD: cardiovascular disease; eGFR: estimated glomerular filtration rate.

<https://doi.org/10.1371/journal.pone.0244402.t002>

Table 3. Plasma concentrations of TMAO and related compounds in relation to a decline in eGFR $\geq 30\%$, a decline in eGFR $\geq 40\%$ and a decline in mGFR $\geq 30\%$.

	eGFR decline $\geq 30\%$	p	eGFR decline $\geq 40\%$	p	mGFR decline $\geq 30\%$	p
Numbers available for analyses	289		289		138	
Events, n (%)	106 (36.7)		78 (25.6)		89 (64.5)	
TMAO—unadjusted	1.14 (0.97–1.37)	0.12	1.04 (0.85–1.28)	0.69	1.07 (0.89–1.27)	0.47
TMAO—adjusted	1.08 (0.90–1.31)	0.42	1.00 (0.80–1.27)	0.96	1.11 (0.92–1.35)	0.28
Choline—unadjusted	1.58 (1.27–1.96)	<0.001	1.69 (1.30–2.20)	<0.001	1.38 (1.04–1.83)	0.03
Choline—adjusted	1.49 (1.18–1.89)	0.001	1.66 (1.25–2.22)	<0.001	1.49 (1.08–2.04)	0.01
Carnitine—unadjusted	1.20 (0.98–1.47)	0.08	1.42 (1.10–1.82)	0.006	1.08 (0.83–1.39)	0.58
Carnitine—adjusted	1.27 (1.02–1.59)	0.03	1.55 (1.17–2.07)	0.003	1.31 (0.99–1.72)	0.06
Betaine—unadjusted	1.10 (0.88–1.39)	0.41	1.16 (0.89–1.52)	0.28	0.96 (0.73–1.27)	0.79
Betaine—adjusted	1.07 (0.82–1.39)	0.63	1.14 (0.82–1.58)	0.43	1.07 (0.78–1.47)	0.69
Weighted sum score—unadjusted	1.40 (1.20–1.64)	<0.001	1.49 (1.23–1.81)	<0.001	1.28 (1.02–1.61)	0.04
Weighted sum score—adjusted	1.36 (1.16–1.60)	<0.001	1.54 (1.25–1.89)	<0.001	1.49 (1.13–1.97)	0.005

Cause-specific proportional hazards models presented as hazard ratios (HR) with 95% confidence intervals for decline in eGFR $\geq 30\%$, decline in eGFR $\geq 40\%$ and decline in mGFR $\geq 30\%$. HRs express risk per 1 standard deviation increase in the log₂ transformed concentrations of the metabolites. Weighted sum score was calculated as a weighted score of the four metabolites. Adjustment included baseline age, sex, HbA_{1c}, systolic blood pressure, body mass index, total cholesterol, smoking, urinary albumin excretion rate and eGFR/mGFR, as appropriate. eGFR: estimated glomerular filtration rate; mGFR: measured glomerular filtration rate.

<https://doi.org/10.1371/journal.pone.0244402.t003>

choline and the weighted sum score were risk markers for a decline in eGFR $\geq 30\%$, in both unadjusted and adjusted models. Higher plasma concentrations of carnitine were also a risk marker in the adjusted model (HR 1.27 [1.02–1.59]), but not unadjusted ($p = 0.08$) (Table 3).

We also evaluated the risk of an eGFR decline $\geq 40\%$ (78 events) presented in Table 3. Higher plasma concentrations of choline, carnitine and the weighted sum score were associated with higher risk of eGFR decline $\geq 40\%$ both unadjusted and after adjustment (adjusted HR choline: 1.66 [1.25–2.22], carnitine: 1.55 [1.17–2.07] and weighted sum score: 1.54 [1.25–1.89]).

In a regression model with yearly change in eGFR ($n = 273$) evaluated over a median of 5.0 (4.0–13.0) years, higher concentrations of choline were associated with a steeper yearly decline in eGFR in both unadjusted ($p = 0.001$) and adjusted ($p < 0.001$) models. Higher carnitine levels were also associated with a steeper yearly decline in eGFR in adjusted analyses ($p = 0.04$). Plasma TMAO and betaine were not associated with yearly change in eGFR ($p \geq 0.18$).

In the subset of participants from the Steno-2 trial with measurements of ⁵¹Cr-EDTA GFR (mGFR) at baseline and during follow-up ($n = 138$), we estimated the risk of mGFR decline $\geq 30\%$ (89 events). In these analyses, higher plasma concentrations of choline and the weighted sum score were risk markers both in unadjusted and adjusted models (adjusted HR 1.49 [1.08–2.06] and 1.49 [1.13–1.96], respectively), Table 3. TMAO and betaine were not associated with mGFR decline $\geq 30\%$.

Additional inclusion of the plasma concentrations of TMAO in the proportional hazards models for choline, carnitine and betaine did not change the overall results for any of the endpoints (mortality, CVD events or renal endpoints).

Additional analyses

In sensitivity analyses, we included additional adjustment for cohort and treatment with RAAS blocking agents, insulin, acetylsalicylic acid and lipid lowering agents. This did not change any of the results. To consider competing risk for the endpoint of CV mortality, a proportional hazard model was performed for non-CV mortality ($n = 62$); None of the




Fig 1. Forrest plot visualizing of plasma concentrations of TMAO and related compounds in relation to clinical endpoints. Proportional hazards models presented as hazard ratios (HRs) with 95% confidence intervals for all-cause mortality ($n = 106$), CV mortality ($n = 44$), CVD ($n = 116$) and eGFR decline $\geq 30\%$ ($n = 106$). HRs express risk per 1 standard deviation increase in the log2 transformed plasma concentrations of the metabolites. Weighted sum score was calculated as a weighted score of the four metabolites. Adjustment included age, sex, HbA_{1c}, systolic blood pressure, body mass index, serum total cholesterol, smoking, urinary albumin excretion rate and eGFR at baseline. Level of significance is denoted as follows: *: $0.01 \leq p < 0.05$, **: $0.001 \leq p < 0.01$, ***: $p < 0.001$. TMAO = trimethylamine N-oxide.

<https://doi.org/10.1371/journal.pone.0244402.g001>

metabolites or weighted metabolite sum score were associated with the risk of non-CV mortality ($p \geq 0.14$).

Discussion

We investigated whether plasma concentrations of TMAO and its metabolic precursors choline, carnitine and betaine as well as a composite score of metabolites, were associated with mortality, adverse CVD events and deterioration in renal function in individuals with T2D and albuminuria.

We found that higher plasma concentrations of choline and carnitine (carnitine, only in adjusted models) were directly associated with deterioration in renal function. Also, when we combined the four examined metabolites in the TMAO pathway into a weighted sum score, a higher score was positively associated with deterioration in renal function. Furthermore, the weighted sum score of metabolites was directly associated with cardiovascular mortality. We could not demonstrate any association between plasma concentrations of TMAO or its metabolic precursors and the risk of CVD.

The TMAO pathway is the first in the literature that links metabolites produced by the gut microbiota to the risk of cardiovascular and renal diseases especially in cohorts with established CVD [3, 4, 7]. Such associations may be of interest in the search for new therapeutic targets for prevention of CVD and progression of renal disease.

In our study we found an association between the weighted sum score and CV mortality, but not when the metabolites were analysed separately. This may indicate that calculating a weighted sum of the TMAO related metabolites is justified potentially by reducing variability in the measurements and day-to-day variability of individual metabolites. In a previous report a comparable sum score including five metabolites in the TMAO pathway (TMAO, choline, phosphocholine, α -glycerophosphocholine, and betaine) was calculated and shown to be linked to an increased risk of CVD events [28].

Previous studies in persons with T2D have demonstrated a positive association between higher plasma concentrations of TMAO and increased risk of CVD events and mortality even after adjustment for traditional CVD risk factors [6, 8, 29, 30]. In addition, in recent meta-analyses, higher concentrations of TMAO were reported to be a risk marker of mortality and CVD risk amongst all subjects [7]. In our cohort, we could not replicate such associations which may be explained by clinical differences between the reported cohorts and ours. Previous studies in T2D included individuals undergoing elective coronary angiography or with already existing CVD at baseline. In the present study, only a few of our included participants had known coronary artery disease at baseline as it was an exclusion criterion for the individuals enrolled in our study during 2007–08. Moreover, the reported studies had a lower median eGFR ranging between 62–82 (50–94) ml/min/1.73 m² [6, 30] compared to our participants with median eGFR 96 (82–104) ml/min/1.73 m². Still, the impact of renal function as a potential mediator or contributing factor of the adverse effect of TMAO is debated [12, 15, 31]. Though it should be noted that in a recent meta-analysis that examined the impact of renal function on TMAO associations with all-cause mortality and CVD, TMAO remained significantly associated with both all-cause mortality and CVD risks in individuals with and without CKD [7]. The level of measured plasma TMAO in our study did not differ substantially from previous studies (5.9 [3.8–9.0] μ M in our study compared with 4.4 [2.8–7.7] μ M and 7.5 [4.4–12.1] μ M, respectively in the other studies [6, 30]). Another limitation to consider is the lack of statistical power. However, due to the long duration of our study (up to 21 years), we have many person-years of follow-up and a relatively high overall event rate in the aging population monitored. Further, we have with previous markers that show strong association with renal function like plasma symmetric and asymmetric dimethylarginine and urinary kidney injury molecule 1 (u-KIM-1), shown an association between these and cardiovascular disease [19, 32, 33].

Consistent with our study, a recent paper also failed to show an association between TMAO and CVD in T2D [34]. This study by Cardona *et al.* was a post-hoc case-control analyses of the ACCORD study including persons with T2D without evident atherosclerotic disease at baseline. With similar plasma concentrations of TMAO and traditional CVD risk factors at baseline between controls and cases the investigators concluded that the lack of a link between plasma concentration of TMAO and CVD might be due to the absence in difference of

traditional CVD risk factors. Another study including individuals with high cardiovascular risk when estimated from conventional CVD risk factors, but free of CVD at baseline also failed to show an association with incident CVD, indicating that plasma TMAO concentration may be a better risk marker in established CVD [28]. In line with this, a very recent larger study of a T2D cohort found an association between higher plasma concentrations of TMAO and cardiovascular endpoints and mortality, however these associations were insignificant in subanalyses excluding participants with known CVD at baseline [8]. It is also of interest to note that a recent experimental and human study indicated that TMAO might be a better marker of instability of established atherosclerotic plaques, than the development of or extent of atherosclerosis [35].

It is worth noticing that in our cohort the individuals with the highest plasma concentrations of TMAO also were those with the highest risk profile of traditional CVD risk factors. Drugs intended to lower these risk factors were also more frequently prescribed in the individuals with highest plasma TMAO levels. This may have attenuated a potential association between TMAO and the outcomes. In this situation it would be difficult to conclude, whether plasma concentrations of TMAO (and its precursors) are a risk factor for CVD or just markers of a high-risk profile based on known risk factors.

The results of previous studies on the association between the other metabolites of the TMAO pathway and risk of mortality and CVD have been inconsistent [4, 9, 14, 28, 36]. In a large population-based study, higher plasma concentration of phosphatidylcholine was directly associated with risk of all-cause mortality, especially CVD related [36]. An association that was stronger among individuals with diabetes than in those without. In contrast, the association between plasma concentrations of choline and betaine and the risk of CVD may be confounded by the plasma level of TMAO as the main driver of the association [14]. In our study, the associations for choline were independent of the level of TMAO and a recent meta-analysis could not demonstrate any association between circulating concentrations of choline or betaine and risk of CVD [37].

A few studies have investigated the association between circulating choline and renal outcome in humans indicating a positive association between choline and an aggravation of renal function decline [9, 38]. In addition, experimental studies in a mouse model have demonstrated that TMAO and choline are associated with renal damage expressed as increased circulating concentrations of Cystatin C and KIM-1 [3].

Potential therapeutic targets of the TMAO pathway have recently been investigated in experimental murine models with promising results. In a mice model of chronic kidney disease, small molecule inhibition of microbial TMA generation (the precursor of TMAO) with a mechanism-based inhibitor iodomethylcholine (a TMA-lyase inhibitor predominately targeting TMA production in the gut lumen) reduced the decline in renal function and tubulointerstitial fibrosis [39]. A general inhibition of the same pathway also attenuates atherosclerosis and thrombosis in mice [40, 41]. The molecular mechanisms of TMAO's adverse effect on kidney impairment has been shown to involve TGF- β induced signalling pathways, and induction of multiple pro-fibrosis forming gene expression changes, and yet the receptor involved in this effect remains unknown and needs further exploration [39]. The composition of diet may also be a regulating factor of TMAO and the impact of CVD and renal impairment. A large observational study (n = 55,113) demonstrated that vegans (e.g. a low choline diet) had a lower prevalence of chronic kidney disease compared to omnivores [42]. Diet restriction may therefore also be considered as an intervention. However, randomised controlled clinical trials are needed to clarify any causal effects of these interventions.

Our study has limitations. The plasma metabolites were not measured in the fasting state and only measured once at baseline and hence day-to-day variability due to for example

expected variation in diet might have influenced our results. Another limitation is that the metabolites were measured in samples stored for a long period. Even though plasma concentrations of TMAO show stability when stored on -80°C over 5 years with multiple freeze thaw cycles [43], stability of the metabolites are not known for follow-up of up to 21 years. Medication use throughout follow-up period was not incorporated into any analyses presented. Moreover, it is possible that estimation of GFR using cystatin C would have increased accuracy, however we do not expect this would have affected our findings significantly [44]. Finally, the study population included 77% males and although we did not find any significant effect modification on sex, there could be variations in the TMAO pathway with sex, which have impacted our results. Strengths of our study include a well-characterized cohort and complete long-time follow-up based on data from Danish registries with high quality.

In conclusion, higher plasma concentrations of choline, carnitine and a weighted sum of the concentrations of four plasma metabolites related to the TMAO pathway were directly associated with a deterioration of renal function in Danish white individuals with T2D and albuminuria. Moreover, a higher weighted sum score of the TMAO pathway-related metabolites was positively associated with risk of CV mortality, whereas individual metabolites were not.

Supporting information

S1 Fig. Pearson correlations between the four log₂-transformed plasma metabolites. Numbers are R^2 and all p-values <0.0001 . TMAO = trimethylamine N-oxide. (TIFF)

Acknowledgments

We thank the lab technicians at Steno Diabetes Center Copenhagen for technical assistance.

Author Contributions

Conceptualization: Hans-Henrik Parving, Oluf Pedersen, Peter Rossing.

Data curation: Jens Christian Øllgaard, Bernt Johan von Scholten, Henrik Reinhard, Zeneng Wang, Peter Gæde, Oluf Pedersen.

Formal analysis: Signe Abitz Winther, Jens Christian Øllgaard, Tine Willum Hansen, Tarunveer Singh Ahluwalia, Stanley Hazen.

Investigation: Oluf Pedersen, Peter Rossing.

Methodology: Signe Abitz Winther, Oluf Pedersen.

Supervision: Tine Willum Hansen, Bernt Johan von Scholten, Tarunveer Singh Ahluwalia, Stanley Hazen, Oluf Pedersen, Peter Rossing.

Validation: Signe Abitz Winther.

Visualization: Signe Abitz Winther, Tarunveer Singh Ahluwalia.

Writing – original draft: Signe Abitz Winther.

Writing – review & editing: Signe Abitz Winther, Jens Christian Øllgaard, Tine Willum Hansen, Bernt Johan von Scholten, Henrik Reinhard, Tarunveer Singh Ahluwalia, Zeneng Wang, Peter Gæde, Hans-Henrik Parving, Stanley Hazen, Oluf Pedersen, Peter Rossing.

References

1. Gregg EW, Li Y, Wang J, Burrows NR, Ali MK, Rolka D, et al. Changes in diabetes-related complications in the United States, 1990–2010. *The New England journal of medicine*. 2014; 370(16):1514–23. <https://doi.org/10.1056/NEJMoa1310799> PMID: 24738668
2. Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nature medicine*. 2013; 19(5):576–85. <https://doi.org/10.1038/nm.3145> PMID: 23563705
3. Tang WH, Wang Z, Kennedy DJ, Wu Y, Buffa JA, Agatista-Boyle B, et al. Gut microbiota-dependent trimethylamine N-oxide (TMAO) pathway contributes to both development of renal insufficiency and mortality risk in chronic kidney disease. *Circulation research*. 2015; 116(3):448–55. <https://doi.org/10.1161/CIRCRESAHA.116.305360> PMID: 25599331
4. Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*. 2011; 472(7341):57–63. <https://doi.org/10.1038/nature09922> PMID: 21475195
5. Oellgaard J, Winther SA, Hansen TS, Rossing P, von Scholten BJ. Trimethylamine N-oxide (TMAO) as a New Potential Therapeutic Target for Insulin Resistance and Cancer. *Current pharmaceutical design*. 2017; 23(25):3699–712. <https://doi.org/10.2174/1381612823666170622095324> PMID: 28641532
6. Tang WH, Wang Z, Li XS, Fan Y, Li DS, Wu Y, et al. Increased Trimethylamine N-Oxide Portends High Mortality Risk Independent of Glycemic Control in Patients with Type 2 Diabetes Mellitus. *Clinical chemistry*. 2017; 63(1):297–306. <https://doi.org/10.1373/clinchem.2016.263640> PMID: 27864387
7. Schiattarella GG, Sannino A, Toscano E, Giugliano G, Gargiulo G, Franzone A, et al. Gut microbe-generated metabolite trimethylamine-N-oxide as cardiovascular risk biomarker: a systematic review and dose-response meta-analysis. *European heart journal*. 2017; 38(39):2948–56. <https://doi.org/10.1093/eurheartj/ehx342> PMID: 29020409
8. Croyal M, Saulnier PJ, Aguesse A, Gand E, Ragot S, Roussel R, et al. Plasma Trimethylamine N-oxide and risk of cardiovascular events in patients with type 2 diabetes. *The Journal of clinical endocrinology and metabolism*. 2020. <https://doi.org/10.1210/clinem/dgaa188> PMID: 32301490
9. Missailidis C, Hallqvist J, Qureshi AR, Barany P, Heimbürger O, Lindholm B, et al. Serum Trimethylamine-N-Oxide Is Strongly Related to Renal Function and Predicts Outcome in Chronic Kidney Disease. *PLoS one*. 2016; 11(1):e0141738. <https://doi.org/10.1371/journal.pone.0141738> PMID: 26751065
10. Zhu W, Gregory JC, Org E, Buffa JA, Gupta N, Wang Z, et al. Gut Microbial Metabolite TMAO Enhances Platelet Hyperreactivity and Thrombosis Risk. *Cell*. 2016; 165(1):111–24. <https://doi.org/10.1016/j.cell.2016.02.011> PMID: 26972052
11. Seldin MM, Meng Y, Qi H, Zhu W, Wang Z, Hazen SL, et al. Trimethylamine N-Oxide Promotes Vascular Inflammation Through Signaling of Mitogen-Activated Protein Kinase and Nuclear Factor-kappaB. *Journal of the American Heart Association*. 2016; 5(2). <https://doi.org/10.1161/JAHA.115.002767> PMID: 26903003
12. Winther SA, Ollgaard JC, Tofte N, Tarnow L, Wang Z, Ahluwalia TS, et al. Utility of Plasma Concentration of Trimethylamine N-Oxide in Predicting Cardiovascular and Renal Complications in Individuals With Type 1 Diabetes. *Diabetes care*. 2019; 42(8):1512–20. <https://doi.org/10.2337/dc19-0048> PMID: 31123156
13. Heianza Y, Ma W, Manson JE, Rexrode KM, Qi L. Gut Microbiota Metabolites and Risk of Major Adverse Cardiovascular Disease Events and Death: A Systematic Review and Meta-Analysis of Prospective Studies. *Journal of the American Heart Association*. 2017; 6(7). <https://doi.org/10.1161/JAHA.116.004947> PMID: 28663251
14. Wang Z, Tang WH, Buffa JA, Fu X, Britt EB, Koeth RA, et al. Prognostic value of choline and betaine depends on intestinal microbiota-generated metabolite trimethylamine-N-oxide. *European heart journal*. 2014; 35(14):904–10. <https://doi.org/10.1093/eurheartj/ehu002> PMID: 24497336
15. Mueller DM, Allenspach M, Othman A, Saely CH, Muendlein A, Vonbank A, et al. Plasma levels of trimethylamine-N-oxide are confounded by impaired kidney function and poor metabolic control. *Atherosclerosis*. 2015; 243(2):638–44. <https://doi.org/10.1016/j.atherosclerosis.2015.10.091> PMID: 26554714
16. Gaede P, Vedel P, Parving HH, Pedersen O. Intensified multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: the Steno type 2 randomised study. *Lancet (London, England)*. 1999; 353(9153):617–22. [https://doi.org/10.1016/S0140-6736\(98\)07368-1](https://doi.org/10.1016/S0140-6736(98)07368-1) PMID: 10030326
17. Gaede P, Oellgaard J, Carstensen B, Rossing P, Lund-Andersen H, Parving HH, et al. Years of life gained by multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: 21 years follow-up on the Steno-2 randomised trial. *Diabetologia*. 2016; 59(11):2298–307. <https://doi.org/10.1007/s00125-016-4065-6> PMID: 27531506

18. Reinhard H, Hansen PR, Persson F, Tarnow L, Wiinberg N, Kjaer A, et al. Elevated NT-proBNP and coronary calcium score in relation to coronary artery disease in asymptomatic type 2 diabetic patients with elevated urinary albumin excretion rate. *Nephrology, dialysis, transplantation: official publication of the European Dialysis and Transplant Association—European Renal Association*. 2011; 26(10):3242–9. <https://doi.org/10.1093/ndt/gfr009> PMID: 21372253
19. von Scholten BJ, Reinhard H, Hansen TW, Lindhardt M, Petersen CL, Wiinberg N, et al. Additive prognostic value of plasma N-terminal pro-brain natriuretic peptide and coronary artery calcification for cardiovascular events and mortality in asymptomatic patients with type 2 diabetes. *Cardiovascular diabetology*. 2015; 14:59. <https://doi.org/10.1186/s12933-015-0225-0> PMID: 25990319
20. Wang Z, Bergeron N, Levison BS, Li XS, Chiu S, Jia X, et al. Impact of chronic dietary red meat, white meat, or non-meat protein on trimethylamine N-oxide metabolism and renal excretion in healthy men and women. *European heart journal*. 2019; 40(7):583–94. <https://doi.org/10.1093/eurheartj/ehy799> PMID: 30535398
21. Carrillo S. Updated Information for IDMS Traceable VITROS[®] Chemistry Products CREA Slides [Internet], Rochester, NY, Ortho Clinical Diagnostics. Available from http://clinicalcomm/Downloads/OrthoClinicalDiagnostics-IDMS_20080612pdf. 2008.
22. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, et al. A new equation to estimate glomerular filtration rate. *Annals of internal medicine*. 2009; 150(9):604–12. <https://doi.org/10.7326/0003-4819-150-9-200905050-00006> PMID: 19414839
23. Andersen TF, Madsen M, Jorgensen J, Møller-Jørgensen L, Olsen JH. The Danish National Hospital Register. A valuable source of data for modern health sciences. *Danish medical bulletin*. 1999; 46(3):263–8. PMID: 10421985
24. Yusuf S, Dagenais G, Pogue J, Bosch J, Sleight P. Vitamin E supplementation and cardiovascular events in high-risk patients. *The New England journal of medicine*. 2000; 342(3):154–60. <https://doi.org/10.1056/NEJM200001203420302> PMID: 10639540
25. Coresh J, Turin TC, Matsushita K, Sang Y, Ballew SH, Appel LJ, et al. Decline in estimated glomerular filtration rate and subsequent risk of end-stage renal disease and mortality. *Jama*. 2014; 311(24):2518–31. <https://doi.org/10.1001/jama.2014.6634> PMID: 24892770
26. Oellgaard J, Gaede P, Rossing P, Persson F, Parving HH, Pedersen O. Intensified multifactorial intervention in type 2 diabetics with microalbuminuria leads to long-term renal benefits. *Kidney international*. 2017; 91(4):982–8. <https://doi.org/10.1016/j.kint.2016.11.023> PMID: 28187983
27. Brochner-Mortensen J. A simple method for the determination of glomerular filtration rate. *Scandinavian journal of clinical and laboratory investigation*. 1972; 30(3):271–4. <https://doi.org/10.3109/00365517209084290> PMID: 4629674
28. Guasch-Ferre M, Hu FB, Ruiz-Canela M, Bullo M, Toledo E, Wang DD, et al. Plasma Metabolites From Choline Pathway and Risk of Cardiovascular Disease in the PREDIMED (Prevention With Mediterranean Diet) Study. *Journal of the American Heart Association*. 2017; 6(11). <https://doi.org/10.1161/JAHA.117.006524> PMID: 29080862
29. Dong Z, Liang Z, Guo M, Hu S, Shen Z, Hai X. The Association between Plasma Levels of Trimethylamine N-Oxide and the Risk of Coronary Heart Disease in Chinese Patients with or without Type 2 Diabetes Mellitus. *Disease markers*. 2018; 2018:1578320. <https://doi.org/10.1155/2018/1578320> PMID: 30159101
30. Lever M, George PM, Slow S, Bellamy D, Young JM, Ho M, et al. Betaine and Trimethylamine-N-Oxide as Predictors of Cardiovascular Outcomes Show Different Patterns in Diabetes Mellitus: An Observational Study. *PloS one*. 2014; 9(12):e114969. <https://doi.org/10.1371/journal.pone.0114969> PMID: 25493436
31. Pelletier CC, Croyal M, Ene L, Aguesse A, Billon-Crossouard S, Krempf M, et al. Elevation of Trimethylamine-N-Oxide in Chronic Kidney Disease: Contribution of Decreased Glomerular Filtration Rate. *Toxins*. 2019; 11(11). <https://doi.org/10.3390/toxins11110635> PMID: 31683880
32. Zobel EH, von Scholten BJ, Reinhard H, Persson F, Teerlink T, Hansen TW, et al. Symmetric and asymmetric dimethylarginine as risk markers of cardiovascular disease, all-cause mortality and deterioration in kidney function in persons with type 2 diabetes and microalbuminuria. *Cardiovascular diabetology*. 2017; 16(1):88. <https://doi.org/10.1186/s12933-017-0569-8> PMID: 28697799
33. Rotbain Curovic V, Hansen TW, Eickhoff MK, von Scholten BJ, Reinhard H, Jacobsen PK, et al. Urinary tubular biomarkers as predictors of kidney function decline, cardiovascular events and mortality in microalbuminuric type 2 diabetic patients. *Acta diabetologica*. 2018; 55(11):1143–50. <https://doi.org/10.1007/s00592-018-1205-0> PMID: 30105469
34. Cardona A, O'Brien A, Bernier MC, Somogyi A, Wysocki VH, Smart S, et al. Trimethylamine N-oxide and incident atherosclerotic events in high-risk individuals with diabetes: an ACCORD trial post hoc

- analysis. *BMJ open diabetes research & care*. 2019; 7(1):e000718. <https://doi.org/10.1136/bmjdr-2019-000718> PMID: 31798892
35. Koay YC, Chen YC, Wali JA, Luk AWS, Li M, Doma H, et al. Plasma Levels of TMAO can be Increased with 'Healthy' and 'Unhealthy' Diets and Do Not Correlate with the Extent of Atherosclerosis but with Plaque Instability. *Cardiovascular research*. 2020.
 36. Zheng Y, Li Y, Rimm EB, Hu FB, Albert CM, Rexrode KM, et al. Dietary phosphatidylcholine and risk of all-cause and cardiovascular-specific mortality among US women and men. *The American journal of clinical nutrition*. 2016; 104(1):173–80. <https://doi.org/10.3945/ajcn.116.131771> PMID: 27281307
 37. Meyer KA, Shea JW. Dietary Choline and Betaine and Risk of CVD: A Systematic Review and Meta-Analysis of Prospective Studies. *Nutrients*. 2017; 9(7). <https://doi.org/10.3390/nu9070711> PMID: 28686188
 38. Rhee EP, Clish CB, Ghorbani A, Larson MG, Elmariah S, McCabe E, et al. A combined epidemiologic and metabolomic approach improves CKD prediction. *Journal of the American Society of Nephrology: JASN*. 2013; 24(8):1330–8. <https://doi.org/10.1681/ASN.2012101006> PMID: 23687356
 39. Gupta N, Buffa JA, Roberts AB, Sangwan N, Skye SM, Li L, et al. Targeted Inhibition of Gut Microbial TMAO Production Reduces Renal Tubulointerstitial Fibrosis and Functional Impairment in a Murine Model of Chronic Kidney Disease. *Arteriosclerosis, thrombosis, and vascular biology*. 2020: Atvbaha120314139.
 40. Wang Z, Roberts AB, Buffa JA, Levison BS, Zhu W, Org E, et al. Non-lethal Inhibition of Gut Microbial Trimethylamine Production for the Treatment of Atherosclerosis. *Cell*. 2015; 163(7):1585–95. <https://doi.org/10.1016/j.cell.2015.11.055> PMID: 26687352
 41. Roberts AB, Gu X, Buffa JA, Hurd AG, Wang Z, Zhu W, et al. Development of a gut microbe-targeted nonlethal therapeutic to inhibit thrombosis potential. *Nature medicine*. 2018; 24(9):1407–17. <https://doi.org/10.1038/s41591-018-0128-1> PMID: 30082863
 42. Liu HW, Tsai WH, Liu JS, Kuo KL. Association of Vegetarian Diet with Chronic Kidney Disease. *Nutrients*. 2019; 11(2). <https://doi.org/10.3390/nu11020279> PMID: 30691237
 43. Wang Z, Levison BS, Hazen JE, Donahue L, Li XM, Hazen SL. Measurement of trimethylamine-N-oxide by stable isotope dilution liquid chromatography tandem mass spectrometry. *Analytical biochemistry*. 2014; 455:35–40. <https://doi.org/10.1016/j.ab.2014.03.016> PMID: 24704102
 44. Shlipak MG, Matsushita K, Ärnlöv J, Inker LA, Katz R, Polkinghorne KR, et al. Cystatin C versus creatinine in determining risk based on kidney function. *The New England journal of medicine*. 2013; 369(10):932–43. <https://doi.org/10.1056/NEJMoa1214234> PMID: 24004120