

Original Article

Frequency of Malignancy in Solitary Thyroid Nodule and Multi-nodular Goitre

Md. Abul Hossain¹, Md. Zakaria Sarkar², Utpal Kumar Dutta³, Md. Abdul Karim⁴,
Md. Zahedul Alam⁵

Abstract

This cross sectional study with purposive sampling based on 50 cases of solitary thyroid nodule and multi-nodular goitre for operation in department of Otolaryngology and Head-Neck surgery of SSMC & Mitford Hospital, Dhaka and Dhaka Medical college, Dhaka during January 2009 to December 2009. Objectives of this study to find out relative frequency of malignancy in solitary thyroid nodule and multi-nodular goitre (by histopathology). Each case was studied by history taking, physical examination and relevant investigations. In this series, male to female ratio was 1:7. Mean age of the patients of nodular goitre was 36.5+12.33 years and the highest number of patients were found in age group 31-40 years.

After operation, histopathological report analysis was done and found that nodular colloid goitre was 52%, follicular adenoma was 24% and auto-immune thyroiditis was 6%. Among the malignancies papillary carcinoma was 66.66%, follicular carcinoma was 22.22% and anaplastic carcinoma was 11.11%. The relative frequency of malignancy in solitary thyroid nodule was 28% and in multi-nodular goitre was 8%. As there is a chance of malignancy is high so it should get serious medical attention.

Key words: Thyroid nodule, Solitary thyroid nodule, Multi-nodular goitre, Malignancy, Histopathology.

Introduction

Thyroid nodules are common in clinical practice. There may be solitary within a

normal thyroid gland or dominant within a multi-nodular goitre. The incidence of thyroid nodule has been on rise in recent decades mainly due to the wider use of neck imaging¹.

Nodules are usually multiple, forming a multi-nodular goitre. Occasionally, only one macroscopic nodule is found, but microscopic changes will be present throughout the gland, this is one form of clinically solitary nodule. Nodules appear early in endemic goitre and later (between 20 and 30 years) in sporadic goitre².

Palpable thyroid nodules occur in 4-7% of the population but nodules found incidentally on ultrasonography suggest a prevalence of

1. Assistant Professor, National Institute of ENT, Dhaka
2. Assistant Professor, National Institute of ENT, Dhaka
3. Assistant Professor (C.C), National Institute of ENT, Dhaka
4. Junior Consultant, National Institute of ENT, Dhaka
5. Director & Professor, National Institute of ENT, Dhaka

Address of Correspondence: Dr. Md. Abul Hossain, Assistant Professor, National Institute of ENT, Dhaka. E-mail: abulhossain619@gmail.com

19-67%. The majority of the thyroid nodules are asymptomatic. Because about 5% of all palpable nodules are found to be malignant, the main objective of evaluating thyroid nodules is to exclude malignancy³.

Iodine deficiency is the main cause of goitre development in Bangladesh. In Bangladesh, goitre is prevalent in bank of river Jamuna, northern part of the country and also hilly areas in Sylhet and Chittagong. In clinical practice, we consider nodular goitre either with solitary nodule or multiple nodules in different stages of development⁴.

Nodules in the thyroid gland are important for their malignant potential. It is the highest among the cancer affecting endocrine glands. Cancer of the thyroid gland occurs at earlier ages in most parts of the world. It is commonest between 20-40 years of age⁵.

Risk factors that increase the probability of malignancy of a thyroid nodule are age under 30 years or over 60 years, male sex (8% in male versus 4% in female), history of head neck irradiation in childhood and family history of medullary thyroid carcinoma or multiple endocrine neoplasia (MEN) type 2¹.

Early diagnosis helps in early treatment, results in good outcome. But late diagnosis indicates advanced stage of disease with unsatisfactory treatment and poor prognosis. Frequency of malignancy in thyroid nodules varies among different studies in our country & abroad. In our country, one of the study was found, the percentage of malignancy 21.44% & 8.1% in solitary nodule & multi-nodular goitre respectively⁵. Another study showed chance of malignancy 9.89%⁶. One of the study in abroad found, 5% of the thyroid nodules as malignant⁷.

Purpose of this study was to find out the relative frequency of malignancy in solitary thyroid nodule and multi-nodular goitre.

Methods

This cross-sectional comparative study was done in the department of Otolaryngology &

Head-Neck surgery of Sir Salimullah Medical College Hospital and Dhaka Medical College Hospital during the calendar year 2009. 50 admitted cases of solitary thyroid nodules and multi nodular goiter matching the inclusion and exclusion criteria were included in this study.

Results

Table- I

Age distribution in solitary and multi-nodular goitre, (n=50)

Age (years)	Solitary thyroidnodule	Multi-nodulargoitre	Total (%)
11-20	5	2	7 (14%)
21-30	3	2	5(10%)
31-40	8	13	21(42%)
41-50	7	5	12(24%)
51-60	2	1	3 (6%)
61-70	0	2	2 (4%)
Total	25	25	50 (100%)

Table- II

Sex distribution in solitary and multi-nodular goitre, (n=50)

Sex	Solitary thyroidnodule	Multi-nodulargoitre	Total (%)
Male	4	2	6 (12%)
Female	21	23	44 (88%)
Total	25	26	50(100%)

Table- III

Distribution of habitat of patients

Habitat	Solitary thyroidnodule	Multi-nodulargoitre	Total (%)
Urban	6	5	11 (22%)
Rural	19	20	39 (78%)
Total	25	25	50 (100%)

Table- IV
Histopathological diagnosis

Diagnosis	Solitary thyroid nodule	Multi-nodular goitre	Total (%)
Multi-nodular colloid goitre	0	15	15 (30%)
Single nodular colloid goitre	11	0	11 (22%)
Follicular Adenoma	7	5	12 (24%)
Papillary carcinoma	6	0	6 (12%)
Follicular carcinoma	1	1	2 (4%)
Medullary carcinoma	0	1	1 (2%)
Anaplastic carcinoma	0	1	1 (2%)
Auto immune thyroiditis	0	3	3 (6%)
Total	25	25	50 (100%)

Table- V
Frequency of malignancy in solitary thyroid nodule & multi-nodular goitre.

Types	Malignant	Non malignant	Total (%)	Relative frequency (%)
Solitary thyroid nodule	7	18	25 (50%)	28%
Multi-nodular goitre	2	23	25 (50%)	8%

$z = 2$ $p < 0.05$

Table-VI
Distribution of thyroid malignancy (n = 14)

Name of malignancy	Number of cases		Percentage
	Solitary thyroid	Multi-nodular goitre	
Papillary carcinoma	6	0	66.66%
Follicular carcinoma	1	1	22.22%
Anaplastic carcinoma	0	1	11.11%

Discussion

In this study, 50 patients with thyroid swelling (solitary & multi-nodular goitre) were studied. Mean age of the patients of nodular goitre was 36.5 ± 12.33 years and the highest frequency (42%) was in 31-40 years. Which correlated with study of Rahman. Nath, Sattar^{5,8}. The youngest patient in this study

was a girl of 15 years with a papillary carcinoma and the oldest patients was a lady of 70 years with medullary carcinoma. The youngest patient and oldest patients of this study both had been suffering from malignant thyroid disease, the extreme of ages show less incidence of thyroid disease but has a more chance to be malignant.

In this series, out of 50 patients, male were 6 (12%) and female were 44 (88%). Male female ratio was (1:7.31). This ratio was shown 1:5 by Rahman⁵ (2000), 1:4 by Welkar³, 1:2.5 to 4.1 Zuberi⁹. This female preponderance is reflected in all studies including the present. The cause of high female to male ratio in this series can be explained by most of the patients are from non-endemic area. Here we may recall the finding of Kilpatrick et al who found a male to female ratio of 1: 14 in non endemic area, which was 1:1 in endemic area.

In this study the commonest occupational group was house wife (58%). It was due to the fact that thyroid disorders are female prone owing to the presence of estrogen receptors in the thyroid tissue².

The most of the patients in this series came from rural areas (88%) having a socio-economic status < 10,0007 per month (54%).

In this series, after operation, histopathological report analysis was done and found that nodular colloid goitre was 52% (30% + 22%), follicular adenoma was 24%, papillary carcinoma was 12%, follicular carcinoma was 4% & autoimmune thyroiditis was 6%. Among the malignancies papillary carcinoma was 66.66%, follicular carcinoma was 22.22% and anaplastic carcinoma each was 11.11%. It was almost similar to study of Zygmunt and Meckenzic¹⁰.

In this series, relative frequency of malignancy in solitary thyroid nodule was 28% and in multi-nodular goitre was 8% which correlate with study of Asraf,^{5,6,8} Rahman⁵ and Satter. In study of Rahman in solitary thyroid nodule, malignancy was 21.44% and in multi-nodular goitre was 8.1%. Another study shown & chance of malignancy 9.89%⁶. One of the study in abroad found 5% of thyroid nodule was malignant⁷. It is observed that relative

frequency of malignancy was higher than that of others. It might be happened due to small sample size, sampling bias, by chance, advanced stage of disease our patients took surgical treatment in advanced stage of disease. In this study frequency of malignancy differs significantly with age & sex, older patients and male sex are more prone to develop malignancy ($p < 0.05$).

In this study observed that follicular carcinoma occurred in the age group 31-40 years but follicular carcinoma usually occurs in the age group 50-59 years. So it is a matter of thinking that follicular carcinoma is occurring in early age group. But it may be a good sign that our patients are becoming aware regarding thyroid nodule and are attending in the hospital for early surgical treatment.

Summary

Clinically nodular goitre is the most common problem of thyroid disease in all ages. Clinical importance of thyroid nodule is exclusion of malignancy. Female was more commonly affected than male. The highest number of patients were found in age group 31-40 years. All the patients were in euthyroid state clinically & also biochemically. Clinically solitary nodular goitre was found 56% and rest was multi-nodular goitre.

After operation histopathological examination (gold standard investigation) revealed nodular colloid goitre 52%, follicular adenoma 34%, papillary carcinoma 12%, follicular carcinoma 4%, anaplastic carcinoma 2% and autoimmune thyroiditis 6%. Among malignancies papillary carcinoma was 66.66%, follicular carcinoma 22.22%, anaplastic carcinoma each was 11.11%. Relative frequency of malignancy in solitary thyroid nodule and in multi-nodular goitre was 28% and 8% respectively.

Conclusion

A significant proportion of solitary thyroid nodule & multi-nodular goitre may be malignant. So, it should get appropriate medical attention. Our people should be offered the early treatment of thyroid nodule is better and late treatment of even asymptomatic thyroid nodule may be fatal.

References

1. Polyzos SA, Kita M & Avramidis. Thyroid nodules-Stepwise diagnosis and Management. *Hormones*. 2007; 6 (2): 101-119
2. Krukowski ZH. The thyroid gland and thyroglossal tract, Baily & Love's short practice of surgery. 24th ed. London. Hodder education, 2004;776-804.
3. Welker MJ & Lov. Thyroid nodules. *American Family / Physician*, Feb. 1: 1-11.
4. Alam MM, Hossain D, Datta PG & Alauddin M. Study of thyroid carcinoma in solitary thyroid nodule; *Bangladesh J of otorhinolaryngology*, 2004; 10 (1) : 7-12.
5. Rahman MJ & Mustafa MG. Comparative study of cancer developing in solitary thyroid nodule and multi-nodular goitre. *Bangladesh J of otorhinolaryngology*. 2000; 6(11): 6-12.
6. Ashraf SA & Matin SA MR. A Review of thyroid diseases in Bangladesh. *Journal of BCPS*. 21996; 2(1): 6-10.
7. Gandolfi PP, Frisina A & Raffa M The incidence of thyroid carcinoma in multi-nodular goitre, *Acta Bio Medica Ateneo parmense*, 2004; 75: 114-117.
8. Sattar MA. Alam MR & Haider A. Clinico Pathological study of solitary cold thyroid nodule, *Bangladesh J of otorhinolaryngology*, 2003; 9 (1):24-27.
9. Zuberi LM, Yawar A, Islam & Jabbar A, Clinical presentation of thyroid cancer patients in Pakistan- AKUH Experience. *Journal of Pakistan Association*, 2009; Available from <http://Jpma.org.pk/viewv> Article, accessed 7 July.
10. Mackenzie EJ & Mortimer RH. Thyroid nodules and thyroid Cancer. *The medical Journal of Australia*, 2004; 180 (5): 242-247.