



Epidemiological analysis of drinking water-type fluorosis areas and the impact of fluorosis on children's health in the past 40 years in China

Feiqing Wang · Yanju Li · Dongxin Tang ·
Jianing Zhao · Bo Yang · Chike Zhang · Min Su ·
Zhixu He · Xiaodong Zhu · Dong Ming · Yang Liu

Received: 31 December 2022 / Accepted: 28 September 2023 / Published online: 31 October 2023
© The Author(s) 2023

Abstract This study analyzed the effect of China's fluorosis prevention and control program, which has been in effect for more than 40 years, and the impact of fluorosis on children's health. Relevant research studies were retrieved from the following online databases from the time of their inception to May 2022: PubMed, ScienceDirect, Embase, Cochrane, China National Knowledge Infrastructure, and Wanfang. The Review Manager 5.3 software was used in statistical analyses. This article included seventy studies: Thirty-eight studies reported the effect of improving water quality and reducing fluoride content, the

incidence rate of dental fluorosis in children, and the level of urinary fluoride, and thirty-two studies reported the intelligence quotient (IQ) and health status of children. Following water improvement strategies, the fluoride levels in drinking water decreased significantly; urinary fluoride levels and dental fluorosis decreased significantly in children. With regard to the effect of fluorosis on the IQ of children, the results showed that the IQ of children in areas with a high fluoride of fluorosis was lesser than that in areas with a low fluoride, and this difference was significant. Based on the prevalence of dental fluorosis and its effect on the intelligence of children, it appears that reducing fluoride levels in drinking water and monitoring water quality are important strategies for the prevention and treatment of fluorosis.

Feiqing Wang and Yanju Li have contributed equally to this work.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10653-023-01772-9>.

F. Wang · X. Zhu (✉) · D. Ming
Academy of Medical Engineering and Translational
Medicine, Tianjin University, No. 92 Weijin Road, Nankai
District, Tianjin City 300072, China
e-mail: zxdttju@163.com

D. Ming
e-mail: mindtju@126.com


F. Wang · D. Tang · J. Zhao · B. Yang · Y. Liu (✉)
Clinical Research Center, The First Affiliated Hospital
of Guizhou University of Traditional Chinese Medicine,
No. 71 Bao Shan North Road, Yunyan District,
Guiyang 550001, Guizhou Province, China
e-mail: ly7878@163.com

Y. Li · C. Zhang
Clinical Research Institute, Affiliated Hospital of Guizhou
Medical University, Guiyang 550004, Guizhou Province,
China

M. Su · Z. He · Y. Liu
National and Guizhou Joint Engineering Laboratory
for Cell Engineering and Biomedicine Technique,
Guiyang 550004, Guizhou Province, China

X. Zhu
Neurological Institute, Tianjin Medical University General
Hospital, Tianjin 300072, China

Graphical abstract



Keywords Drinking water fluorosis · Defluoridation · Intelligence quotient · Health status · Fluorosis area

Introduction

According to the World Health Organization (WHO), fluorosis has become a major health problem worldwide, as over 260 million people globally are drinking water from sources with high-fluoride concentrations (Amini et al., 2008). Drinking water is the most important pathway of fluoride intake, as it accounts for 75–90% of fluoride intake (Habtmu et al., 2019; Kloos & Haimanot, 1999). China has the highest prevalence of fluorosis and faces the most serious harmful effects of fluorosis in the world. This is because China is located in the fluoride belt. Excess

fluoride levels have been detected in the groundwater in areas of China where the fluoride content is high in rocks or soil. Fluorosis caused by the consumption of drinking water with high-fluoride content, which will henceforth be referred to as drinking water fluorosis, is one of the most important types of endemic fluorosis in China, and it is prevalent in 27 provinces, cities, and autonomous regions across the North and South of China (Li et al., 2020a, 2020b, 2020c; Wang et al., 2012a, 2012b).

The severity of fluorosis is associated with the concentration of fluoride in drinking water, daily intake, duration of exposure, and climatic conditions (Li et al., 2021). Exposure to high-fluoride levels for a prolonged period results in dental fluorosis, skeletal fluorosis, and non-skeletal fluorosis, which manifests as muscle weakness, tiredness, anemia, dyspepsia, male infertility, decrease in IQ, and other symptoms

(Everett, 2011; Kumar et al., 2009; Shashi et al., 2008; Wei et al., 2019). Epidemiological investigations have found that fluorosis can severely affect any age group and can range from mild dental fluorosis to crippling skeletal fluorosis. Its effects are especially harmful to developing children, and its adverse effects are irreversible (Alexandra et al., 2021; Alvee et al., 2021). In 1937, Murray first reported the manifestation of nervous system dysfunction in patients with endemic fluorosis (Murray, 1937). Since then, other studies have found that long-term excessive intake of fluoride can cause demyelinating changes in the cerebral cortex and subcortical areas and lead to hypothyroidism; this may explain the decline in the intelligence level of children in high-fluoride areas (Fan, 2020; Saeed et al., 2020; Wang et al., 2021a, 2021b, 2021c). As a result of these findings, and the fact that approximately 100 million people in 1115 counties are presently affected by fluoride threats in China (Wang et al., 2020), the functional and organic effects of fluorosis on the central nervous system, especially on children's intellectual development, learning, and memory, have been receiving increasing attention.

The most effective way to prevent and treat fluorosis is to improve water quality and reduce water fluoride content (James et al., 2021; Wang et al., 2021a, 2021b, 2021c). So far, there is no effective clinical treatment for endemic fluorosis, and reducing fluoride exposure level and reducing fluoride intake are the main measures adopted by most countries (Malin et al., 2019; Zhao et al. 2013; Mohd Nor et al., 2021). In China, since 1978, fluoride-safe drinking water supply schemes have been implemented to provide fluoride-safe drinking water to populations in high-fluoride areas and fluorosis endemic areas (Li et al., 2020a, 2020b, 2020c). Through decades of effort on the part of the government, the fluoride concentrations in the drinking water supplied to many villages have been under control. However, due to differences in the implementation periods and management strategies, there are still significant differences in the effects of these water improvement and defluoridation projects among various provinces. At present, the effects of the water improvement and fluoride reduction projects that have been implemented in China for more than half a century have not been systematically analyzed and studied. Therefore, this study was undertaken to comprehensively assess the effects

of these projects on the prevention and treatment of drinking water-borne endemic fluorosis and the effects of fluorosis on the development and intelligence levels of children. These findings are important from the perspective of monitoring the effects of these projects and highlighting potential areas of improvement.

Methods

Literature retrieval strategy

The PubMed, ScienceDirect, Embase, Cochrane, China National Knowledge Infrastructure (CNKI), and Wanfang databases were searched for relevant articles deposited from their inception to May 2022, using the following search terms in the title, keyword, or abstract: water improvement, defluorination, dental fluorosis, skeletal fluorosis, children, student, intelligence, intelligence quotient, nervous system, and behavior. Additionally, we performed a backward literature search by manually retrieving the references included in the studies and related reviews.

Inclusion and exclusion criteria

Three investigators independently performed the data extraction. This study evaluated the content and scope of previously published reports and generated inclusion and exclusion criteria. The following were the inclusion criteria: publication from the inception of each database to May 2022, which belongs to the exposure group (fluoride content in drinking water > 1.0 mg/L), dental fluorosis in children, skeletal fluorosis in adults, and data on urinary fluoride levels in children and/or adults. Combined Raven's test (CRT) measures a child's IQ (Julvez et al., 2021; Theobald et al., 2018). Based on the IQ level, the data were divided into seven categories: very excellent (IQ > 130), excellent (IQ = 120–129), upper middle (IQ = 110–119), medium (IQ = 90–109), lower middle (IQ = 80–89), marginal (IQ = 70–79), and mental retardation (IQ < 69). The main advantage of the combined Raven test is that it assesses the basic factors that represent children's intelligence structure, and it is therefore suitable for collective testing.

Data analysis

This research evaluated the study content and scope of previously published reports. A list of relevant data from each study was created, including publication time, type of study, time and place of investigation, duration of water treatment, prevalence of dental fluorosis in children between the ages of 8 and 15 years, prevalence of skeletal fluorosis in adults, fluoride levels in drinking water, fluoride levels in urine and child's IQ, etc. When there was heterogeneity, sensitivity analysis or appropriate statistical methods were used to estimate the effect. The heterogeneity of the research indicators was analyzed to determine whether the indicators could be pooled for analysis. The data included.

Statistical analysis

All statistical analyses were completed utilizing the Review Manager version 5.3. The chi-square test (χ^2) was used to evaluate heterogeneity among the studies ($P=0.05$). Heterogeneity among studies was assessed using the I^2 statistic. 25%, 50%, and 75% were designated as low, medium, and high for I^2 values, respectively. We performed the study using a fixed-effect model, where with $I^2 > 50\%$ we used random-effect model. Weighted mean difference (WMD) and standardized mean difference (SMD) were used for continuous variables, and odds ratio (OR) was used for binary variables and reported pooled estimates and the corresponding 95% confidence interval (CI). Study quality assessment and risk of bias were assessed for all studies. In the study, differences were considered statistically significant when the p value was ≤ 0.05 .

Results

Literature search results

Out of a total of 2736 studies that were retrieved, 119 were obtained from PubMed, ScienceDirect, Embase, and Cochrane and 2617 were obtained from CNKI and Wanfang. Out of the 2736 studies, 1589 were duplicates: 57 were included in PubMed, ScienceDirect, Embase, and Cochrane, and 329 were included in CNKI and Wanfang. After application of the exclusion

criteria, a total of seventy studies were finally included in the meta-analysis: thirty-eight studies reported improvement in the incidence of fluorosis and thirty-two studies reported on children's intelligence levels in fluorosis areas (Fig. 1).

The seventy included research articles were cross-sectional studies published between 1995 and 2021. Thirty-eight studies reported on water improvement projects that were carried out over a period of 5–40 years and covered 22 provinces, cities, and counties. Thirty-two studies reported on children's intelligence levels in fluorosis areas and covered 14 provinces, cities, and counties. Thirty-eight of the studies reported the prevalence of dental fluorosis in children in 32,042 cases, and thirty-four studies reported the fluoride levels in drinking water (Table S1). Twenty-seven studies reported intelligence levels in 14,617 children in fluorosis areas, and eleven studies report children's IQ scores (Table S1).

Epidemiological analysis of reducing water fluoride content in fluorosis areas

Thirty-four studies (Table S1) reported fluoride levels in drinking water, and the results showed that the fluoride content in water decreased significantly (95% CI – 2.72 to – 1.96, $P < 0.05$; Table 1). For subgroup analysis, the studies were divided into those from South China (95% CI – 4.69 to – 1.55) and North China (95% CI – 2.38 to – 1.81), and the studies were further divided into six areas of China: North China (95% CI – 2.86, – 1.49), Northeast China (95% CI – 2.10 to – 0.68), East China (95% CI 3.26 to – 1.81), Central China (95% CI – 8.01 to 0.25), South China (95% CI – 2.97 to – 1.49), and Northwest China (95% CI – 3.59 to – 1.02), and the fluoride content in water decreased significantly (all $P < 0.05$; Table 1). Subgroup analysis was also conducted according to the duration of the water improvement: 0–10 years (95% CI – 1.95 to 1.29), 11–20 years (95% CI – 7.17 to – 1.37), and 21–30 years (95% CI – 2.82 to – 1.61), the differences were statistically significant ($P < 0.05$; Table 1).

Epidemiological analysis of urinary fluoride levels and incidence of skeletal fluorosis in adults after water improvement

Six studies (Table S1) reported urinary fluoride levels in adults. The research findings




Fig. 1 Flowchart depicting the study selection process

showed (WMD = −2.01, 95% CI −2.81 to −1.21, $P < 0.00001$) there was a statistically significant decrease in urinary fluoride levels ($P < 0.05$; Fig. 2). Eleven studies (Table S1) reported the prevalence of skeletal fluorosis in adults. The research findings showed (OR = 0.33, 95% CI 0.20–0.55, $P < 0.0001$) a statistically significant decrease in skeletal fluorosis ($P < 0.05$; Fig. 2).

Epidemiological analysis of dental fluorosis prevalence in children after water improvement

Thirty-eight studies (Table S1) reported data on dental fluorosis in children. These studies also showed


(OR = 0.13, 95% CI 0.10–0.18, $P < 0.00001$) a significant difference in the prevalence of decreased dental fluorosis in children ($P < 0.05$; Table 2). Using subgroup analysis, the following results were obtained: South China (95% CI 0.06–0.14) and North China (95% CI 0.11–0.23), and the differences between the two regions were statistically significant ($P < 0.05$; Table 2). The studies were further divided into six areas: North China (95% CI 0.15–0.41), North-east China (95% CI 0.10–0.35), East China (95% CI 0.05–0.19), Central China (95% CI 0.04–0.18), South China (95% CI 0.01–0.38), and Northwest China (95% CI 0.03–0.44) (all $P < 0.05$; Table 2). The studies were divided based on the duration of the

Table 1 Comparison of water improvement and fluoride reduction among in different areas

Area	Test for heterogeneity		Analysis model	Test for overall effect		WMD or SMD	95% CI
	I^2 (%)	P		Z	P		
Total	100	<0.00001	Random	12.02	<0.00001	−2.34	(−2.72, −1.96)
Two areas							
South	100	<0.00001	Random	3.89	=0.0001	−3.12	(−4.69, −1.55)
North	99	<0.00001	Random	12.31	<0.00001	−2.09	(−2.38, −1.81)
Six areas							
North China	99	<0.00001	Random	6.23	<0.00001	−2.17	(−2.86, −1.49)
Northeast China	98	<0.00001	Random	3.85	=0.0001	−1.39	(−2.10, −0.68)
East China	99	<0.00001	Random	6.85	<0.00001	−2.53	(−3.26, −1.81)
Central China	100	<0.00001	Random	1.84	$P=0.07$	−3.88	(−8.01, 0.25)
South China	93	<0.00001	Random	5.89	<0.00001	−2.23	(−2.97, −1.49)
Northwest China	99	<0.00001	Random	3.51	=0.0004	−2.31	(−3.59, −1.02)
Time							
0–10 Years	100	<0.00001	Random	9.54	<0.00001	−1.62	(−1.95, 1.29)
11–20 Years	100	<0.00001	Random	2.89	=0.004	−4.27	(−7.17, −1.37)
21–30 Years	98	<0.00001	Random	7.13	<0.00001	−2.22	(−2.82, −1.61)

WMD weighted mean difference, SMD standardized mean difference, 95% CI 95% confidence intervals

A



B


**Fig. 2** Urinary fluoride levels and incidence of skeletal fluorosis in adults. **A** Urinary fluoride levels in adults. **B** Prevalence of skeletal fluorosis in adults

Table 2 Comparison of the prevalence of dental fluorosis in children among in different areas

Area	Test for heterogeneity		Analysis model	Test for overall effect		Odds ratio	95% CI	Unimproved	Improved
	I^2 (%)	P		Z	P			Total/events	Total/events
Total	99	<0.00001	Random	12.63	<0.00001	0.13	(0.10, 0.18)	96,442/52217	121,699/25918
Two areas									
North	97	<0.00001	Random	11.05	<0.00001	0.09	(0.06, 0.14)	48,553/22452	111,212/23385
South	99	<0.00001	Random	9.25	<0.00001	0.16	(0.11, 0.23)	47,889/29765	10,487/2533
Six areas									
North China	99	<0.00001	Random	5.26	<0.00001	0.24	(0.15, 0.41)	26,122/12658	54,084/17912
Northeast China	98	<0.00001	Random	5.34	<0.00001	0.19	(0.10, 0.35)	11,276/2381	42,619/2892
East China	96	<0.00001	Random	7.11	<0.00001	0.10	(0.05, 0.19)	18,765/10176	7912/1664
Central China	94	<0.00001	Random	6.23	<0.00001	0.08	(0.04, 0.18)	2677/1985	1720/404
South China	85	=0.001	Random	3.05	=0.002	0.07	(0.01, 0.38)	27,572/18337	1968/641
Northwest China	100	<0.00001	Random	3.11	=0.002	0.11	(0.03, 0.44)	10,030/6680	13,396/2405
Time									
0–10 Years	98	<0.00001	Random	8.10	<0.00001	0.22	(0.15, 0.32)	23,964/10715	33,636/5423
11–20 Years	100	<0.00001	Random	4.04	<0.0001	0.11	(0.04, 0.32)	41,689/27056	38,028/15426
21–30 Years	99	<0.00001	Random	4.47	<0.00001	0.07	(0.02, 0.22)	24,437/11934	11,847/1951
Degree of dental fluorosis in children									
Total	98	<0.00001	Random	3.54	=0.0004	0.65	(0.51, 0.82)	154,932/25823	316,962/52974
Normal	96	<0.00001	Random	10.16	<0.00001	5.07	(3.71, 6.94)	25,822/12156	52,827/43583
Suspect	97	<0.00001	Random	1.19	=0.23	1.30	(0.84, 2.02)	25,822/2437	52,827/4389
Very mild	93	<0.00001	Random	3.37	=0.0007	0.61	(0.46, 0.81)	25,822/5290	52,827/3408
Mild	86	<0.00001	Random	8.25	<0.00001	0.29	(0.22, 0.39)	25,822/3477	52,827/1240
Moderate	77	<0.00001	Random	8.26	<0.00001	0.16	(0.10, 0.25)	25,822/1725	52,827/298
Severe	87	<0.00001	Random	5.39	<0.00001	0.05	(0.02, 0.14)	25,822/738	52,827/56

95% CI 95% confidence intervals

water improvement projects into 0–10, 11–20, and 21–30 years, and the differences between the three groups were statistically significant (all $P < 0.05$; Table 2). Nineteen studies (Table S1) reported the degree of dental fluorosis in children after water improvement. The studies showed (OR=0.65, 95% CI 0.51–0.82, $P=0.0004$) a statistically significant damage decrease in the degree of dental fluorosis in children ($P < 0.05$; Table 2).

Epidemiological analysis of Urinary fluoride levels in children after water improvement

Fifteen studies (Table S1) reported urinary fluoride levels, and the research findings showed (WMD=−2.54, 95% CI−3.12 to−1.95,

$P < 0.00001$) a statistically significant decrease in urinary fluoride levels in children after implementation of the water improvement ($P < 0.05$; Fig. 3).

Epidemiological analysis of IQ level in children in fluorosis areas

Twenty-seven studies (Table S1) reported IQ levels in children in normal areas ($n=7133$) and fluorosis areas ($n=7484$). The results showed children living in fluorosis areas had significantly lower IQ levels than children living in normal areas ($P < 0.05$, 95% CI−8.95 to−5.16, Table 3). For subgroup analysis, the studies were divided into mild-to-moderate and severe fluorosis areas, for which the following results were obtained: mild-to-moderate fluorosis areas:




Fig. 3 Urinary fluoride levels in children

WMD=6.64, 95% CI−9.04 to −4.24, $P < 0.00001$, and severe fluorosis areas: WMD=−7.47, 95% CI−10.20 to −4.75, $P < 0.00001$. The differences between the fluorosis and normal areas were statistically significant ($P < 0.05$; Table 3). The studies were also divided into those reporting drinking water-borne fluorosis and coal burning-associated fluorosis, for which the results were WMD=−6.11, $P < 0.00001$ and WMD=−13.59, $P < 0.00001$, respectively. The differences between the groups were statistically significant ($P < 0.05$; Table 3).

Analysis of the IQ scores of children in fluorosis areas

Eleven studies (Table S1) reported the degree of IQ in children. When the number of children under each IQ category was compared between children in normal

areas ($n = 3361$) and children in severe fluorosis areas ($n = 4027$), a significantly lower number of children from normal areas fell under the lower IQ categories. However, a significantly higher number of children from normal areas than fluorosis areas fell under the higher IQ categories ($P < 0.00001$; Table 4).

Analysis of IQ in children with dental fluorosis in fluorosis areas

Seven studies (Table S1) reported the IQ levels in children with normal dentition ($n = 1185$) and dental fluorosis ($n = 1169$) in fluorosis areas. The results showed there was a statistically significant decrease in IQ levels between children with dental fluorosis and those with normal dentition in fluorosis areas ($P < 0.05$; 95% CI−13.68 to −8.20, Fig. 4). Additionally, three studies (Table S1) reported the

Table 3 Comparison of IQ level children in fluorosis area and normal area

Class	Test for heterogeneity		Analysis model	Test for overall effect		WMD or SMD	95% CI
	I^2 (%)	P		Z	P		
Total	94	<0.00001	Random	7.29	<0.00001	−7.06	(−8.95, −5.16)
Fluorosis area							
Mild to moderate	95	<0.00001	Random	5.43	<0.00001	−6.64	(−9.04, −4.24)
Severe	91	<0.00001	Random	5.38	<0.00001	−7.47	(−10.20, −4.75)
Type							
Drinking water	93	<0.00001	Random	6.93	<0.00001	−6.11	(−7.84, −4.38)
Coal burning	74	=0.009	Random	5.56	<0.00001	−13.59	(−18.38, −8.80)

WMD weighted mean difference, SMD standardized mean difference, 95% CI 95% confidence intervals

Table 4 Comparison of IQ scores of children in fluorosis area and normal area

IQ class	Test for heterogeneity		Analysis model	Test for overall effect		Odds ratio	95% CI	Fluorosis Total/events	Normal Total/events
	I^2 (%)	P		Z	P				
< 69	0	= 0.65	Fixed	6.56	< 0.00001	3.02	(2.17, 4.20)	4027/213	3361/45
70–79	41	= 0.07	Fixed	1.35	= 0.0007	1.35	(1.09, 1.69)	4027/276	3361/135
80–89	27	= 0.19	Fixed	2.48	= 0.01	1.22	(1.04, 1.43)	4027/498	3361/303
90–109	49	= 0.03	Fixed	0.94	= 0.35	0.96	(0.87, 1.05)	4027/1759	3361/1520
110–119	24	= 0.21	Fixed	0.18	= 0.86	0.99	(0.88, 1.11)	4027/806	3361/744
120–129	30	= 0.16	Fixed	2.69	= 0.007	0.81	(0.70, 0.94)	4027/389	3361/434
> 130	0	= 0.75	Fixed	6.13	< 0.00001	0.44	(0.34, 0.57)	4027/86	3361/180

95% CI 95% confidence intervals

degree of IQ in the two groups of children, and statistically significant differences were found for all IQ levels ($P < 0.05$; Table S2).

Effects of fluorosis on thyroid hormone levels, language IQ, weight, and height

Five studies (Table S1) reported thyroid hormones levels in children from normal areas and fluorosis areas. Significant differences were found for TSH ($P = 0.009$, 95% CI 0.29–1.99), T4 ($P < 0.0001$, 95% CI –4.30 to –3.09), and FT4 ($P < 0.0001$, 95% CI –1.54 to –0.54), but no significant differences were found for T3 ($P = 0.52$) and FT3 ($P = 0.81$) (Table 5). Two studies (Table S1) reported language IQ and performance IQ in children from normal areas and fluorosis areas, and significant differences were found for all categories (Table 5). Six studies (Table S1) reported height and weight in children from normal areas and fluorosis areas, but there was no statistically significant difference between the two groups.

Discussion

This is the first study to retrospectively analyze the relationship between water fluoride content, urine fluoride, dental fluorosis, project duration, fluorosis incidence, and children's health and IQ in fluorosis areas after the implementation of China's water improvement and defluoridation project. The findings provide important insight into the impact of these programs on the incidence of fluorosis and the effect of fluorosis on the intelligence level of children. They also highlight some areas of the project that need attention in the future.

The present data show that there was a significant reduction in water fluoride levels following the implementation of the water improvement and defluoridation projects in different provinces in China, which is in accordance with WHO standards. The findings showed that the fluoride content of water in the South was higher than that in the North. After water improvement and defluoridation, the fluoride content of water in the South was lower than that in the North. According to the results for different districts, the water fluoride content in Central China was the



Fig. 4 IQ levels in children with normal teeth and dental fluorosis in fluorosis areas

Table 5 Analysis of thyroid hormone level and language IQ of children in fluorosis area

IQ class	Test for heterogeneity		Analysis model	Test for overall effect		WMD or SMD	95% CI
	I^2 (%)	P		Z	P		
Thyroid hormones							
TSH	97	<0.00001	Random	2.62	=0.009	1.14	(0.29, 1.99)
T3	81	=0.005	Random	0.64	=0.52	0.08	(−0.16, 0.32)
T4	0	=0.82	Random	11.97	<0.00001	−3.70	(−4.30, −3.09)
FT3	73	=0.05	Random	0.24	=0.81	0.05	(−0.36, 0.46)
FT4	0	=0.60	Random	4.09	<0.0001	−1.04	(−1.54, −0.54)
Language IQ performance IQ							
Common sense	0	=0.59	Random	5.33	<0.00001	−2.62	(−3.59, −1.66)
Similar	80	=0.03	Random	3.27	=0.001	−4.41	(−7.05, −1.77)
Arithmetic	65	=0.09	Random	5.28	<0.00001	−5.70	(−7.81, −3.58)
Vocabulary	0	=0.51	Random	8.66	<0.00001	−4.21	(−5.17, −3.26)
Comprehension	80	=0.03	Random	9.41	<0.0001	−3.34	(−4.94, −1.73)

WMD weighted mean difference, SMD standardized mean difference, 95% CI 95% confidence intervals

highest, and it was followed in descending order by East China, North China, Northwest China, and South China. The water fluoride content in Northeast China was the lowest. This study found that while there were significant regional differences in the fluoride content of drinking water in China (Wu et al., 2021), the water improvement and defluoridation project effectively reduced the fluoride content in drinking water across all regions.

Skeletal fluorosis mostly occurs in young adults (Meena & Gupta, 2021; Srivastava & Flora, 2020). It is a chronic invasive systemic bone disease caused by long-term excessive intake of fluoride and is an important indicator of the effect of water improvement and defluoridation strategies (WHO, 2014; Idowu et al., 2019). The present results showed that the prevalence of skeletal fluorosis and urinary fluoride levels in adults and children decreased significantly after implementation of the water improvement projects in China. The findings of this study indicate that the fluoride content in the human body can be effectively reduced by water improvement and defluoridation strategies.

Dental fluorosis in children showed that water improvement strategies decreased significantly. The results showed that the incidence rate of dental fluorosis in children was higher in the South than that in the North, and this is consistent with the trend in water fluoride content. This implies that the incidence rate of dental fluorosis in children was positively

correlated with the water fluoride content. The decrease in the incidence of dental fluorosis among children was most pronounced in Northeast China. According to Dean's fluorosis index, the findings of the current analysis indicate that the severity of dental fluorosis in children was significantly reduced following the implementation of the water improvement and defluoridation project, with the percentage of children classified as normal increasing significantly. Overall, the water improvement and fluoride reduction project in China has been implemented for more than 40 years and has resulted in a 30% decrease in the incidence of dental fluorosis in children.

Long-term excessive intake of fluoride does not only cause extensive damage to hard tissues such as teeth and bones, but also causes extensive damage to soft tissues to varying degrees (Niu et al., 2018; Quadri et al., 2018; Riddell et al., 2019; Sabine et al., 2020; Zhou et al., 2015). Most studies have reported that the IQ of children with long-term high-fluoride intake was significantly lower than that of children with normal fluoride intake (Dec et al., 2017; Lee et al., 2016; Gong and James 2020). On the contrary, a study conducted in New Zealand concluded that there was no evidence to indicate that fluoride exposure affects the neurologic development or IQ of children (Broadbent et al., 2015). Therefore, there is some controversy about the influence of fluorosis on children's intelligence. Our results showed that the IQ of children in fluorosis areas was lower than

that of children in normal areas. There was a negative correlation between children's IQ and fluoride intake, with the IQ of children in high-fluoride areas lower than that of children in low fluoride areas, IQ of children with dental fluorosis lower than that of children with normal teeth in fluorosis areas, and IQ of children related to the source of fluorosis. The study found that there were significant differences in the IQ scores between normal areas and fluorosis areas, with a significantly higher percentage of children in the fluorosis areas having IQ scores below 89 and a significantly higher percentage of children in normal areas having IQ scores higher than 120. In addition, the IQ scores of children with dental fluorosis were significantly lower than those of children with normal teeth. This has important implications, as this has been a controversial issue in the literature. Thus, these findings further emphasize the impact of fluoride exposure on intelligence levels in children.

So far, thyroid hormone levels have been identified as markers of mental retardation in children exposed to fluoride (Broadbent et al., 2015; Xu et al., 2020). Our results showed that the TSH level of children in fluorosis areas was significantly higher than that in normal areas, and the T4, FT4, and language IQ levels were significantly lower than those in normal areas. This is consistent with the results of Hara (1980).

The provinces this study have included are relatively large and time long, which results in significant heterogeneity in the analysis results. Through sensitivity and subgroup analysis, it was found that heterogeneity had a small impact on the results of this study. This study is the first to systematically analyze the effect of water quality improvement and fluoride reduction strategies in China over the last 40 years. The findings confirm all the reports so far that water improvement and fluoride reduction are effective measures to prevent endemic fluorosis caused by drinking water and can also effectively reduce the impact of fluorosis on children's intelligence. However, the findings also point to some weak links and problems that need to be resolved in this regard. Due to the limited service life of water improvement facilities, poor management, and inadequate health supervision, a considerable number of water improvement and fluoride reduction projects have been discontinued or scrapped altogether. However, it is necessary to continue to invest in these projects in

fluorosis endemic areas and strengthen the management and maintenance of the fluoride reduction and water improvement project, in order to ensure that residents of these areas have access to drinking water with low fluoride content. This will help to effectively control the disease and continue to reap the benefits of the prevention and control measures that have been in place over the last few decades.

Conclusions

This is the first retrospective analysis of the effect of the water improvement and fluoride reduction project in China, which has been implemented over the past 40 years. The findings demonstrate that the water fluoride content, the incidence of dental fluorosis, and the incidence of skeletal fluorosis were effectively reduced as a result of this project. The findings also highlighted obvious regional differences in fluorosis areas, as the water fluoride content and the incidence rate of dental fluorosis were significantly higher in the South than in the North. This implies that efforts for the prevention and control of fluorosis in the South should be further strengthened. The present results also suggest that fluorosis can cause a reduction in children's IQ, and this effect was found to be associated with the type of fluoride exposure. Based on the prevalence of dental fluorosis and its effect on the intelligence of children, it appears that reducing fluoride levels in drinking water and monitoring water quality are important strategies for the prevention and treatment of dental fluorosis.

Author contributions All authors contributed to the study conception and design; they had full access to all data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. FW and YL wrote the report. XZ, DM, YL, and DT critically revised the report. JZ, BY, CZ, MS, and ZH performed the statistical analysis. The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding This study was supported by the National Natural Science Foundation of China (Nos. 82160519, 31660326); the Natural Science Foundation of Guizhou Province (Nos. QianKeHe Support [2022]181, QianKeHe Basics -ZK[2023] Key 042); the Natural Science Foundation of Guiyang City (Nos. [2022]4-3-2, [2022]4-3-10); Research project of Education Department of Guizhou Province (No. QianJiaoJi [2023]037); and Subject Excellent Reserve Talent Project (No. gyfyxkrc-2023-14). The funders of the study had no role in

study design, data collection, data analysis, data interpretation, or writing of the report.

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Conflict of interest The authors declare that they have no known competing financial interests or personal relationships that could have influenced the work reported in this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alexandra, S., Diego, F. R., Manuel, R., et al. (2021). Clinical changes in the severity of dental fluorosis: a longitudinal evaluation. *BMC Oral Health*, 21(1), 366. <https://doi.org/10.1186/s12903-021-01729-3>
- Alvee, S., Rajinder, K. D., Ravi, S. S., et al. (2021). Teaching neuroimage: Fluorosis: A forgotten cause of compressive myelopathy. *Neurology*, 97(19), e1973–e1974. <https://doi.org/10.1212/WNL.00000000000012389>
- Amini, M., Mueller, K., Abbaspour, K. C., et al. (2008). Statistical modeling of global geogenic fluoride contamination in groundwaters. *Environmental Science & Technology*, 42(10), 3662–3668. <https://doi.org/10.1021/es071958y>
- Broadbent, J. M., Thomson, W. M., Ramrakha, S., et al. (2015). Community water fluoridation and intelligence: prospective study in New Zealand. *American Journal of Public Health*, 105(1), 72–76. <https://doi.org/10.2105/AJPH.2013.301857>
- Chang, Z. L., Yang, X. J., Zhao, C. X., et al. (2018). Analysis of monitoring data on drinking-water-borne endemic fluorosis from 2012–2016 in Inner Mongolia Autonomous Region. *Chinese Journal of Endemiology*, 37(06), 485–488.
- Chen, M. S., Guo, X. C., Wang, R. Y., et al. (2000). Effect of altering water sources to lower fluoride level at hot spring fluorosis areas. *Chinese Journal of Endemiology*, 19(04), 47–48.
- Chen, Z. (2003). Analysis on monitoring results of water improvement and fluoride reduction in Nanzhuang village of Shan county for 10 years. *Henan Journal of Preventive Medicine*, 06, 349–350.
- Dec, K., Łukomska, A., Maciejewska, D., et al. (2017). The influence of fluorine on the disturbances of homeostasis in the central nervous system. *Biological Trace Element Research*, 177(2), 224–234. <https://doi.org/10.1007/s12011-016-0871-4>
- Dong, J. M., Zhou, X. M., Wu, T. H., et al. (2011). An analysis of the incidence of dental fluorosis of children aged 8 to 12 years in endemic fluorosis villages that changed water in Lianyungang city. *Chinese Journal of Control of Endemic Diseases*, 26(04), 278–280.
- Dong, L., Yao, P. J., Chen, W., et al. (2018). An investigation of children's dental fluorosis and intelligence in drinking water-type of endemic fluorosis area in Xi'an. *Chinese Journal of Endemiology*, 37(01), 45–48.
- Everett, E. T. (2011). Fluoride's effects on the formation of teeth and bones, and the influence of genetics. *Journal of Dental Research*, 90, 552–560. <https://doi.org/10.1177/0022034510384626>
- Fan, S. L. (2020). Current situation of prevention and control of endemic fluorosis in China during the 13th Five Year Plan Period. *Journal of Occupational and Environmental Medicine*, 37(12), 1219–1223. <https://doi.org/10.13213/j.cnki.jeom.2020.20274>
- Gong, C. X., & James, N. E. I. (2020). Association between maternal fluoride exposure and child IQ. *JAMA Pediatrics*, 174(2), 212–213. <https://doi.org/10.1001/jamapediatrics.2019.5239>
- Guo, S. H., Hou, X. Y., Wang, Y. J., et al. (2015). Analysis of drinking water fluorosis monitoring results in Weinan City in 2015. *Foreign Medical Sciences*, 36(3), 190–192.
- Guo, X. C., Wang, R. Y., Huang, Y. L., et al. (2002). Effect of water improvement to reduce fluoride in drinking-water type fluorosis areas in Hunan province. *Chinese Journal of Endemiology*, 21(05), 49–51.
- Habtamu, D., Abebe, B., Zewdu, A., et al. (2019). Fluoride concentration in ground water and prevalence of dental fluorosis in Ethiopian Rift Valley: Systematic review and meta-analysis. *BMC Public Health*, 19, 1298. <https://doi.org/10.1186/s12889-019-7646-8>
- Hara, K. (1980). Studies on fluorosis especially effects of fluoride on thyroid metabolism. *Shikoku Eisei Gakkai Zasshi*, 30(1), 42–57. <https://doi.org/10.5834/jdh.30.42>
- He, N., Wang, R., Wang, Y. C., et al. (2021). Analysis of monitoring results of drinking water type endemic fluorosis in Fucheng County, Hengshui City. *Chinese Journal of Control of Endemic Diseases*, 36(01), 52–54.
- He, X. M., Gu, J. H., & Lin, H. Z. (2011). Monitoring and analysis of drinking water type endemic fluorosis of central transfer payment project in Xianyou County from 2008 to 2009. *Strait Journal of Preventive Medicine*, 17(04), 60–61.
- Hong, F. G., Wang, H., Yang, D., et al. (2001). A study of fluoride effects on children's intelligence development under different environments. *Chinese Primary Health Care*, 15(01), 12–14.
- Huang, W. Y., Huang, K., Xu, Z. Q., et al. (2016). Effects of water improvement projects on children's dental fluorosis in drinking water-born endemic fluorosis areas in

- Yangchun City, Guangdong. *Modern Preventive Medicine*, 43(21), 3869–3879.
- Idowu, O. S., Azevedo, L. B., & Valentine, R. A. (2019). The use of urinary fluoride excretion to facilitate monitoring fluoride intake: A systematic scoping review. *PLoS ONE*, 14(9), e0222260. <https://doi.org/10.1371/journal.pone.0222260>
- James, P., Harding, M., Beecher, T., et al. (2021). Impact of reducing water fluoride on dental caries and fluorosis. *Journal of Dental Research*, 100(5), 507–514. <https://doi.org/10.1177/0022034520978777>
- Jia, L. H., Ma, J., Du, Y. G., et al. (2011). The investigation of drinking-water-borne endemic fluorosis in Hebei province in 2009. *Chinese Journal of Endemiology*, 30(02), 184–187.
- Jian, H. H., Chen, Q. W., Zhang, Y., et al. (2013). Analysis of monitoring results of drinking water type endemic fluorosis in Fuyang city in 2012. *Anhui Journal of Preventive Medicine*, 19(04), 251–253.
- Jin, X., Cui, Y. S., & Cao, L. C. (2020). The relationship between thyroid hormone abnormality and DIO1 gene polymorphism in children with dental fluorosis. *Tianjin Medical Journal*, 37(08), 717–721.
- Julvez, J., López-Vicente, M., Warembourg, C., et al. (2021). Early life multiple exposures and child cognitive function: A multi-centric birth cohort study in six European countries. *Environmental Pollution*, 284, 117404. <https://doi.org/10.1016/j.envpol.2021.117404>
- Kan, Z. Y., Zheng, Z. X., Wang, J. H., et al. (2021). Investigation on control and prevention of drinking water endemic fluorosis in Liaoning. *China Tropical Medicine*, 21(11), 1032–1046. <https://doi.org/10.13604/j.cnki.46-1064/r.2021.11.03>
- Kloos, H., & Haimanot, R. T. (1999). Distribution of fluoride and fluorosis in Ethiopia and prospects for control. *Tropical Medicine & International Health*, 4(5), 355–364. <https://doi.org/10.1046/j.1365-3156.1999.00405.x>
- Kou, B. Z. (2015). Analysis of surveillance results of endemic fluorosis in Baicheng City from 2002 to 2007. *China Health Industry*, 12(11), 9–10.
- Kumar, H., Boban, M., & Tiwari, M. (2009). Skeletal fluorosis causing high cervical myelopathy. *Journal of Clinical Neuroscience*, 16, 828–830. <https://doi.org/10.1016/j.jocn.2008.08.028>
- Lee, J., Han, Y. E., Favorov, O., et al. (2016). Fluoride induces a volume reduction in CA1 hippocampal slices via MAP kinase pathway through volume regulated anion channels. *Experimental Neurobiology*, 25(2), 72–78. <https://doi.org/10.5607/en.2016.25.2.72>
- Li, F. H., Chen, X., Huang, R. J., et al. (2009). Intelligence impact of children with endemic fluorosis caused by fluoride from coal burning. *Journal of Environmental Health*, 26(4), 338–340.
- Li, M. J., Qu, X. N., Hong, M., et al. (2020a). Spatial distribution of endemic fluorosis caused by drinking water in a high-fluorine area in Ningxia, China. *Environmental Science and Pollution Research International*, 27(16), 20281–20291. <https://doi.org/10.1007/s11356-020-08451-7>
- Li, P. F., Wang, Z. H., Wu, M., et al. (2014). Monitoring and analysis of drinking water type endemic fluorosis in Shanxi Province in 2012. *Bulletin Disease Control and Prevention*, 29(01), 16–32. <https://doi.org/10.13215/j.cnki.jbyfkztb.1311012>
- Li, Q. R., Shen, J. Q., Qin, T., et al. (2021). A qualitative and comprehensive analysis of caries susceptibility for dental fluorosis patients. *Antibiotics (basel)*, 10(9), 1047. <https://doi.org/10.3390/antibiotics10091047>
- Li, X. J., Jiao, B. R., & Li, Y. S. (2020b). Condition survey and analysis of relevant factors on 12–15 years old youths with dental fluorosis after water-improving and defluoridation in Jiashi general farm of the third agricultural division of Xinjiang. *Chinese Journal of Aesthetic Medicine*, 29(06), 142–146.
- Li, X. L., & Zhang, F. L. (2016). Effects of fluorosis on intelligence, physical development and serum bone morphogenetic protein levels in children. *Chinese Journal of Control of Endemic Diseases*, 31(08), 895–896.
- Li, X. S., Zhi, J. L., & Gao, R. O. (1995). Effect of fluoride exposure on intelligence in children. *Fluoride*, 28(4), 189–192.
- Li, Y. H., Li, Z. L., Zhang, X. Q., et al. (2017). Analysis of effect of water improvement projects in drinking water type fluorosis areas in Inner Mongolia. *Journal of Medical Pest Control*, 33(01), 12–14.
- Li, Y. P., Quan, X. Y., Chen, D., et al. (2003). Effect of endemic fluorosis on children's intelligence development in Baotou City. *Chinese Journal PHM*, 19(4), 337–338.
- Li, Y., Wang, F., Feng, J., et al. (2020c). Health risk in children to fluoride exposure in a typical endemic fluorosis area on Loess Plateau, north China, in the last decade. *Chemosphere*, 243, 125451. <https://doi.org/10.1016/j.chemosphere.2019.125451>
- Liu, S. S., Lv, Y., Sen, Z. S., et al. (2000). Investigation on intelligence level of children in high fluoride area. *Chinese Journal of Control of Endemic Diseases*, 15(4), 231–232.
- Liu, X. L., Fan, Z. X., Hua, J. L., et al. (1999). Determination of T3, T4 and TSH in children in endemic fluorosis area. *Bulletin of Endemic Diseases*, 14(01), 16–17.
- Liu, Z. H., Li, W. F., Cui, Y. H., et al. (2019). Effects of different ways and years of waterimproving on children with dental fluorosis in Tianjin. *Chinese Journal of Endemiology*, 38(01), 36–40.
- Luo, C. (2018). *Correlation analysis between fluoride exposure and intelligence quotients as well as mitochondrial DNA copy number among children*. Master's degree thesis of Huazhong University of Science and Technology.
- Luo, Y., Ma, R. R., Liu, Z. L., et al. (2018). Investigation of children's intelligence in coal-burning fluorosis areas and the significance of Forensic Medicine. *Chinese Journal of Forensic Medicine*, 33(06), 590–593. <https://doi.org/10.13618/j.issn.1001-5728.2018.06.007>
- Lv, F. Q., Guo, L. L., He, C. S., et al. (2014). Analysis of monitoring results of drinking water type endemic fluorosis after water improvement and fluoride reduction in Pingliang City from 2004 to 2013. *Bulletin Disease Control and Prevention*, 29(01), 18–20. <https://doi.org/10.13215/j.cnki.jbyfkztb.1310010>
- Ma, G. Q. (2010). Analysis of monitoring results of drinking-water fluorosis in Anyang city in 2009. *China Healthcare Nutrition Volume (academic Edition)*, 10(219), 12–130.

- Ma, Q. (2019). *Screening and validation of NNAT gene methylation marker related to intellectual development in Children's exposed to fluoride*. Zhengzhou University for the degree of Master.
- Malin, A. J., Lesseur, C., Busgang, S. A., et al. (2019). Fluoride exposure and kidney and liver function among adolescents in the United States: NHANES, 2013–2016. *Environment International*, 132, 105012. <https://doi.org/10.1016/j.envint.2019.105012>
- Meena, L., & Gupta, R. (2021). Skeletal fluorosis. *The New England Journal of Medicine*, 385(16), 1510. <https://doi.org/10.1056/NEJMcim2103503>
- Mi, P. (2019). *Effects of High fluoride on Children's intelligence and programmed death in neurotoxicity in rats*. Master's Thesis of Tianjin Medical University.
- Mohd Nor, N. A., Chadwick, B. L., Farnell, D. J. J., et al. (2021). Factors associated with dental fluorosis among Malaysian children exposed to different fluoride concentrations in the public water supply. *Journal of Public Health Dentistry*, 81(4), 270–279. <https://doi.org/10.1111/jphd.12448>
- Murray, M. (1937). Fluorine intoxication. *Nature*, 140, 483–484. <https://doi.org/10.1038/140483a0>
- Ni, N., Wu, C. F., Li, C. H., et al. (2020). Analysis of dental fluorosis in 2393 school aged children. *Chinese Journal of Control of Endemic Diseases*, 35(05), 568–569.
- Niu, R., Chen, H., Manthari, R. K., et al. (2018). Effects of fluoride on synapse morphology and myelin damage in mouse hippocampus. *Chemosphere*, 194, 628–633. <https://doi.org/10.1016/j.chemosphere.2017.12.027>
- Pang, W. L., Ge, J. H., & Xu, C. F. (2011). Analysis on the effect of improving water quality, reducing fluoride and preventing endemic fluorosis in Tiantai County. *Zhejiang Preventive Medicine*, 23(04), 38–39.
- Pu, G. L., Lu, Q., Yang, P. Z., et al. (2019). Analysis of monitoring results of dental fluorosis in children with drinking water fluorosis in Qinghai Province from 2009 to 2017. *Chinese Journal of Endemiology*, 38(07), 562–565.
- Quadri, J. A., Sarwar, S., Sinha, A., et al. (2018). Fluoride-associated ultrastructural changes and apoptosis in human renal tubule: A pilot study. *Human & Experimental Toxicology*, 37, 1199–1206. <https://doi.org/10.1177/0960327118755257>
- Riddell, J. K., Malin, A. J., Flora, D., et al. (2019). Association of water fluoride and urinary fluoride concentrations with attention deficit hyperactivity disorder in Canadian youth. *Environment International*, 133(1), 105190. <https://doi.org/10.1016/j.envint.2019.105190>
- Sabine, G., Stephanie, H., Angelika, R., et al. (2020). Toxicity of fluoride: Critical evaluation of evidence for human developmental neurotoxicity in epidemiological studies, animal experiments and in vitro analyses. *Archives of Toxicology*, 94(5), 1375–1415. <https://doi.org/10.1007/s00204-020-02725-2>
- Saeed, M., Malik, R., & Kamal, A. (2020). Fluorosis and cognitive development among children (6–14 years of age) in the endemic areas of the world: A review and critical analysis. *Environmental Science and Pollution Research International*, 27(3), 2566–2579. <https://doi.org/10.1007/s11356-019-06938-6>
- Seraj, B., Shahrabi, M., Shadfar, M., et al. (2012). Effect of high water fluoride concentration on the intellectual development of Children in Makoo/Iran. *Journal of Dentistry (tehran)*, 9(3), 221–229.
- Shashi, A., Kumar, M., & Bhardwaj, M. (2008). Incidence of skeletal deformities in endemic fluorosis. *Tropical Doctor*, 38, 231–233. <https://doi.org/10.1258/td.2008.070379>
- Shi, M. L., Yao, M. Q., Cui, Y. X., et al. (2011). Monitoring and evaluation of the defluoridation effect by water improvement in shihezi area. *Modern Preventive Medicine*, 38(21), 4355–4357.
- Shu, C. L., Wang, C. S., Wang, Y., et al. (2013). Analysis of monitoring results of drinking water endemic fluorosis in Jiangsu Province in 2009. *Chinese Journal of Endemiology*, 32(06), 662–667.
- Srivastava, S., & Flora, S. J. S. (2020). Fluoride in drinking water and skeletal fluorosis: A Review of the global impact. *Current Environmental Health Reports*, 7(2), 140–146. <https://doi.org/10.1007/s40572-020-00270-9>
- Sun, G. D., Zhang, X., Su, Y. N., et al. (2011). Investigation and analysis on the effect of water improvement and fluoride reduction in five counties of Shandong Province in 2009. *Journal of Preventive Medicine and Public Health*, 22(04), 112–113.
- Theobald, S., Brandes, N., Gyapong, M., et al. (2018). Implementation research: new imperatives and opportunities in global health. *Lancet*, 392(10160), 2214–2228. [https://doi.org/10.1016/S0140-6736\(18\)32205-0](https://doi.org/10.1016/S0140-6736(18)32205-0)
- Wang, R. (2012). *Joint effects of excessive fluoride and iodine in drinking water on children's thyroid and intelligence development*. Master's Thesis of Tianjin Medical University.
- Wang, C., Gao, Y. H., Wang, W., et al. (2012a). A national cross-sectional study on effects of fluoride-safe water supply on the prevalence of fluorosis in China. *British Medical Journal Open*, 2(5), e001564. <https://doi.org/10.1136/bmjopen-2012-001564>
- Wang, F. Q., Li, Y. J., Tang, D. X., et al. (2021a). Effects of water improvement and defluoridation on fluorosis-endemic areas in China: A meta-analysis. *Environmental Pollution*, 270, 116227. <https://doi.org/10.1016/j.envpol.2020.116227>
- Wang, G. J., Zhang, M. F., Wang, Q. S., et al. (2017). The correlation between the level of serum fluoride and intelligence quotient among children before and after defluoridation. *Capital Journal of Public Health*, 11(06), 274–277.
- Wang, J. H., Zheng, Z. X., Liu, W., et al. (2008). Endemic fluorosis: Prevalence and prevention in Liaoning Province. *Chinese Journal of Endemiology*, 27(06), 663–667.
- Wang, J. X., Yue, B. J., Zhang, X. H., et al. (2021b). Effect of exercise on microglial activation and transcriptome of hippocampus in fluorosis mice. *Science of the Total Environment*, 760, 143376. <https://doi.org/10.1016/j.scitotenv.2020.143376>
- Wang, M., Liu, L., Li, H., Li, Y., Liu, H., Hou, C., Zeng, Q., Li, P., Zhao, Q., Dong, L., & Zhou, G. (2020). Thyroid function, intelligence, and low-moderate fluoride exposure among Chinese school-age children. *Environment International*, 134, 105229. <https://doi.org/10.1016/j.envint.2019.105229>

- Wang, Q. J., Gao, M. X., Zhang, M. F., et al. (2012b). Study on the correlation between daily total fluoride intake and children's intelligence quotient. *Journal of Southeast University (medical Science Edition)*, 31(06), 743–746.
- Wang, R., He, N., Wang, Y. C., et al. (2021c). Investigation and analysis on children's dental fluorosis and IQ level in high-fluoride areas of Hengshui City. *Journal of Medical Pest Control*, 37(08), 796–800.
- Wang, S. Q. (2021). Analysis on prevention and control status of drinking water borne fluorosis, Zhoucun district Zibo city. *Preventive Medicine Tribune*, 27(07), 551–554. <https://doi.org/10.16406/j.pmt.issn.1672-9153.2021.07.022>
- Wang, S. X., Wang, Z. H., Cheng, X. T., et al. (2005). Investigation and evaluation on intelligence and growth of children in endemic fluorosis and arsenism areas. *Chinese Journal of Endemiology*, 24(2), 179–182.
- Wang, S. X., Wang, Z. H., Cheng, X. T., et al. (2007). Arsenic and fluoride exposure in drinking water: Children's IQ and growth in Shan yin county, Shanxi province, China. *Environmental Health Perspectives*, 115(4), 643–647.
- Wang, S., & Zhu, X. H. (2012). Investigation and analysis on intelligence level of children in high fluoride area. *Chinese Journal of Control of Endemic Diseases*, 27(01), 67–68.
- Wang, Z. H., Wang, S. X., Zhang, X. D., et al. (2006). Investigation on children's growth and development under long term fluoride exposure. *Chinese Journal of Control of Endemic Diseases*, 21(4), 239–241.
- Wei, W., Pang, S. J., & Sun, D. J. (2019). The pathogenesis of endemic fluorosis: Research progress in the last 5 years. *Journal of Cellular and Molecular Medicine*, 23(4), 2333–2342. <https://doi.org/10.1111/jcmm.14185>
- WHO (World Health Organization). (2014). *Basic methods for assessment of renal fluoride excretion in community prevention programmes for oral health*. WHO, Geneva.
- Wu, L. W., Fan, C. L., Zhang, Z. H., et al. (2021). Association between fluoride exposure and kidney function in adults: A cross-sectional study based on endemic fluorosis area in China. *Ecotoxicology and Environmental Safety*, 225, 112735. <https://doi.org/10.1016/j.ecoenv.2021.112735>
- Xiang, J., Tu, Q. Y., Chen, C., et al. (2019). Effect of high fluoride intake on TT3, TT4 and TSH levels in children. *Jiangsu Journal of Preventive Medicine*, 30(04), 383–392.
- Xiang, Q., Liang, Y., Chen, L., et al. (2003). Effect of fluoride in drinking water on children's intelligence. *Fluoride*, 36(02), 84–94.
- Xu, K. H., An, N., Huang, H., et al. (2020). Fluoride exposure and intelligence in school-age children: Evidence from different windows of exposure susceptibility. *BMC Public Health*, 20, 1657. <https://doi.org/10.1186/s12889-020-09765-4>
- Xu, X. H. (2005). Surveillance of prevention and control of endemic fluorosis by water-improving and defluoridation in Liangshan County for 20 years. *Chinese Medicine and Health*, 17(02), 63–64.
- Yang, S. Y., Deng, Y., Yao, L. M., et al. (1997). Comparative analysis of physical and mental development of children in fluorosis areas with and without water improvement. *Literature and Information on Preventive Medicine*, 3(1), 42–43.
- Yao, L. M., Zhou, J. L., Wang, X. L., et al. (1996). TSH and intelligence level of children with dental fluorosis in high fluoride area. *Literature and Information on Preventive Medicine*, 2(1), 26–27.
- Yu, J., Chen, S. X., Li, B. L., et al. (2004). Evaluation of the effect of water- improving in high-fluoride districts in Guangzhou, Huizhou, and Chaozhou. *Chinese Rural Health Service Administration*, 24(03), 35–37.
- Yu, L. P., Zhang, X. L., Zhang, S. W., et al. (2021). Analysis of trace elements calcium and zinc in hair of children in fluorosis and non fluorosis areas. *Chinese Journal of Control of Endemic Diseases*, 36(04), 360–362.
- Yu, L. Y., Liu, H. L., Wang, R., et al. (2014). Joint effects of excessive fluoride and iodine on thyroid function among children in Tianjin. *Chinese Journal of Public Health*, 30(02), 212–215.
- Yu, Q. L., Yang, Y., Chun, Z. M., et al. (2008). Control effect of endemic fluorosis in Dunhuang City on water-improving and defluoridation project. *Occupational Health*, 24(06), 566–567.
- Yu, X., Chen, J., Li, Y., et al. (2018). Threshold effects of moderately excessive fluoride exposure on children's health: A potential association between dental fluorosis and loss of excellent intelligence. *Environment International*, 118, 116–124. <https://doi.org/10.1016/j.envint.2018.05.042>
- Zhang, J. J. (2002). Analysis of the effect of altering water resources to lower fluoride level in Ma Chang Jing Village Etuoke Qian Qi Inner Mongolia Autonomus Region. *Inner Mongolia Medical Journal*, 34(05), 427–429.
- Zhang, J. W., Yao, H., & Chen, Y. (1998). Investigation on the effect of high arsenic and high fluoride on intelligence development of offspring. *Chinese Journal of Public Health*, 17(02), 57–58.
- Zhang, M. (2010). Analysis of key surveillance results of endemic fluorosis in Hebei Province. *Hebei Medicine*, 16(12), 1522–1523.
- Zhang, N. (2018). *Investigation on the effect of reducing fluoride and improving drinking water on the prevalence of dental fluorosis in children in drinking water fluorosis areas of Anshan City*. Master's Degree Thesis of China Medical University.
- Zhang, P. H., & Cheng, L. (2015). Effect of coal-burning endemic fluorosis on children's physical development and intellectual level. *Chinese Journal of Control of Endemic Diseases*, 30(06), 458–459.
- Zhang, S., Zhang, X., Liu, H., et al. (2015). Modifying effect of COMT gene polymorphism and a predictive role for proteomics analysis in children's intelligence in endemic fluorosis area in Tianjin, China. *Toxicological Sciences: an Official Journal of the Society of Toxicology*, 144(2), 238–245. <https://doi.org/10.1093/toxsci/kfu311>
- Zhang, X. F. (2012a). *Studies of relationships between the polymorphism of COMT gene and plasma proteomic profiling and children's intelligence in high fluoride areas*. Master's Degree Thesis of Huazhong University of Science and Technology.
- Zhang, X. F. (2012b). *Study on the relationship between COMT gene polymorphism, plasma proteomics and intelligence of children in high fluoride areas*. Master's

- Degree Thesis of Huazhong University of Science and Technology.
- Zhao, L. J., Pei, J. R., Zhang, W., et al. (2016). Surveillance on drinking-water-born endemic fluorosis in China, 2013. *Zhonghua Liu Xing Bing Xue Za Zhi*, 37(6), 816–820.
- Zhao, Y., Zhang, C. N., Kang, J., et al. (2015). Monitoring and analysis of endemic fluorosis of drinking water type in Heilongjiang province in 2014. *Chinese Journal of PHM*, 31(1), 84–85. <https://doi.org/10.19568/j.cnki.23-1318.2015.01.036>
- Zhao, Y. N., Zhou, J. C., & Li, S. L. (2012). Investigation on the effect of water control in endemic fluorosis areas in Qingtongxia City. *Chinese Journal of School Health*, 33(09), 1108–1110. <https://doi.org/10.16241/j.cnki.1001-5914.2012.05.036>
- Zheng, Z. G., Chen, Y., & Ling, S. Q. (2010). Investigation on prevention and treatment of drinking water fluorosis in Bobai County. *Journal of Applied Preventive Medicine*, 16(02), 101–102.
- Zhi, D., & Wang, L. Y. (2020). Effect of water improvement and fluoride reduction on dental fluorosis in school-age children in Shenyang and its risk factors. *Chinese Journal of Public Health Engineering*, 19(01), 57–59.
- Zhou, B. H., Zhao, J., Liu, J., et al. (2015). Fluoride-induced oxidative stress is involved in the morphological damage and dysfunction of liver in female mice. *Chemosphere*, 139, 504–511. <https://doi.org/10.1016/j.chemosphere.2015.08.030>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.