

# Sources of Carbon for Hepatic Glycogen Synthesis in the Conscious Dog

Doss W. Neal,\* Christine Badet,<sup>§</sup> and Gerald I. Shulman\*

\*Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, Tennessee 37222;

way contribution (see Discussion). The weights of the animals ranged from 18.7 to 23.9 kg (mean $\pm$ SEM 21.2 $\pm$ 0.8 kg). They were maintained

cess took no more than 10 min from the time of sacrifice. The position of the catheter tips was then verified to ensure proper placement.

Diet No. 5006, Ralston Purina Co., St. Louis, MO) composed of 34% after a 42-h fast, and liver samples were taken just as in the experimen-

blood glucose values per se, but the variance was reduced because of the accuracy of plasma glucose A-V differences, which do not require a denroteinization step. The use of whole blood glucose ensures accurate

Results

Blood glucose concentrations: hepatic glucose load balance


cose entry into the erythrocyte.

portal vein whole-blood glucose levels, arterial and portal vein

Intraduodenal Glucose Infusion

Mean net hepatic lactate uptake during the control period was





Glycerol →

within the glucose space, leaving 44% of the net absorbed glucose, or 13 g. to be disposed of by nonhepatic tissues. If the

*Table II. Comparison of Net Hepatic Balance of Gluconeogenic Substrates with Net Hepatic Glcogen Synthesis and NMR-determined*

*Contribution of the Direct Pathway during the Glucose Infusion Period*

<sup>13</sup>C-NMR and GC-MS data

Dog no.	carbon*	lactate	C <sub>1</sub>	C <sub>6</sub>	PV glucose	CF	pathway	synthesis
---------	---------	---------	----------------	----------------	------------	----	---------	-----------

synthesis are not clear. Although there were no correlations

A second possibility is that lactate carbon was derived from

between hormonal levels and changes and the pathways of glycogen synthesis, it is notable that dog 3, which had an unusually large mass of glycogen deposited via the direct pathway.

nonhepatic tissues. This implies that there was uptake of lactate by the liver, even though there was net lactate output, suggesting that the total hepatic lactate production was greater

experienced an increase in plasma insulin concentration (to a

than the net production. Net balance measurements underesti-

level that was fourfold the mean of the other dogs) during the

mate absolute uptake or production of substrates which are

final hour of the glucose infusion.

both taken up and released by an organ, since they reflect the

The current data are consistent with the hypothesis (33)

quantitatively dominant process. Thus, we cannot rule out this

that the three-carbon substrates used for glycogen synthesis by

explanation of our findings. The lactate being taken up by the



case in our animals. Nevertheless, these investigators con-

sidered the effect of galactose on carbohydrate and substrate metabolism in healthy man. *Metab. Clin.*

34. Cahill, G. F., Jr., A. B. Hastings, J. Ashmore, and S. Zottu. 1958. Studies

40. Johnson, J. L., and G. J. Bagby. 1988. Gluconeogenic pathway in liver and

on carbohydrate metabolism in rat liver slices. X. Factors in the regulation of  
pathways of glucose metabolism. *J. Biol. Chem.* 230:125-135.

muscle glycogen synthesis after exercise. *J. Appl. Physiol.* 64:1591-1599.

41. Marin, P., M. Rebuffe-Scrive, U. Smith, and P. Bjorntorp. 1987. Glucose