# Design, Synthesis, Molecular Modeling, *In Silico* ADME Studies and Anti-HIV-1 Assay of New Diazocoumarin Derivatives

Zeynab Alimi Livani , Mahdieh Safakish , Zahra Hajimahdi , Sepehr Soleymani , Rezvan Zabihollahi , Mohammad Reza Aghasadeghi , Eskandanr Alipour and Afshin Zarghi \*

#### Abstract

Some new diazo incorporated coumarin compounds were designed and synthesized to evaluate their anti-HIV activity. Overall, compounds were active against HIV at 100  $\mu$ M. Additionally, no cytotoxic effect was observed at this concentration. The compound with 4-chlorobenzyl group indicated the best anti-HIV activity (52%). Docking studies using the later crystallographic data available for PFV integrase showed similar binding modes to HIV-1 integrase inhibitors. On the basis of these data, nitrogen atoms of 1,3,4-oxadiazole ring have been involved in the Mg<sup>2+</sup> chelation and 4-chlorobenzyl group occupies the same position as 4-flourobenzyl group of raltegravir in the active site. In addition, ADME assay demonstrated favorable physicochemical properties for the new designed compounds. Thus, synthesized structures could be introduced as a novel template for designing safe anti-HIV compounds with integrase inhibitory potential.

Keywords: Anti-HIV; Diazocoumarin; 1,3,4-Oxadiazole; Synthesis, Docking, ADME.

### Introduction

Immunodeficiency Acquired Syndrome (AIDS) is the second cause of death from infectious diseases in low income countries (1). Human Immunodeficiency Virus (HIV) is recognized as the infective agent. Current anti-HIV drugs target different stages of HIV life cycle including absorption, fusion, reverse transcription, integration, and proteolytic cleavage (2-4). Although there is a plethora of anti-HIV drugs, drug toxicity, viral resistance, and patient incompliance has been limited antiHIV therapy (5). Altogether, there is a demand for novel drugs from alternative scaffolds. HIV-1 integrase is an essential enzyme which catalyzes the integration of viral DNA into the host cellular DNA. This process is separated into 2 steps. First, 3'-processing resulting in the creation of free hydroxyl groups at reversetranscribed proviral cDNA 3'-ends. During the second step (strand transfer), integrase catalyzes the insertion of the 3'-processed viral DNA into the host cell genome (6, 7). FDAapproved integrase inhibitors (raltegravir (8, 9), elvitegravir (10, 11) and dolutegravir (12, 13) in Figure 1) compete with host DNA in the IN active site, thus specifically inhibiting the strand transfer step (14). Development of integrase

<sup>\*</sup> Corresponding author:

E-mail: zarghi@sbmu.ac.ir



Dolutegravir

Figure 1. FDA-approved INSTIs.

strand transfer inhibitors (INSTIs) represents an attractive research area in the field of HIV-1 drug discovery because there is no host cell equivalent for HIV-1 integrase enzyme (15).

The integrase strand transfer inhibitors (INSTIs) bind in the interface of the viral DNA and catalytic DDE motif. For this reason, they could be classified as interfacial inhibitors. The DDE motif contains a triad of three conserved acidic residues (Asp-64, Asp-116, and Glu-152) coordinating 2 Mg<sup>2+</sup> cofactors which are critical for the catalytic activity (16). There are various chemical classes of INSTIs such as oligonucleotides, peptides, hydroxylated aromatics, hydrazides, quinoline, and quinolones. All specific INSTIs feature a metal chelating motif that complexes 2 Mg<sup>2+</sup> cofactors and a hydrophobic terminal benzyl moiety (17-19).

Coumarin is a well-known drug scaffold which displays versatile biological activities especially substantial anti-HIV activity (20). Notably, IN inhibitory activity has been reported for coumarin dimmers (1, Figure 2) (21) and 4-hydroxy-5-azacoumarin-3-carbox(thio)amides (2, Figure 2) (22). Furthermore, the anti-HIV activity of a bis-azo compound (FP-21399, Figure 2) has been evidenced (23). With regard to these data we selected coumarin scaffold as a core moiety to design new antiHIV compounds. The decision for selecting substitutions on coumarin ring was carried out from the information obtained through our previous studies. Recently, we reported development of 8-phenylquinolin-4(1)-one derivatives possessing 1,3,4-oxadiazole ring as metal-chelating moiety along with a benzyl part necessary for HIV-1 IN inhibitory activity. Among them, compound 3 (Figure 2) displayed promising anti-HIV-1 activity with  $EC_{50} = 50$  $\mu$ M and no considerable cytotoxicity ( $CC_{50}$  > 100  $\mu$ M) (24). SAR analysis showed that the incorporation of electronegative substitutions like halogens (-F, -Cl) at proper positions of benzyl ring would improve inhibitory activity (25). In this study, we merged coumarin core with a 5-(halo-substituted) benzylthio-1,3,4oxadiazole motif through a diazo-phenyl linker (Designed structure, Figure 2) to present new structure for anti-HIV activity. Some compounds containing alkyl substitution such as methyl, ethyl, and propyl instead of benzyl ring at C-5 position of 1,3,4-oxadiazol ring were also designed. Using wide range of substitutions would help us to perform a SAR analysis on this series.

The overall shape and functionalities of designed structure particularly resemble those of compounds 1 and 3, suggesting that our



Figure 2. Lead anti-HIV compounds (1, 2, 3 and FP-21399) and our designed structure.

novel designed compounds possess ability to be promising anti-HIV agents.

## **Experimental**

3-((4-(5-(substitutedaryl/alkyl)-1,3,4oxadiazol-2-yl) phenyl) diazenyl)-4hydroxy-2 -chromen-2-one derivatives were synthesized according to the scheme 1. First, Ethyl 4-aminobenzoate was prepared from 4-aminobenzoic acid (4) and ethanol in a routine Fischer esterification reaction under sulfuric acid catalysis. Ethyl 4-aminobenzoate (5) was converted to 4-aminobenzohydrazide (6) by refluxing in absolute ethanol and excess of hydrazine hydrate. The 2-mercaptho-1,3,4-oxadiazole ring (7) was closed on the carbohydrazide intermediate in the presence of carbon disulfide and alcoholic potash according to procedure described in references (26, 27). S-alkylation/benzylation reaction was performed with alkyl/ substituted benzyl halides in methanol as the solvent and 10% sodium hydroxide solution to afford derivatives **8a-h.** The moderate basicity of the reaction medium is essential to gain the mono S-alkyl/ benzyl product rather than both S- and N-alkyl/ benzyl substituted product. The diazonium salts of compounds (**9a-h**) were prepared in 6 M hydrochloric acid and 10% sodium nitrite aqueous solution at 0-5 °C. The final derivatives **11a-h** were obtained through coupling reaction of the diazonium salts with 4-hydroxycoumarin (**10**) in aqueous sodium carbonate solution at neutral pH.

All reagents purchased from the Aldrich (USA) or Merck (Germany) Chemical Company and were used without further purifications.

Melting points (mp) were determined using a Thomas Hoover capillary apparatus (Philadelphia, USA). Infrared spectra were acquired on a Perkin-Elmer 1420 ratio recording spectrometer. A Bruker FT-500 MHz instrument (Brucker Biosciences, USA) was used to acquire <sup>1</sup>HNMR; Chloroform-D used as solvent.



**Scheme 1.** Reagents and conditions: (a) ethanol, sulfuric acid, reflux, 24h (b) ethanol,  $NH_2$ - $NH_2$ -OH (6 eq.), reflux, 96h (c) ethanol,  $CS_2$ , KOH, reflux, 4h (d) methanol, 10% aq. NaOH, substituted alkyl/aryl halides, r.t (e) 1. 6M hydrochloric acid 2. 10% sodium nitrite, 0-5 °C (f) 10% aq. Na, CO<sub>3</sub>, stir, 0-5 °C.

Coupling constant () values are estimated in hertz (Hz) and spin multiples are given as brs (broad singlet), s (singlet), d (double), t (triplet), q (quartet), m (multiplet), and br (broad). The mass spectral measurements were performed on an 6410 Agilent LCMS triple quadrupole mass spectrometer (LCMS) with an electrospray ionization (ESI) interface.

# 5

Yield 74%; White crystalline powder; mp 89.2-91.0 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 3424-3348 (NH<sub>2</sub>), 1682 (C=O); LC-MS (ESI) : 166.1 (M+1, 100).

#### 6

Yield 82%; White crystalline powder; mp 216.8-218.3 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 3432-3350 (NH<sub>2</sub>), 3307-3278 (-CO-NH), 3239 (-CO-NH<sub>2</sub>), 1603 (C=O); LC-MS (ESI) : 152.0 (M+1, 100).

### 7

To a solution of compound **6** (6 mmol) and potassium hydroxide (1.23 g) in absolute ethanol (50 mL), carbon disulfide (8.2 mL) was added in an ice bath. The mixture was refluxed at 50-60 °C for 4 hours. Then, it was cooled to room temperature and the solvent was evaporated to dryness. The crude suspended in 7 mL water and acidified with 3 M hydrochloric acid until pH 4.5. The yellow precipitate was filtered and washed with water. Yield 73%; Yellow powder; mp 197-199 °C; IR (KBr disk): v (cm<sup>-1</sup>) 3356-3440 (NH<sub>2</sub>); LC-MS (ESI) : 194.0 (M+1, 100).

#### 8a-h

To a solution of compound 7 (1.55 mmol) in methanol (4 mL) sodium hydroxide (10%, 0.12 mL) was added. The mixture was cooled in

an ice bath and substituted benzyl/alkyl halide (1.55 mmol) was added dropwise and stirred at room temperature. After completion of the reaction, the precipitate was filtered, rinsed with water, and crystallized from ethanol.

### 8*a*

Yield 87%; Yellow crystalline powder; mp 105.8-107.6 °C; IR (KBr disk): v (cm<sup>-1</sup>) 3327-3425 (NH<sub>2</sub>); LC-MS (ESI) : 207.9 (M+1, 100).

### **8**b

Yield 73%; Yellow crystalline powder; mp 80.4-83.2 °C; IR (KBr disk): υ (cm<sup>-1</sup>) 3374-3444 (NH<sub>2</sub>); LC-MS (ESI) : 221.9 (M+1, 100).

### 8*c*

Yield 67%; Yellow crystalline powder; mp 105.1-106.8 °C; IR (KBr disk): v (cm<sup>-1</sup>) 3333-3478 (NH<sub>2</sub>); LC-MS (ESI) : 236.9 (M+1, 100).

### 8*d*

Yield 61%; Yellow crystalline powder; mp 120.8-124 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 3359-3493 (NH<sub>2</sub>); LC-MS (ESI) : 284.9 (M+1, 100).

# 8e

Yield 59%; Yellow crystalline powder; mp 117.8-118.9 °C; IR (KBr disk): v (cm<sup>-1</sup>) 3337-3399 (NH<sub>2</sub>); LC-MS (ESI) : 318.9 (M+1, 100).

# 8f

Yield 65%; Yellow crystalline powder; mp 141.8-144.3 °C; IR (KBr disk): v (cm<sup>-1</sup>) 3349-3417 (NH<sub>2</sub>); LC-MS (ESI) : 318.9 (M+1, 100).

# 8g

Yield 55%; Yellow crystalline powder; mp

121.8-123.8 °C; IR (KBr disk): υ (cm<sup>-1</sup>) 3373-3506 (NH<sub>2</sub>); LC-MS (ESI) : 302.9 (M+1, 100).

# 8*h*

Yield 59%; Yellow crystalline powder; mp 104-106 °C; IR (KBr disk): υ (cm<sup>-1</sup>) 3353-3422 (NH<sub>2</sub>); LC-MS (ESI) : 302.9 (M+1, 100).

### 11a-h

To a cooled solution of substituted compounds **8a-h** (1 mmol) in 6 M hydrochloric acid (2.7 mL), 0.7 mL cooled solution of 10% sodium nitrite was added dropwise in order to keep the temperature less than 5 °C. Then, the

7.3 Hz, H<sub>7</sub>), 7.78 (2H, d, = 8.5 Hz, phenyl H<sub>2</sub> & H<sub>6</sub>), 8.09-8.14 (3H, m, H<sub>5</sub> & phenyl H<sub>3</sub> & H<sub>5</sub>); LC-MS (ESI) : 395.0 (M+1, 100). Anal. Calcd. for  $C_{19}H_{14}N_4O_4S$ ; C, 57.86; H, 3.58; N, 14.21. Found: C, 57.66; H, 3.41; N, 14.01.

#### 11c

Yield 71%; Orange powder; mp 205.5-206.8 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 1732 (C=O); <sup>1</sup>HNMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  (ppm) 1.09 (3H, t, = 7.3 Hz, -CH<sub>3</sub>), 1.89 (2H, sixtet, = 7.3 Hz, CH<sub>3</sub>-C ), 3.30 (2H, t, = 7.2 Hz, CH<sub>2</sub>-C ), 7.30 (1H, d, = 8.3 Hz, H<sub>8</sub>), 7.34 (1H, t, = 7.5 Hz, H<sub>6</sub>), 7.70 (1H, t, = 8.4 Hz, 7.2, H<sub>7</sub>), 7.78 (2H, d, = 8.6 Hz, phenyl H<sub>2</sub> & H<sub>6</sub>), 8.09-8.1 (3H, m, H<sub>8</sub> & phenyl H<sub>3</sub> & H<sub>5</sub>); LC-MS (ESI) : 409.2 (M+1, 100). Anal. Calcd. for C<sub>20</sub>H<sub>16</sub>N<sub>4</sub>O<sub>4</sub>S; C, 58.82; H, 3.95; N, 13.72. Found: 58.61; H, 3.77; N, 13.55.

### 11d

Yield 75%; Orange powder; mp 243.1-243.8 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 1737 (C=O); <sup>1</sup>HNMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  (ppm) 4.58 (2H, s, -CH<sub>2</sub>), 7.35-7.39 (5H, m, H<sub>6</sub> & H<sub>8</sub> & benzyl H<sub>3</sub> & H<sub>4</sub> & H<sub>5</sub>), 7.50 (2H, d, = 7.2 Hz, benzyl H<sub>2</sub> & H<sub>6</sub>), 7.74 (1H, m, H<sub>7</sub>), 7.82 (2H, d, = 8.2 Hz, phenyl H<sub>2</sub> & H<sub>6</sub>), 8.13-8.15 (3H, m, H<sub>5</sub> & phenyl H<sub>3</sub> & H<sub>5</sub>); LC-MS (ESI) : 284.1 (M+1, 100). Anal. Calcd. for C<sub>24</sub>H<sub>16</sub>N<sub>4</sub>O<sub>4</sub>S; C, 63.15; H, 3.53; N, 12.27. Found: C, 62.95; H, 3.41; N, 12.57.

### 11e

Yield 71%; Orange powder; mp 236.8-237.5 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 1755 (C=O); <sup>1</sup>HNMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  (ppm) 4.69 (2H, s, -CH<sub>2</sub>), 7.30-7.45 (5H, m, H<sub>6</sub> & H<sub>8</sub> & 2-Cl-benzyl H<sub>4</sub> & H<sub>5</sub> & H<sub>6</sub>), 7.67-7.73 (2H, m, H<sub>7</sub> & 2-Cl-benzyl H<sub>3</sub>), 7.82 (2H, m, phenyl H<sub>2</sub> & H<sub>6</sub>), 8.13-8.4 (3H, m, H<sub>5</sub> & phenyl H<sub>3</sub> & H<sub>5</sub>), 16.26 (1H, s, -OH); LC-MS (ESI) : 491.2 (M+1, 100). Anal. Calcd. for C<sub>24</sub>H<sub>15</sub>ClN<sub>4</sub>O<sub>4</sub>S; C, 58.72; H, 3.08; N, 11.41. Found: C, 58.62; H, 3.28; N, 11.25.

### 11f

Yield 74%; Orange powder; mp 260.8-263.8 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 1748 (C=O); <sup>1</sup>HNMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  (ppm) 4.49 (2H, s, -CH<sub>2</sub>), 7.30-7.36 (4H, m, H<sub>6</sub> & H<sub>8</sub> & 4-Cl-benzyl H<sub>2</sub> & H<sub>6</sub>), 7.41 (2H, d, = 8.5, 4-Cl-benzyl H<sub>3</sub> & H<sub>5</sub>), 7.70 (1H, m, H<sub>7</sub>), 7.78 (2H, d, = 10, phenyl H<sub>2</sub> & H<sub>6</sub>), 8.09 (3H, m, H<sub>5</sub> & phenyl H<sub>3</sub> & H<sub>5</sub>); LC-MS (ESI) : 514.3 (M+23, 100). Anal. Calcd. for C<sub>24</sub>H<sub>15</sub>ClN<sub>4</sub>O<sub>4</sub>S; C, 58.72; H, 3.08; N, 11.41. Found: C, 58.98; H, 3.23; N, 11.30.

### 11g

Yield 58%; Orange powder; mp 240.1-242.2 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 1752 (C=O); <sup>1</sup>HNMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  (ppm) 4.61 (2H, s, -CH<sub>2</sub>), 7.11-7.17 (2H, m, 2-F-benzyl H<sub>3</sub> & H<sub>4</sub>), 7.34-7.51 (3H, m, H<sub>8</sub> & 2-F-benzyl H<sub>5</sub> & H<sub>6</sub>), 7.60 (1H, t, =7.5 Hz, H<sub>6</sub>), 7.74 (1H, t, = 7.2 Hz, H<sub>7</sub>), 7.83 (2H, d, = 8.5 Hz, phenyl H<sub>2</sub> & H<sub>6</sub>), 8.14-8.16 (3H, m, H<sub>5</sub> & phenyl H<sub>3</sub> & H<sub>5</sub>), 16.31 (1H, s, -OH); LC-MS (ESI) : 475.0 (M+1, 100). Anal. Calcd. for C<sub>24</sub>H<sub>15</sub>FN<sub>4</sub>O<sub>4</sub>S; C, 60.76; H, 3.19; N, 11.81. Found: C, 60.90; H, 3.03; N, 11.97.

### 11h

Yield 64%; Orange powder; mp 246.6-251.6 °C; IR (KBr disk):  $\upsilon$  (cm<sup>-1</sup>) 1763 (C=O); <sup>1</sup>HNMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  (ppm) 4.56 (2H, s, -CH<sub>2</sub>), 7.07 (2H, t, = 8.5 Hz, 4-F-benzyl H<sub>3</sub> & H<sub>5</sub>), 7.35-7.41 (2H, m, H<sub>6</sub> & H<sub>8</sub>), 7.49-7.51 (2H, dd, = 8.2 Hz, 5, 4-F-benzyl H<sub>2</sub> & H<sub>6</sub>), 7.75 (1H, t, = 7.5 Hz, H<sub>7</sub>), 7.83 (2H, d, = 8.9 Hz, phenyl H<sub>2</sub> & H<sub>6</sub>), 8.14-8.15 (3H, m, H<sub>8</sub> & phenyl H<sub>3</sub> & H<sub>5</sub>), 16.31 (1H, s, -OH); LC-MS (ESI) : 475.0 (M+1, 100). Anal. Calcd. for C<sub>24</sub>H<sub>15</sub>FN<sub>4</sub>O<sub>4</sub>S; C, 60.76; H, 3.19; N, 11.81. Found: C, 61.01; H, 3.37; N, 11.69.

Molecular modeling was performed using the Autodock Vina (28). According to the literature, X-ray crystallographic structure of prototype Table 1. Biological assay and docking results of compounds 11a-h.



		Single cycle HIV-1 replicat				
Compounds	R	(%) Inhibition rate of p24 expression	% Cell viability	Affinity binding (kcal/mol)		
11a	methyl	79	51	-9.5		
11b	ethyl	5	100	-9.5		
11c	propyl	32	93	-9.5		
11d	benzyl	6	94	-9.5		
11e	2-chlorobenzyl	33	75	-9.8		
11f	4-chlorobenzyl	52	100	-9.8		
11g	2-fluorobenzyl	22	74	-9.0		
11h	4-fluorobenzyl	25	100	-9.8		
AZT		58	92	-		

foamy virus (PFV) IN (PDB: 30YA) in complex with DNA, two Mg<sup>2+</sup> cations and raltegravir has many similarities with secondary structure of HIV-IN (RMSD 1.04 Å). Thus, it could be considered as a good model for the development of INSTIs (29). The protein and ligands were prepared in Autodock tools 1.5.6 from MGL Tools package (30). The co-crystallized ligand and water molecules were extracted, Kollman charges were added, nonpolar hydrogens were merged and AutoDock4 atom type assigned to the protein structure. The ligand was created and minimized using HyperChem 8.0 (31). The active site was defined as a Grid box around the crystallographic ligand raltegravir in 20×20×20 dimensions. All molecules were docked in the active site and the bioactive conformations were generated using Autodock Vina.

Anti-HIV-1 activity of synthesized compounds was evaluated in single cycle replication assay which was evaluated in our laboratory and reported previously (32-35). In this assay, the single-cycle replicable HIV NL4-3 virions (200 ng p24) were inoculated concurrently with compounds in different concentrations to the Hela cells. The inhibition rate (%) of p24 expression was measured by capture ELISA (Biomerieux, France) 72 hours after inoculation. Percentage inhibition of p24 expression in treated culture was calculated as inhibition rate of p24 (%). The XTT (sodium3-[1(phenylaminocarbonyl)-3,4-tetrazolium]-bis(4-methoxy-6-nitro)benzene sulfonic acid) proliferation assay was conducted to evaluate the cellular toxicity according to kit instructions (36, 37). Cytotoxicity test was performed exactly after the p24 assay.

#### **Results and Discussion**

Compounds **11a-h** were assayed for anti-HIV-1 activity in single cycle replication method which reflects only one round of infection. Anti-HIV activity of compounds was measured as inhibition rate (%) of p24 expression in Hela cells cultures. Azidothymidine (AZT) was the positive control in the test. The cytotoxic properties were measured in the same cells by XTT proliferation assay and reported as percentage of cell viability. Results are summarized in Table 1. According to our results, all compounds indicated anti-HIV activity in the range of 5-79% at 100  $\mu$ M concentration. AZT as positive control exhibited inhibition rate of 58% in our assay. Generally, all the tested compounds proved no considerable cytotoxic properties (cell viability > 74%) except compound 11a with 51% cell viability. Hence, it can be concluded that the anti-HIV-1 activity of the compounds was not a result of their cytotoxic effects. As it was expected, compounds containing halobenzyl substitutions exhibited better anti-HIV activity as compared to alkyl substitutions. Although compound 11a containing methyl substitution displayed inhibition rate of 79%, its cell viability was 51% which indicated that its anti-HIV-1 activity might be derived from cytotoxic property. Among alkyl substitutions, propyl (compound 11c) was reasonably well tolerated, giving similar results to the 2-chlorobenzyl substitution. Introduction of an unsubstituted benzyl group (compound 11d) was not well tolerated, resulting in reduction in anti-HIV-1 activity (inhibition rate = 6%). Introduction of fluorine atom in position 2 and 4 of benzyl ring (compound 11g and 11h, respectively) gave the similar inhibition rates (22 and 25%, respectively). However, compound 11g with fluorine atom in position 2 showed less %cell viability. The best anti-HIV-1 activity was observed with compounds containing chlorobenzyl substitution (compounds 11e and 11f). The shift of the chlorine atom from position 4 (compound **11f**) to position 2 (compound 11e) was associated with decrease in anti-HIV activity (inhibition rate from 52 to 33%) and increase in cytotoxicity (cell viability from 100 to 75%). It can be concluded that 4-chlorobenzyl moiety at C-5 position of 1,3,4-oxadiazol ring was optimal in terms of anti-HIV activity and cell viability.

A molecular docking study was conducted in order to evaluate synthesized compounds interactions with the integrase active site. To validate the docking study, the co-crystalized ligand, raltegravir was redocked under the same condition and superimposed on co-crystalized ligand pose (RMSD = 0.001), and displayed high affinity binding (-12.8 kcal/mol).

Docking results revealed that all the docked compounds accommodated well in the active site with affinity binding energy range from -9.0 to -9.8 kcal/mol (Table 1). Compound 11f showed high binding affinity (-9.8 kcal/mol). The 3D and 2D alignment of compound 11f in the active site was shown in Figure 3. The nitrogen atoms of 1,3,4-oxadiazole ring have been located in the 2.28 and 2.83 Å distance from Mg<sup>2+</sup> essential cofactors and was involved in metal acceptor interaction with the Mg<sup>2+</sup> ions. Phenyl and 1,3,4-oxadiazole motifs of the structure have been interacted via  $\pi$ -stacking bonding with deoxy adenosine (DA-17) presenting in the active site and could be expected to displace it. In addition, the 4-hydroxycoumarin motif was able to participate in  $\pi$ -stacking bonding with the Tyr212 residue.

Orientation compound of **11f** and co-crystalized raltegravir in the active site shown in Figure 4 clearly emphasized that both compounds occupy the same place and are engaged in similar interactions with active site residues. Nitrogen atoms of 1,3,4-oxadiazole ring of compound 11f and chelating motif of raltegravir are nearly superimposable. 4-Chlorobenzyl group of compound 11f and 4-florobenzyl group of raltegravir have been similarly oriented towards the hydrophobic pocket. 4-Hydroxycoumarin motif of compound 11f and oxadiazole ring of raltegravir occupy the same area as well.

Collectively, the docking results of tested compounds were in a good agreement with IN inhibitors, suggesting that their anti-HIV activity may be due to IN inhibition.

Oral bioavailability is considered as an important factor in design of new molecules as therapeutic agents. Thus, in drug design it is valuable to gain sufficient information about the molecular properties that limit oral bioavailability. There are some analyses that have correlated physical properties of molecules





Figure 3. (a) 2D alignment of best docked conformer of compound 11f and (b) 3D alignment of best docked conformer of compound 11f (shown in pink) in the PFV IN active site.

Alimi Livani Z / JJPR (2018), 17 (Special Issue 2): 65-77

No.	Log P	MW	HBA	HBD	Violations	nROTB	TPSA	ABS%	Volume	Log S
11a	3.32	380.38	8	1	0	4	114.09	69.64	306.47	-6.300
11b	3.82	394.41	8	1	0	5	114.09	69.64	323.27	-6.532
11c	4.27	408.44	8	1	0	6	114.09	69.64	340.07	-6.802
11d	4.81	456.48	8	1	0	6	114.09	69.64	378.12	-7.881
11e	5.41	490.93	8	1	1	6	114.09	69.64	391.65	-8.617
11f	5.41	490.93	8	1	1	6	114.09	69.64	391.65	-8.617
11g	4.91	474.47	8	1	0	6	114.09	69.64	383.05	-8.195
11h	4.91	474.47	8	1	0	6	114.09	69.64	383.05	-8.195
RAL	-0.69	444.42	11	3	0	6	152.25	56.47	375.58	-1.978
ELV	3.34	447.89	6	2	0	7	88.77	78.37	383.21	-5.655
DTG	0.12	419.38	8	2	0	3	100.87	74.20	345.56	-2.579

Table 2.molecular properties calculations of the synthesize compounds 11a-h.

TPSA: Polar surface area ( $Å^2$ ), %ABS: Absorption percentage, Vol: Volume ( $Å^3$ ), HBA: Number of hydrogen bond acceptor, HBD: Number of hydrogen bond donor, Violation: Number of violation from Lipinski's rule of five., Log P: Calculated lipophilicity., Log S: Solubility parameter, RAL= raltegravir, ELV= elvitegravir, DTG= dolutegravir.

with successful drug development. Among them, Lipinski's rule of 5 (38) and Veber's criteria (39) are the common principles utilized to evaluate the drug-like properties of a compound. In this study, Molinspiration online property calculation toolkit (40) was used to calculate the Lipinski's molecular properties and the number of rotatable bonds (nROTB), together with the topographical polar surface area (TPSA; a sum of polar atoms' surfaces: a descriptor for drug absorption, penetrability and bioavailability), the percentage of absorption (ABS%) calculated as (ABS% =109 - 0.345  $\times$  TPSA) (41) and the molecular volume (a determinant of the transport characteristics). Lipinski parameters and Veber's criteria of three FDA-approved INSTIs, raltegravir (RAL), elvitegravir (ELV), and dolutegravir (DTG) were calculated in a same way for comparison.

All compounds showed log values ranged from 3.32-5.41, MW ranging from 380.38-490.93, HBA value of 8, HBD value of 1, nROTB values ranging from 3-7 (<10), TPSA values lower than 140 Å<sup>2</sup> (114.09 Å<sup>2</sup>) which all suggesting that the evaluated compounds have acceptable flexibility and are expected to possess good permeability and oral bioavailability (39). According to these

calculations, all the compounds displayed the percentage of absorption (ABS%) about 70 which is similar to ABS% of RAL, ELV, and DTG. Additionally, aqueous solubility (Log S) of the tested compounds was determined using OSIRIS Property Explorer (Data Warrior) software (42). As can be seen from Table 2, all the compounds exhibited weak to moderate aqueous solubility with Log S values ranging between -6.300 and -8.617 mol/L which are lower than solubility of RAL, ELV, and DTG.

### Conclusions

The principal objective of this study was to investigate a new chemical scaffold with anti-HIV activity. A novel structure was constructed by hybridization of fragments from well-known anti-HIV-1 lead compounds. Key to the design was the introduction of a 1,3,4-oxadiazole ring as a chelating motif associated with a halobenzyl moiety to mimic pharmacophore of INSTIs. SAR analysis revealed that compound **11f** containing 4-chlorobenzyl ring exhibited best anti-HIV-1 activity among synthesized compounds with inhibition rate of 52% near inhibition rate of AZT (58%). Docking studies showed that designed compounds bind to IN with a mode similar to co-crystalized raltegravir. Moreover,

computation of ADME properties of designed compounds showed non-violations of Lipinski's rule of 5 and Veber's criteria, suggesting that the designed compounds possess favorable physicochemical properties near to those of FDA-approved INSTIS, RAL, ELV, and DTG. However, the designed compounds displayed low aqueous solubility as compared to FDA-approved INSTIS which may be the main reason of their moderate anti-HIV-1 activity. So, these compounds could undergo further structural modifications to improve their anti-HIV-1 activity.

#### References

- Agharbaoui FE, Hoyte AC, Ferro S, Gitto R, Buemi MR, Fuchs JR, Kvaratskhelia M and De Luca L. Computational and synthetic approaches for developing Lavendustin B derivatives as allosteric inhibitors of HIV-1 integrase (2016) 123: 673-83.
- (2) Yedavalli VRK and Jeang K-T. Methylation: a regulator of HIV-1 replication? (2007)4: 9.
- (3) Engelman A and Cherepanov P. The structural biology of HIV-1: mechanistic and therapeutic insights (2012) 10: 279.
- (4) Kinch MS and Patridge E. An analysis of FDAapproved drugs for infectious disease: HIV/AIDS drugs (2014) 19: 1510-3.
- (5) Frentz D, Boucher CAB and Van De Vijver DAMC. Temporal changes in the epidemiology of transmission of drug-resistant HIV-1 across the world (2012) 14: 17-27.
- (6) Zeinalipour-Loizidou E, Nicolaou C, Nicolaides A and Kostrikis LG. HIV-1 integrase: from biology to chemotherapeutics (2007) 5: 365-88.
- (7) Pommier Y, Pilon AA, Bajaj K, Mazumder A and Neamati N. HIV-1 integrase as a target for antiviral drugs (1997) 8: 463-83.
- (8) FDA, Isentress (Raltegravir) New Drug Application (NDA) Approval Letter, Food and Drug Administration, Rockville, MD 20857, 2007. Available from: http://www.accessdata.fda.gov/drugsatfda\_docs/ appletter/2007/022145s000tr.pdf.
- (9) Cocohoba J and Dong BJ. Raltegravir: the first HIV integrase inhibitor (2008) 30: 1747-65.
- (10) FDA Stribild (elvitegravir, cobicistat, emtricitabine, tenofovir disoproxil fumarate) new drug application (NDA) approval letter [Internet], Food and Drug Administration, Silver Spring, MD 20993, 2012. Available from: http://www.accessdata.fda.gov/ drugsatfda\_docs/appletter/2012/0203100Orig1s000ltr.

pdf.

- (11) Shimura K and Kodama EN. Elvitegravir: a new HIV integrase inhibitor. (2009)20: 79-85.
- (12) FDA, Tivicay (Dolutegravir) New Drug Application (NDA) Approval Letter [Internet], Food and Drug Administration Silver, Spring MD, 2013, 20993. Available from: http://www.accessdata.fda.gov/ drugsatfda\_docs/appletter/2013/204790Orig1s000ltr. pdf.
- (13) Vandekerckhove L. GSK-1349572, a novel integrase inhibitor for the treatment of HIV infection. (2010) 11: 203-12.
- (14) Hazuda DJ, Felock P, Witmer M, Wolfe A, Stillmock K, Grobler JA, Espeseth A, Gabryelski L, Schleif W, Blau C and Miller MD. Inhibitors of Strand Transfer That Prevent Integration and Inhibit HIV-1 Replication in Cells (2000) 287: 646-50.
- (15) Pommier Y, Johnson AA and Marchand C. Integrase inhibitors to treat HIV/Aids (2005) 4: 236.
- (16) Rogolino D, Carcelli M, Sechi M, and Neamati N. Viral enzymes containing magnesium: Metal binding as a successful strategy in drug design (2012) 256: 3063-86.
- (17) Marchand C, Maddali K, Metifiot M and Pommier Y. HIV-1 IN inhibitors: 2010 update and perspectives (2009) 9: 1016-37.
- (18) Hajimahdi Z and Zarghi A. Progress in HIV-1 integrase inhibitors: A review of their chemical structure diversity (2016) 15: 595-628.
- (19) Deng J, Sanchez T, Neamati N and Briggs JM. Dynamic pharmacophore model optimization: identification of novel HIV-1 integrase inhibitors. (2006) 49: 1684-92.
- (20) Olmedo D, Sancho R, Bedoya LM, Lopez-Perez JL, Del Olmo E, Munoz E, Alcami J, Gupta MP and San Feliciano A. 3-Phenylcoumarins as inhibitors of HIV-1 replication (2012) 17: 9245-57.
- (21) Mao PC-M, Mouscadet J-F, Leh H, Auclair C and Hsu L-Y. Chemical modification of coumarin dimer and HIV-1 integrase inhibitory activity (2002) 50: 1634-7.
- (22) Lee SU, Park JH, Kwon TH, Yoo YJ, Lee JY, Shin C-G, Yoo KH and Lee YS. Synthesis and HIV-1 integrase inhibitory activities of 4-hydroxy-5-azacoumarin 3-carboxamides (2007) 28: 1510-4.
- (23) Zhang JL, Choe H, Dezube BJ, Farzan M, Sharma PL, Zhou XC, Chen LB, Ono M, Gillies S, Wu Y, Sodroski JG and Crumpacker CS. The Bis-Azo Compound FP-21399 Inhibits HIV-1 Replication by Preventing Viral Entry (1998) 244: 530-41.
- (24) Parizadeh N, Alipour E, Soleymani S, Zabihollahi R, Aghasadeghi MR, Hajimahdi Z and Zarghi A. Synthesis of novel 3-(5-(alkyl/arylthio)-1,3,4-oxadiazol-2-yl)-8-phenylquinolin-4(1)-one derivatives as anti-HIV agents (2018) 193: 225-31.

- (25) Hajimahdi Z, Ranjbar A, Suratgar AA and Zarghi A. QSAR Study on anti-HIV-1 activity of 4-oxo-1,4-dihydroquinoline and 4-oxo-4 -pyrido[1,2-] pyrimidine derivatives using SW-MLR, artificial neural network and filtering methods (2015) 14 (Supplement): 69-75.
- (26) Zarghi A and Hajimahdi Z. Substituted oxadiazoles: a patent review (2010 -2012) (2013) 23: 1209-32.
- (27) Zarghi A, Hajimahdi Z, Mohebbi S, Rashidi H, Mozaffari S, Sarraf S, Feizi M, Tabatabaee SA and Shafiee A. Design and synthesis of new 2-substituted-5-[2-(2-halobenzyloxy)phenyl]-1, 3,4-oxadiazoles as anticonvulsant agents (2008) 56: 509-12.
- (28) Trott O and Olson AJ. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading (2010) 31: 455-61.
- (29) Hare S, Gupta SS, Valkov E, Engelman A and Cherepanov P. Retroviral intasome assembly and inhibition of DNA strand transfer (2010) 464: 232-6.
- (30) Morris GM, Goodsell DS, Halliday RS, Huey R, Hart WE, Belew RK and Olson AJ. Automated Docking Using a Lamarckian Genetic Algorithm and and Empirical Binding Free Energy Function (1998) 19: 1639-62.
- (31) HyperChem(TM) Professional 8.0, Hypercube, Inc., 1115 NW 4th Street, Gainesville, Florida 32601, USA.
- (32) Hajimahdi Z, Zabihollahi R, Aghasadeghi MR and Zarghi A. Design, synthesis and docking studies of new 4-hydroxyquinoline-3-carbohydrazide derivatives as anti-HIV-1 agents (2013) 63: 192–7.
- (33) Hajimahdi Z, Zarghi A, Zabihollahi R and Aghasadeghi MR. Synthesis, biological evaluation, and molecular modeling studies of new 1,3,4-oxadiazole- and 1,3,4-thiadiazole-substituted 4-oxo-4 -pyrido[1,2-] pyrimidines as anti-HIV-1 agents (2013) 22: 2467–75.
- (34) Hajimahdi Z, Zabihollahi R, Aghasadeghi MR, Ashtiani SH and Zarghi A. Novel quinolone-3-carboxylic acid derivatives as anti-HIV-1 agents: design, synthesis,

and biological activities 1861–76.

(35) Safakish M, Hajimahdi Z, Zabihollahi R, Aghasadeghi MR, Vahabpour R and Zarghi A. Design, synthesis, and docking studies of new 2-benzoxazolinone derivatives as anti-HIV-1 agent (2017) 26: 2718–26.

(2016) 25:

- (36) Lin C-C, Cheng H-Y, Yang C-M and Lin T-C. Antioxidant and antiviral activities of L. (2002) 9: 656–64.
- (37) Scudiero DA, Shoemaker RH, Paull KD, Monks A, Tierney S, Nofziger TH, Currens MJ, Seniff D and Boyd MR. Evaluation of a soluble tetrazolium/ formazan assay for cell growth and drug sensitivity in culture using human and other tumor cell lines (1988) 48: 4827-33.
- (38) Lipinski CA, Lombardo F, Dominy BW and Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings (1997) 23: 3-25.
- (39) Veber D, Johnson S, Cheng H, Smith B, Ward K and Kopple K. Molecular properties that influence the oral bioavailability of drug candidates (2002) 45: 2615-23.
- (40) Borrenberghs D, Dirix L, De Wit F, Rocha S, Blokken J, De Houwer S, Gijsbers R, Christ F, Hofkens J, Hendrix J and Debyser Z. Dynamic Oligomerization of Integrase Orchestrates HIV Nuclear Entry (2016) 6.
- (41) Zhao Y, Abraham M, Lee J, Hersey A, Luscombe N, Beck G, Sherborne B and Cooper I. Rate-limited steps of human oral absorption and QSAR studies (2002) 19: 1446-57.
- (42) Haddad N, Mangunuru HPR, Fandrick KR, Qu B, Sieber JD, Rodriguez S, Desrosiers JN, Patel ND, Lee H, Kurouski D, Grinberg N, Yee NK, Song JHJ and Senanayake CH. Reengineered BI-DIME Ligand Core Based on Computer Modeling to Increase Selectivity in Asymmetric Suzuki-Miyaura Coupling for the Challenging Axially Chiral HIV Integrase Inhibitor (2016) 358: 3522-7.

This article is available online at http://www.ijpr.ir