



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Published Online
January 24, 2020
[https://doi.org/10.1016/S0140-6736\(20\)30185-9](https://doi.org/10.1016/S0140-6736(20)30185-9)

This online publication has been corrected. The corrected version first appeared at [thelancet.com](https://www.thelancet.com) on January 29, 2020

See [Articles](#) pages 497 and 514

For GISAID platform see
<https://www.gisaid.org/>

In December, 2019, Wuhan, Hubei province, China, became the centre of an outbreak of pneumonia of unknown cause, which raised intense attention not only within China but internationally. Chinese health authorities did an immediate investigation to characterise and control the disease, including isolation of people suspected to have the disease, close monitoring of contacts, epidemiological and clinical data collection from patients, and development of diagnostic and treatment procedures. By Jan 7, 2020, Chinese scientists had isolated a novel coronavirus (CoV) from patients in Wuhan. The genetic sequence of the 2019 novel coronavirus (2019-nCoV) enabled the rapid development of point-of-care real-time RT-PCR diagnostic tests specific for 2019-nCoV (based on full genome sequence data on the Global Initiative

than those of patients in hospitals who became infected and had comorbidities such as diabetes or chronic respiratory disease.⁵

Today, the epidemiology of both SARS and MERS viruses is mostly understood, and the same will be true for the current outbreak of 2019-nCoV if data continue to be freely shared and used to provide real-time guidance. These Articles and others being rapidly disseminated and shared will have a major role in assuring that this global collaboration occurs.

I declare no competing interests.

David L Heymann
david.heyman@lshtm.ac.uk

Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK

- 1 Chan JF-W, Yuan S, Kok K-H, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(20\)30154-9](https://doi.org/10.1016/S0140-6736(20)30154-9).
- 2 Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- 3 Outbreak of SARS-like pneumonia being investigated in China. The Japan Times, Dec 31, 2019. <https://www.japantimes.co.jp/news/2019/12/31/asia-pacific/science-health-asia-pacific/outbreak-sars-like-pneumonia-investigated-china/#.XijKzsj7TD4> (accessed Jan 23, 2020).
- 4 Heymann DL, Rodier G. SARS: lessons from a new disease. In: Knobler S, Mahmoud A, Lemon S, et al, eds. Learning from SARS: preparing for the next disease outbreak: workshop summary. Washington, DC, USA: National Academic Press, 2004.
- 5 Petersen E, Hui DS, Perlman S, Zumla A. Middle East respiratory syndrome—advancing the public health and research agenda on MERS—lessons from the South Korea outbreak. *Int J Infect Dis* 2015; 36: 54–55.

on Sharing All Influenza Data [GISAID] platform). Cases of 2019-nCoV are no longer limited to Wuhan. Nine exported cases of 2019-nCoV infection have been reported in Thailand, Japan, Korea, the USA, Vietnam, and Singapore to date, and further dissemination through air travel is likely.^{1–5} As of Jan 23, 2020, confirmed cases were consecutively reported in 32 provinces, municipalities, and special administrative regions in China, including Hong Kong, Macau, and Taiwan.³

but at th
and perha
reported
appears to
Health
any emer
be initiall
treat patie
handwash
measures
at great

Jan 23, 2020, a total of 835 cases with laboratory-confirmed 2019-nCoV infection have been detected in China, of whom 25 have died and 93% remain in hospital (figure).³

In *The Lancet*, Chaolin Huang and colleagues⁷ report clinical features of the first 41 patients admitted to the designated hospital in Wuhan who were confirmed to be infected with 2019-nCoV by Jan 2, 2020. The study findings provide first-hand data about severity of the emerging 2019-nCoV infection. Symptoms resulting from 2019-nCoV infection at the prodromal phase, including fever, dry cough, and malaise, are non-specific. Unlike human coronavirus infections, upper respiratory symptoms are notably infrequent. Intestinal presentations observed with SARS also appear to be uncommon, although two of six cases reported by Chan and colleagues had diarrhoea.⁶ Common laboratory findings on admission to hospital include lymphopenia and bilateral ground-glass opacity or consolidation in chest CT scans. These clinical presentations confounded early detection of infected cases, especially against a background of ongoing influenza and circulation of other respiratory viruses. Exposure history to the Huanan Seafood Wholesale market served as an important clue at the early stage, yet its value has decreased as more secondary and tertiary cases have appeared.

Of the 41 patients in this cohort, 22 (55%) developed severe dyspnoea and 13 (32%) required admission to an intensive care unit, and six died.⁷ Hence, the case-fatality proportion in this cohort is approximately 14.6%, and the overall case fatality proportion appears to be closer to 3% (table). However, both of these estimates should be treated with great caution because not all patients have concluded their illness (ie, recovered or died) and the true number of infections and full disease spectrum are unknown. Importantly, in emerging viral infection outbreaks the case-fatality ratio is often overestimated in the early stages because case detection is highly biased towards the more severe cases. As further data on the spectrum of mild or asymptomatic infection becomes available, one case of which was documented by Chan and colleagues,⁶ the case-fatality ratio is likely to decrease. Nevertheless, the 1918 influenza pandemic is estimated to have had a case-fatality ratio of less than 5%¹³ but had an enormous impact due to widespread transmission, so there is no room for complacency.




Figure: Timeline of early stages of 2019-nCoV outbreak
2019-nCoV=2019 novel coronavirus.

	2019-nCoV*	MERS-CoV	SARS-CoV
Demographic			
Date	December, 2019	June, 2012	November, 2002
Location of first detection	Wuhan, China	Jeddah, Saudi Arabia	Guangdong, China
Age, years (range)	49 (21–76)	56 (14–94)	39.9 (1–91)
Male:female sex ratio	2.7:1	3.3:1	1.1:2.5
Confirmed cases	835†	2494	8096
Mortality	25† (2.9%)	858 (37%)	744 (10%)
Health-care workers	16‡	9.8%	23.1%
Symptoms			
Fever	40 (98%)	98%	99–100%
Dry cough	31 (76%)	47%	29–75%
Dyspnoea	22 (55%)	72%	40–42%
Diarrhoea	1 (3%)	26%	20–25%
Sore throat	0	21%	13–25%
Ventilatory support	9.8%	80%	14–20%

Data are n, age (range), or n (%) unless otherwise stated. 2019-nCoV=2019 novel coronavirus. MERS-CoV=Middle East respiratory syndrome coronavirus. SARS-CoV=severe acute respiratory syndrome coronavirus. *Demographics and symptoms for 2019-nCoV infection are based on data from the first 41 patients reported by Chaolin Huang and colleagues (admitted before Jan 2, 2020). †Case numbers and mortalities are updated up to Jan 21, 2020) as disclosed by the Chinese Health Commission. ‡Data as of Jan 23, 2020. §Data as of Jan 21, 2020.⁹

Table: Characteristics of patients who have been infected with 2019-nCoV, MERS-CoV, and SARS-CoV^{18, 10–12}

As an RNA virus, 2019-nCoV still has the inherent feature of a high mutation rate, although like other coronaviruses the mutation rate might be somewhat lower than other RNA viruses because of its genome-encoded exonuclease. This aspect provides the possibility for this newly introduced zoonotic viral pathogen to adapt to become more efficiently transmitted from person to person and possibly become more virulent.

Two previous coronavirus outbreaks had been reported in the 21st century. The clinical features of 2019-nCoV, in comparison with SARS-CoV and Middle East respiratory syndrome (MERS)-CoV, are summarised in the table. The ongoing 2019-nCoV outbreak has undoubtedly caused the memories of the SARS-CoV outbreak starting 17 years ago to resurface in many people. In November, 2002, clusters of pneumonia of unknown cause were reported in Guangdong province, China, now known as the SARS-CoV outbreak. The number of cases of SARS increased substantially in the next year in China and later spread globally,¹⁴ infecting at least 8096 people and causing 774 deaths.¹² The international spread of SARS-CoV in 2003 was attributed to its strong transmission ability under specific circumstances and the insufficient preparedness and implementation of infection control practices. Chinese public health and scientific capabilities have been greatly transformed since 2003. An efficient system is ready for monitoring and responding to infectious disease outbreaks and the 2019-nCoV pneumonia has been quickly added to the Notifiable Communicable Disease List and given the highest priority by Chinese health authorities.

The increasing number of cases and widening geographical spread of the disease raise grave concerns about the future trajectory of the outbreak, especially with the Chinese Lunar New Year quickly approaching. Under normal circumstances, an estimated 3 billion trips would be made in the Spring Festival travel rush this year, with 15 million trips happening in Wuhan. The virus might further spread to other places during this festival period and cause epidemics, especially if it has acquired the ability to efficiently transmit from person to person.

Consequently, the 2019-nCoV outbreak has led to implementation of extraordinary public health measures to reduce further spread of the virus within China and elsewhere. Although WHO has not recommended any international travelling restrictions so far,¹⁵ the local government in Wuhan announced on Jan 23, 2020, the suspension of public transportation, with closure of airports, railway stations, and highways in the city, to prevent further disease transmission.¹⁶ Further efforts in travel restriction might follow. Active surveillance for new cases and close monitoring of their contacts are being implemented. To improve detection

efficiency, front-line clinics, apart from local centres for disease control and prevention, should be armed with validated point-of-care diagnostic kits.

Rapid information disclosure is a top priority for disease control and prevention. A daily press release system has been established in China to ensure effective and efficient disclosure of epidemic information. Education campaigns should be launched to promote precautions for travellers, including frequent hand-washing, cough etiquette, and use of personal protection equipment (eg, masks) when visiting public places. Also, the general public should be motivated to report fever and other risk factors for coronavirus infection, including travel history to affected area and close contacts with confirmed or suspected cases.

Considering that substantial numbers of patients with SARS and MERS were infected in health-care settings, precautions need to be taken to prevent nosocomial spread of the virus. Unfortunately, 16 health-care workers, some of whom were working in the same ward, have been confirmed to be infected with 2019-nCoV to date, although the routes of transmission and the possible role of so-called super-spreaders remain to be clarified.⁹ Epidemiological studies need to be done to assess risk factors for infection in health-care personnel and quantify potential subclinical or asymptomatic infections. Notably, the transmission of SARS-CoV was eventually halted by public health measures including elimination of nosocomial infections. We need to be wary of the current outbreak turning into a sustained epidemic or even a pandemic.

The availability of the virus' genetic sequence and initial data on the epidemiology and clinical consequences of the 2019-nCoV infections are only the first steps to understanding the threat posed by this pathogen. Many important questions remain unanswered, including its origin, extent, and duration of transmission in humans, ability to infect other animal hosts, and the spectrum and pathogenesis of human infections. Characterising viral isolates from successive generations of human infections will be key to updating diagnostics and assessing viral evolution. Beyond supportive care,¹⁷ no specific coronavirus antivirals or vaccines of proven efficacy in humans exist, although clinical trials of both are ongoing for MERS-CoV and one controlled trial of ritonavir-boosted lopinavir

monotherapy has been launched for 2019-nCoV (ChiCTR2000029308). Future animal model and clinical studies should focus on assessing the effectiveness and safety of promising antiviral drugs, monoclonal and polyclonal neutralising antibody products, and therapeutics directed against immunopathologic host responses.

We have to be aware of the challenge and concerns brought by 2019-nCoV to our community. Every effort should be given to understand and control the disease, and the time to act is now.

FGH reports personal fees from University of Alabama Antiviral Drug Discovery and Development Consortium, and is a non-compensated consultant for Gilead Sciences, Regeneron, and SAB Biotherapeutics, which have investigational therapeutics for coronavirus infections. All other authors declare no competing interests.

*Chen Wang, Peter W Horby, Frederick G Hayden, George F Gao
cyh-birm@263.net

Department of Pulmonary and Critical Care Medicine, Center of Respiratory Medicine, China-Japan Friendship Hospital, Beijing 100029, China (CW); National Clinical Research Center for Respiratory Diseases, Beijing, China (CW); Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China (CW); Institute of Respiratory Medicine, Chinese Academy of Medical Sciences, Beijing, China (CW); Department of Respiratory Medicine, Capital Medical University, Beijing, China (CW); Centre for Tropical Medicine and Global Health, University of Oxford, Oxford, UK (PWH); Department of Medicine, University of Virginia School of Medicine, Charlottesville, VA, USA (FGH); and National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention (China CDC), Beijing, China (GFG)

- 1 WHO. Novel coronavirus – Thailand (ex-China). Geneva: World Health Organization, Jan 14, 2020. <https://www.who.int/csr/don/14-january-2020-novel-coronavirus-thailand/en/> (accessed Jan 23, 2020).
- 2 WHO. Novel Coronavirus – Japan (ex-China). Geneva: World Health Organization, Jan 16, 2020. <https://www.who.int/csr/don/16-january-2020-novel-coronavirus-japan-ex-china/en/> (accessed Jan 23, 2020).
- 3 China National Health Commission. Update on the novel coronavirus pneumonia outbreak (Jan 24, 2020). Beijing: China National Health Commission, 2020. <http://www.nhc.gov.cn/xcs/yqfkdt/202001/c5da49c4c5bf4bcfb320ec2036480627.shtml> (accessed Jan 24, 2020).
- 4 WHO. Novel coronavirus – Republic of Korea (ex-China). Geneva: World Health Organization, 2020. <https://www.who.int/csr/don/21-january-2020-novel-coronavirus-republic-of-korea-ex-china/en/> (accessed Jan 24, 2020).

- 5 US Centers for Disease Control and Prevention. First travel-related case of 2019 novel coronavirus detected in United States. Atlanta, GA: US Centers for Disease Control and Prevention, 2020. <https://www.cdc.gov/media/releases/2020/p0121-novel-coronavirus-travel-case.html> (accessed Jan 24, 2020).
- 6 Chan JF-W, Yuan S, Kok K-H, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(20\)30154-9](https://doi.org/10.1016/S0140-6736(20)30154-9).
- 7 Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- 8 Leung GM, Hedley AJ, Ho LM, et al. The epidemiology of severe acute respiratory syndrome in the 2003 Hong Kong epidemic: an analysis of all 1755 patients. *Ann Intern Med* 2004; **141**: 662–73.
- 9 WHO. Novel coronavirus (2019-nCoV) situation report – 2 (22 January 2020). Geneva: World Health Organization, 2020. <https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200122-sitrep-2-2019-ncov.pdf> (accessed Jan 23, 2020).
- 10 WHO. Middle East respiratory syndrome coronavirus (MERS-CoV). Geneva: World Health Organization, 2020. <http://www.who.int/emergencies/mers-cov/en/> (accessed Jan 12, 2020).
- 11 Assiri A, Al-Tawfiq JA, Al-Rabeeh AA, et al. Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: a descriptive study. *Lancet Infect Dis* 2013; **13**: 752–61.
- 12 WHO. Summary of probable SARS cases with onset of illness from 1 November 2002 to 31 July 2003. Geneva: World Health Organization, 2004. http://www.who.int/csr/sars/country/table2004_04_21/en/ (accessed Jan 12, 2020).
- 13 Viboud C, Eisenstein J, Reid AH, Janczewski TA, Morens DM, Taubenberger JK. Age- and sex-specific mortality associated with the 1918–1919 influenza pandemic in Kentucky. *J Infect Dis* 2013; **207**: 721–29.
- 14 Zhong NS, Zheng BJ, Li YM, et al. Epidemiology and cause of severe acute respiratory syndrome (SARS) in Guangdong, People's Republic of China, in February, 2003. *Lancet* 2003; **362**: 1353–58.
- 15 WHO. Statement on the meeting of the International Health Regulations (2005) Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV). Geneva, World Health Organization, Jan 23, 2020. [https://www.who.int/news-room/detail/23-01-2020-statement-on-the-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/23-01-2020-statement-on-the-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov)) (accessed Jan 24, 2020).
- 16 Announcement from the Headquarter for novel coronavirus pneumonia prevention and control (No. 1). Beijing: China National Health Commission, 2020. http://www.gov.cn/xinwen/2020-01/23/content_5471751.htm (accessed Jan 23, 2020).
- 17 WHO. Clinical management of severe acute respiratory infection when novel coronavirus (nCoV) infection is suspected. Geneva, World Health Organization, 2020. [https://www.who.int/internal-publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/internal-publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected) (accessed Jan 22, 2020).

Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury



The 2019 novel coronavirus (2019-nCoV) outbreak is a major challenge for clinicians. The clinical course of patients remains to be fully characterised, little data are available that describe the disease pathogenesis, and no pharmacological therapies of proven efficacy yet exist.

Corticosteroids were widely used during the outbreaks of severe acute respiratory syndrome (SARS)-CoV¹ and

Middle East respiratory syndrome (MERS)-CoV,² and are being used in patients with 2019-nCoV in addition to other therapeutics.³ However, current interim guidance from WHO on clinical management of severe acute respiratory infection when novel coronavirus (2019-nCoV) infection is suspected (released Jan 28, 2020) advises against the use of corticosteroids unless indicated for

Published Online
February 6, 2020
[https://doi.org/10.1016/S0140-6736\(20\)30317-2](https://doi.org/10.1016/S0140-6736(20)30317-2)