

CORRECTION

Open Access



Correction: Methyl β -Cyclodextrin-sperm-mediated gene editing (MBCD-SMGE): a simple and efficient method for targeted mutant mouse production

Parisa Moradbeigi¹, Sara Hosseini^{2,3}, Mohammad Salehi^{2,4*} and Asghar Moghiseh¹

Correction: Biol Proced Online 26, 3 (2024)

<https://doi.org/10.1186/s12575-024-00230-9>

The original article has been corrected.

Following publication of the original article [1], the author reported two errors:

Published online: 16 February 2024

1. The error in Fig. 3 displaying blastocysts obtained from the 2mM CAG-GFP group instead of the intended 2mM gRNA-Cas9 group, as clarified in the figure legend within the article. The correct figure is shown in this article. Also, in the figure legend, the word 'SMGT' has been appropriately changed to 'SMGE' to reflect the accurate terminology.
2. The repetition of the title 'argeted indel was validated in blastocysts derived from sperm incubated with 2 mM MBCD and 20 ng/ μ l gRNA-Cas9 plasmid constructs' under Experiment 5 in the results section.

Reference

1. Moradbeigi P, Hosseini S, Salehi M, et al. Methyl β -Cyclodextrin-sperm-mediated gene editing (MBCD-SMGE): a simple and efficient method for targeted mutant mouse production. *Biol Proced Online*. 2024;26:3. <https://doi.org/10.1186/s12575-024-00230-9>.

The original article can be found online at <https://doi.org/10.1186/s12575-024-00230-9>.

*Correspondence:

Mohammad Salehi
salehiemb@gmail.com

¹ Department of Clinical Sciences, School of Veterinary Medicine, Shiraz University, P. O. Box: 7144169155, Shiraz, Iran

² Cellular and Molecular Biology Research Center, Shahid Beheshti University of Medical Sciences, P.O. Box: 193954717, Tehran, Iran

³ Hasti Noavaran Gene Royan Co, Tehran, Iran

⁴ Department of Biotechnology, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 3 Murine GFP-positive blastocyst production using the MBCD-SMGE and IVF-IVC methods. **A** Bright field and **(B)** fluorescent field images of day 4 mouse blastocysts obtained from the MBCD-SMGE and conventional IVF-IVC methods. In the MBCD-SMGE method, sperm were incubated in the c-TYH medium supplemented with 2 mM MBCD and 20 ng/μl pgRNA-Cas9. **C** Bright field and **(D)** fluorescent field images of day 4 mouse negative control blastocysts. Scale bar size: 100 μm