

RESEARCH

Open Access



The knowledge, attitude and behavior on the palliative care among neonatal nurses: what can we do

Yilan Yan^{1†}, Jiahui Hu^{1†}, Fei Hu^{1†} and Longyan Wu^{1*}

Abstract

Background Neonatal nurses should provide timely and high-quality palliative care whenever necessary. It's necessary to investigate the knowledge, attitude and behavior of palliative care among neonatal nurses, to provide references and evidences for clinical palliative care.

Methods Neonatal intensive care unit (NICU) nurses in a tertiary hospital of China were selected from December 1 to 16, 2022. The palliative care knowledge, attitude and behavior questionnaire was used to evaluate the current situation of palliative nursing knowledge, attitude and behavior of NICU nurses. Univariate analysis and multivariate logistic regression analysis were used to analyze the influencing factors.

Results 122 nurses were finally included. The average score of knowledge in neonatal nurses was 7.68 ± 2.93 , the average score of attitude was 26.24 ± 7.11 , the score of behavior was 40.55 ± 8.98 , the average total score was 74.03 ± 10.17 . Spearman correlation indicated that score of knowledge, attitude and behavior of palliative care in neonatal nurses were correlated with the age ($r = 0.541$), year of work experience ($r = 0.622$) and professional ranks and titles ($r = 0.576$) (all $P < 0.05$). Age (OR = 1.515, 95%CI: 1.204 ~ 1.796), year of work experience (OR = 2.488, 95%CI: 2.003 ~ 2.865) and professional ranks and titles (OR = 2.801, 95%CI: 2.434 ~ 3.155) were the influencing factors of score of knowledge, attitude and behavior of palliative care (all $P < 0.05$).

Public contribution NICU nurses have a positive attitude towards palliative care, but the practical behavior of palliative care is less and lack of relevant knowledge. Targeted training should be carried out combined with the current situation of knowledge, attitude and practice of NICU nurses to improve the palliative care ability and quality of NICU nurses.

Keywords Knowledge, Attitude, Behavior, Palliative care, Neonate, Nurse

[†]Yilan Yan, Jiahui Hu and Fei Hu are equal contributors.

*Correspondence:

Longyan Wu
oogw90@sina.com

¹Department of Neonatology, Children's Hospital of Nanjing Medical University, No. 72, Guangzhou Road, Gulou District, Nanjing, Jiangsu Province, China



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Neonatal intensive care unit (NICU) is a centralized ward for the treatment of neonatal critical diseases, which is established for continuous monitoring and timely and effective rescue treatment and nursing of high-risk newborns. Its purpose is to reduce neonatal mortality and promote neonatal growth and development. NICU provides life support for high-risk newborns, including rescue and treatment of neonatal diseases, neonatal respiratory management and premature infant management [1]. Previous studies [2, 3] have shown that the mortality of neonate with NICU can be as high as 8.16%. The illness and death of newborns can bring great pain and psychological burden to their families. Therefore, it is necessary to carry out corresponding palliative care for neonates and their families.

As an extension of high-quality clinical nursing service, palliative care centered on newborns, is the overall behavior of neonatal nursing service, which must be taken care of by a competent and qualified professional nursing team [4–6]. How to improve the knowledge, attitude and behavior ability of NICU nurses in palliative care is a very important issue that the hospital managers and nurses must pay attention to.

As a hot spot of social concern, palliative care for newborns has attracted more and more attention in recent years. Palliative care in China is in the initial state of development. Kain et al. in Australia [7], Wright team [8] and Kyc team [9] in the United States, Chen et al. in Taiwan [10], Forouzi et al. in southeastern Iran [11] have used the Neonatal Palliative Care attitude scale to investigate the attitude of NICU nurses towards the obstacles of providing palliative care for newborns. These studies reported some obstacles, including insufficient staff, lack of ideal implementation environment, technical requirements, parents' requirements, nurses' inability to express their opinions, lack of education, lack of effective counseling, and medical staff's personal values and attitude in palliative care for newborns [12, 13]. In order to understand the obstacles that affect palliative care for newborns, it's necessary to investigate the knowledge, attitude and behavior of neonatal nurses. The purpose of this survey was to investigate the current situation of nurses' knowledge, attitude and behavior of palliative care for newborns, and to evaluate the influence of neonatal nurses' personal and professional characteristics on their knowledge, attitude and behavior, to provide insights to the clinical palliative care for newborns.

Methods.

This study was a cross sectional survey design. The study has been reviewed and approved by the ethics committee of Children's Hospital of Nanjing Medical University (No. 202304068-1). And written informed consents had been obtained from all the nurses.

According to the principle of the application of psychometrics, the sample size in the survey is related to the number of variables, and the sample size should be about 10 to 15 times the number of variables, in order to improve the stability of the questionnaire structure and facilitate factor analysis [14]. The number of items used in this survey was 6, and 15 times the number of items was taken as the sample size of the survey, and then considering the 20% loss rate and invalid questionnaire, the required sample size is $6 \times 15 \times 1.2 = 108$. Therefore, at least 108 nurses should be included in this survey.

In this study, targeted NICU nurses in a tertiary hospital of China were surveyed from December 1 to 16, 2022. The inclusion criteria of nurses in the survey were as follows: formal employees of NICU; whose age was ≥ 18 years old, the nurses knew and agree to participate in this survey. Exclusion criteria: trainee nurses, nurses in other department, volunteers.

We designed a general information questionnaire, including gender, age, nationality, professional title, working years and so on. In addition, we used the knowledge, attitude and measured performance questionnaire [15] to evaluate the current situation of palliative nursing knowledge, attitude and behavior of NICU nurses. The questionnaire consists of 40 items, which is divided into three dimensions: knowledge, attitude and behavior. Among them, there are 20 items in the knowledge dimension, with a score of "1" for correct answers and a score of "0" for wrong or unknown answers. The higher the score, the better the nurses' knowledge. There were 12 items in the attitude dimension, which were scored by Likert-5 points (1=strong disagreement, 2=disagreement, 3=neutral, 4=consent, 5=strong agreement). The higher the score, the better the nurses' cognitive attitude. There were 8 items in the behavior dimension, which were scored by Likert5 points (1=almost none, 2=less, 3=uncertain, 4=able to do, 5=often). The higher the score, the more palliative nursing behaviors of nurses. The Cronbach α coefficients of the questionnaire were 0.758, 0.794 and 0.910, respectively [16]. The internal consistency of the questionnaire was good to evaluate the level of knowledge, attitude and behavior of palliative care of nurses.

We explained the purpose, significance and matters needing attention of this survey to the NICU nurses. After the respondents signing the informed consent form, the investigators distributed the questionnaire uniformly. We collected it on the spot after the nurse filled out the questionnaire.

SPSS 23.0 software was used for statistical analysis, the counting data were described by frequency and percentage, the measurement data were expressed by mean \pm standard deviation, and the t and F tests were used between groups. Univariate and multivariate logistic regression analysis were used to analyze the influencing

Table 1 The characteristics of included nurses

Characteristic	Cases(n = 122)	Percentage
Gender		
Female	111	90.98%
Male	11	9.02%
Age(y)		
18 ~ 30	40	32.79%
31 ~ 35	48	39.34%
> 35	34	27.87%
Education level		
Junior college degree	45	36.89%
Bachelor degree	75	61.48%
Master degree	2	1.63%
Year of work experience		
< 5	39	31.87%
5 ~ 10	36	29.51%
11 ~ 20	37	30.33%
> 20	10	8.19%
Professional ranks and titles		
Junior nurse	31	25.41%
Senior nurse	56	45.90%
Nurse in charge	38	31.15%
Deputy chief nurse	5	4.11%
Chief nurse	2	1.63%
Marital status		
Unmarried	52	42.62%
Married	70	57.38%

Table 2 The average score of knowledge, attitude and behavior of palliative care for neonatal nurses

Items	Average score	Minimum score	Maximum score
Knowledge section	7.68 ± 2.93	3	14
Attitude section	26.24 ± 7.11	18	39
Behavior section	40.55 ± 8.98	28	51
Total score	74.03 ± 10.17	55	8

factors. Spearman correlation analysis was used to analyze the correlation between nurses' characteristics and their knowledge, attitude and practice. In this study, $P < 0.05$ as the difference between groups was statistically significant.

Results

A total of 122 nurses were included. As presented in Table 1, the included nurses were mainly female, most of them were under the age of 35, their academic qualifications are mainly bachelor's degree, their working years were less than 10 years, and their professional titles were mostly junior and senior nurses.

As presented in Table 2, The average score of knowledge in neonatal nurses was 7.68 ± 2.93 , the average score of attitude was 26.24 ± 7.11 , the score of behavior was 40.55 ± 8.98 , the average total score was 74.03 ± 10.17 .

As indicated in Table 3, there were significant differences in the score of knowledge, attitude and behavior of palliative care in nurses with different age, year of work experience and professional ranks and titles (all $P < 0.05$). No significant differences in the score of knowledge, attitude and behavior of palliative care in nurses with different gender, education level, and marital status were detected (all $P > 0.05$).

As presented in Table 4, Spearman correlation indicated that score of knowledge, attitude and behavior of palliative care in neonatal nurses were correlated with the age ($r = 0.541$), year of work experience ($r = 0.622$) and professional ranks and titles ($r = 0.576$) (all $P < 0.05$).

As shown in Table 5, multivariate logistic regression analysis indicated that age (OR = 1.515, 95%CI: 1.204 ~ 1.796), year of work experience (OR = 2.488, 95%CI: 2.003 ~ 2.865) and professional ranks and titles (OR = 2.801, 95%CI: 2.434 ~ 3.155) were the influencing factors of score of knowledge, attitude and behavior of palliative care in neonatal nurses (all $P < 0.05$).

Discussions

The results of this study shows that although most nurses recognize the importance of neonatal palliative care, they do not know much about it, which leads to some doubts and deficiencies in their clinical palliative care behavior. There is a lack of standardized palliative care practice guidelines and quality assessment standards and tools, and the contents and items of neonatal palliative care are not standardized [17]. Most nurses have a positive attitude towards the implementation of palliative care for newborns in this study, but it has been reported that some nurse have a negative attitude towards palliative care, mostly worried that after the implementation of palliative care, they will increase their work burden because of insufficient manpower deployment [18, 19]. It is of great significance to find out the promoting factors and obstacle factors that affect nurses' knowledge, attitude and practice [20]. Most nurses do not have enough knowledge of palliative care, especially for newborns [21]. In addition, palliative nursing education is also inadequate. Most nurses learn very few knowledge and skill of palliative nursing [9, 11].

We have found that neonatal nurses lack of knowledge related to palliative care. The knowledge of palliative nursing is the basis of nursing intervention. The evaluation of palliative care for children by the National Medical institutions Review Committee mainly includes two aspects: one is that medical institutions have the conditions to implement palliative care, and the other is that staff have the ability to implement palliative care to meet the needs of dying children and their families [22, 23]. The hospitals should provide support to nurses to enhance the competence of professionals and apply social

Table 3 The score of knowledge, attitude and behavior of palliative care in neonatal nurses with different characteristics

Characteristic	Cases(n= 122)			Knowledge			Attitude			Behavior			
		Score	t/F	P	Score	t/F	P	Score	t/F	P	Score	t/F	P
Gender													
Female	111	7.67 ±2.61	1.113	0.125	26.41±7.75	5.204	0.152	39.81±8.03	6.054	0.137			
Male	11	7.72 ±2.58			26.15±7.81			40.16±9.19					
Age(y)													
18 ~ 30	40	6.49 ±2.58	3.114	0.030	25.04±7.87	4.125	0.021	37.36±9.05	5.351	0.014			
31 ~ 35	48	7.66 ±2.91			26.14±7.53			40.22±9.85					
> 35	34	8.84 ±2.65			27.60±7.17			41.43 ±10.18					
Education level													
Junior college degree	45	7.22 ±2.48	2.126	0.098	26.13±7.86	5.102	0.143	39.05±9.64	6.312	0.057			
Bachelor degree	75	7.69 ±2.50			26.21±7.92			40.75±9.21					
Master degree	2	7.71 ±2.36			26.30±8.88			41.83±8.75					
Year of work experience													
< 5	39	6.41 ±2.46	1.486	0.016	23.13±6.75	6.078	0.044	38.03±9.54	7.172	0.036			
5 ~ 10	36	7.59 ±2.47			24.30±7.94			40.60±8.43					
11 ~ 20	37	7.74 ±2.55			25.04±6.81			41.01±9.15					
> 20	10	8.83 ±2.49			25.19±7.05			41.66±9.54					
Professional ranks and titles													
Junior nurse	31	6.50 ±2.85	3.123	0.041	23.85±8.02	5.458	0.013	38.04±8.29	8.146	0.009			
Senior nurse	56	7.62 ±2.54			24.16±9.13			40.39±9.05					
Nurse in charge	38	7.87 ±2.61			24.29±8.85			41.92±8.57					
Deputy chief nurse	5	8.07 ±2.51			25.19±7.72			41.96±9.23					
Chief nurse	2	8.89 ±2.42			25.35±7.68			42.03±8.84					
Marital status													
Unmarried	52	7.66 ±2.56	1.278	0.207	23.95±6.81	6.794	0.287	40.27±9.84	8.094	0.114			
Married	70	7.68 ±2.44			24.78±5.99			40.68±8.09					

Table 4 Analysis of the correlation between total score and characteristics of neonatal nurses

Characteristic	<i>r</i>	<i>P</i>
Gender	0.143	0.095
Age(y)	0.541	0.021
Education level	0.108	0.167
Year of work experience	0.622	0.015
Professional ranks and titles	0.576	0.037
Marital status	0.113	0.102

Table 5 Multivariate logistic regression analysis on the influencing factors of score of knowledge, attitude and practice of palliative care in neonatal nurses

Variables	β	S^2x	OR	95%CI	<i>P</i>
Age	0.139	0.101	1.515	1.204~1.796	0.031
Year of work experience	0.143	0.102	2.488	2.003~2.865	0.017
Professional ranks and titles	0.149	0.123	2.801	2.434~3.155	0.014

support through research and continuing education [24]. The knowledge, skills and hospital conditions of the parents and medical staff are the important factors affecting the implementation of palliative care in NICU, and the nursing staff are the key factors. Therefore, having certain knowledge and ability to implement palliative care is the basis for the implementation of high-quality palliative care, it is necessary to carry out palliative care training in clinical nursing education.

The neonatal nurses have a positive attitude toward the palliative care in this survey. Some studies [25–27] have shown that the implementation of palliative care is related to nurses’ knowledge, attitude, experience and evaluation ability, and these factors will affect the implementation effect. Some studies [28–30] have shown that the attitude of medical staff, whether positive or negative, can have a lasting impact on parents. A positive attitude can help parents alleviate their grief and retain good memories in the process. Not paying attention to finding psychological problems and giving support will make families feel helpless [31]. Therefore, for NICU nurses, they must be able to face the death of neonate with a correct state of mind, and receive systematic palliative education [32, 33]. The continuing education and training of palliative care should attract the attention of the health care providers and managers of relevant universities and medical institutions [34].

The results of this survey show that the palliative nursing behavior of nurses is insufficient. The concept of Chinese traditional culture has brought challenges to the popularization of palliative care in China [35]. The traditional Chinese view of life and death holds that talking about death is a symbol of misfortune and fear, which makes Chinese people afraid to talk about death [36]. This concept is deeply rooted in the hearts of the people, which not only makes family members avoid

talking about death, but also brings certain psychological pressure for medical staff to talk about “death” and “palliative care” to parents, which hinders the behavior of palliative care for newborns [37–39]. The cause of palliative care for newborns should be raised to the level of safeguarding human dignity, reducing social burden and reflecting human civilization [40]. In addition, it is necessary to improve the relevant policies and regulations, bring palliative care into the basic medical insurance system, improve the professional quality of nursing care, and strengthen the input of manpower and material resources needed for palliative care [41, 42]. Clinical nursing managers should strengthen the training of palliative nursing knowledge and operation of neonatal nurses and encourage nurses’ palliative nursing behavior.

There are some shortcomings in this survey that deserve careful consideration. First of all, this study is a single-center investigation, the study sample size is small, we cannot perform correlation and multi-factor analysis on specific scores of knowledge, attitude and behavior. Secondly, there are few personal characteristic data of nurses included in this study, and other factors affecting nurses’ knowledge, attitude and behavior of palliative care may not be included in the analysis. The sample size should be expanded in future research. More factors were prospectively included to analyze the current situation and influencing factors of neonatal palliative care, so as to provide evidence support for clinical neonatal palliative care.

Conclusion

In summary, the results of this survey have indicated that neonatal nurses lack of knowledge related to palliative care and have a positive attitude, but their behavior is relatively insufficient. And age, year of work experience and professional ranks and titles are the influencing factors of score of knowledge, attitude and behavior of palliative care in neonatal nurses. Palliative care is an important part of NICU care. Through the training of nurses’ professional knowledge and skills, NICU nurses’ understanding of palliative care and clinical nursing ability should be effectively improved. Only when nurses correctly understand the connotation of palliative care and the relevant knowledge and skills of palliative care can they provide high-quality palliative nursing care to newborns and their families.

Abbreviations

NICU neonatal intensive care unit

Acknowledgements

None.

Author contributions

YY, J H designed research; YY, J H, F H, L W conducted research; YY, J H, L W analyzed data; YY, L W wrote the first draft of manuscript; YY, L W had

primary responsibility for final content. All authors read and approved the final manuscript.

Funding

None.

Data availability

All data generated or analyzed during this study are included in this published article. The original data will be available from corresponding authors on reasonable request.

Declarations

Ethics approval and consent to participate

In this study, all methods were performed in accordance with the relevant guidelines and regulations. The study has been reviewed and approved by the ethics committee of Children's Hospital of Nanjing Medical University (No. 202304068-1). And written informed consents had been obtained from all the nurses.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 7 July 2023 / Accepted: 22 May 2024

Published online: 03 July 2024

References

- Carrier L, Dame J, Wright AL, Latimer M. Involving families in the care of indigenous infants: a philosophical exploration of indigenous ways of knowing to inform nursing research and practice in the neonatal intensive care unit. *J Clin Nurs*. 2023;32(7–8):1495–505.
- Yuan J, Zhao F, Wang J. Clinical characteristics and epidemiological investigation of 13,476 intensive care newborns in Yinchuan from 2014 to 2019. *China Maternal Child Health*. 2022;36(3):11–5.
- Jiang L, Li X, Liao Z. Clinical characteristics and death risk factors of neonates in NICU. *Chin J Hosp Infect*. 2023;14(6):10–4.
- Maher-Griffiths C. Perinatal Palliative Care in the neonatal Intensive Care Unit. *Crit Care Nurs Clin North Am*. 2022;34(1):103–19.
- Currie ER, Wolfe J, Boss R, Johnston EE, Paine C, Perna SJ, Buckingham S, KcKillip KM, Li P, Dionne-Odom JN et al. Patterns of pediatric palliative and end-of-life care in neonatal intensive care patients in the Southern U.S. *J Pain Symptom Manage* 2023.
- Barry A, Prentice T, Wilkinson D. End-of-life care over four decades in a quaternary neonatal intensive care unit. *J Paediatr Child Health*. 2023;59(2):341–5.
- Kain V, Gardner G, Yates P. Neonatal palliative care attitude scale: development of an instrument to measure the barriers to and facilitators of palliative care in neonatal nursing. *Pediatrics*. 2009;123(2):e207–213.
- Wright V, Prasun MA, Hilgenberg C. Why is end-of-life care delivery sporadic? A quantitative look at the barriers to and facilitators of providing end-of-life care in the neonatal intensive care unit. *Adv Neonatal Care*. 2011;11(1):29–36.
- Kyc SJ, Bruno CJ, Shabanova V, Montgomery AM. Perceptions of neonatal Palliative Care: similarities and differences between Medical and Nursing Staff in a level IV neonatal intensive care unit. *J Palliat Med*. 2020;23(5):662–9.
- Chen CH, Huang LC, Liu HL, Lee HY, Wu SY, Chang YC, Peng NH. To explore the neonatal nurses' beliefs and attitudes towards caring for dying neonates in Taiwan. *Matern Child Health J*. 2013;17(10):1793–801.
- Azzizadeh Forouzi M, Banazadeh M, Ahmadi JS, Razban F. Barriers of Palliative Care in neonatal intensive care units. *Am J Hosp Palliat Care*. 2017;34(3):205–11.
- Ng SKF, Keenan N, Swart S, Berry MJ. Palliative care in a tertiary neonatal intensive care unit: a 10-year review. *BMJ Support Palliat Care*. 2022;12(e5):e641–5.
- Smith HAB, Besunder JB, Betters KA, Johnson PN, Srinivasan V, Stormorken A, Farrington E, Golianu B, Godshall AJ, Acinelli L, et al. 2022 society of critical Care Medicine Clinical Practice guidelines on Prevention and Management of Pain, agitation, neuromuscular blockade, and Delirium in critically Ill Pediatric patients with consideration of the ICU environment and early mobility. *Pediatr Crit Care Med*. 2022;23(2):e74–110.
- Zheng W, He F. The method of calculating the sample size of current situation survey. *Prev Med*. 2020;32(6):2–6.
- Yajing Z. Knowledge, attitude and behavior of Chinese neonatal pediatric nurses towards palliative care and its influencing factors. Guangzhou: Guangzhou University of traditional Chinese Medicine; 2020.
- Xin Z, Shiyu L, Jingli C. Research progress of hospice care in neonatal intensive care unit. *Chin J Nurs*. 2012;47(6):3–6.
- Mascarenhas D, Goyal M, Nanavati R. Neonatal palliative care practices: an Indian perspective. *Paediatr Int Child Health*. 2022;42(1):22–8.
- Heidari H, Mardani-Hamoooleh M, Fooladi M. Design of a Palliative Care Program for nursing students in the neonatal intensive care unit: a mixed-method study. *Creat Nurs*. 2022;28(2):126–32.
- Torbert N, Taladay C, Kauer T, Hackenburg L, Weaver MS, Kellas JK. Providing compassionate care in the neonatal intensive care unit through infant and family needs-based care. *Am J Perinatol* 2022.
- Girgin BA, Gozen D, Aktas E, Ergun K. Attitudes toward neonatal Palliative Care among Turkish nurses and Physicians: a comparative cross-sectional study. *J Hosp Palliat Nurs*. 2022;24(5):E185–96.
- Abuhammad S, Elayyan M, Ababneh H. Neonatal intensive care unit nurses knowledge and attitude toward neonatal palliative care: review of the literature. *Future Sci OA*. 2023;9(5):FSO856.
- Quinn M, Weiss AB, Crist JD. Early for everyone: reconceptualizing Palliative Care in the neonatal Intensive Care Unit. *Adv Neonatal Care*. 2020;20(2):109–17.
- Forman KR, Thompson-Branch A. Educational perspectives: Palliative Care Education in neonatal-perinatal Medicine Fellowship. *Neoreviews*. 2020;21(2):e72–9.
- Pediatrics AAO: American Academy of Pediatrics, Committee on Bioethics and Committee on Hospital Care. Palliative care for children. *Pediatrics* 2000, 106(2 Pt 1):351–357.
- Cerratti F, Tomietto M, Della Pelle C, Kain V, Di Giovanni P, Rasero L, Cicolini G. Italian nurses' attitudes towards neonatal Palliative Care: a cross-sectional survey. *J Nurs Scholarsh*. 2020;52(6):661–70.
- Cavinder C. The relationship between providing neonatal palliative care and nurses' moral distress: an integrative review. *Adv Neonatal Care*. 2014;14(5):322–8.
- Lee MC, Chen YC, Chen CH, Lu FL, Hsiao CC, Peng NH. Comparison of the Educational needs of neonatologists and neonatal nurses regarding Palliative Care in Taiwan. *Am J Hosp Palliat Care*. 2016;33(3):264–71.
- Allen JD, Shukla R, Baker R, Slaven JE, Moody K. Improving neonatal Intensive Care Unit Providers' perceptions of Palliative Care through a Weekly Case-based discussion. *Palliat Med Rep*. 2021;2(1):93–100.
- Hasanpour M, Pouraboli B, Mohammadpour M, Tahmasebi M, Sabeti F. Challenges of paediatric palliative care in the intensive care unit during the COVID-19 pandemic. *Int J Palliat Nurs*. 2021;27(6):303–15.
- Rusalef F, Cavicchiolo ME, Lago P, Salvadori S, Benini F. Perinatal palliative care: a dedicated care pathway. *BMJ Support Palliat Care*. 2021;11(3):329–34.
- Mohammadi A, Tahmasebi M, Mojen LK, Rassouli M, Ashrafzadeh H. Evaluation of Care Providers' attitude toward Perinatal Palliative Care and its challenges in the selected teaching hospitals of Tehran in 2019. *Indian J Palliat Care*. 2021;27(4):513–20.
- Zhong Y, Black BP, Kain VJ, Song Y. Facilitators and barriers affecting implementation of neonatal Palliative Care by nurses in Mainland China. *Front Pediatr*. 2022;10:887711.
- Chin SDN, Paraszczuk AM, Eckardt P, Bressler T. Neonatal nurses' perceptions of Palliative Care in the neonatal Intensive Care Unit. *MCN Am J Matern Child Nurs*. 2021;46(5):250–7.
- Zhu L, Zhang N, Hu Y, Xu Y, Luo T, Xiang Y, Jiang S, Zhang Z, Chen M, Xiong Y. Influencing factors of knowledge, attitude and behavior in children's palliative care among pediatric healthcare workers: a cross-sectional survey in China. *BMC Palliat Care*. 2023;22(1):67.
- Liu M, Zhao H, Hu D. Qualitative study on palliative care cognition and behavior of medical staff in ICU. *J Qingdao Univ*. 2022;58(4):4–6.
- Zhang A, Bing L, Mi Q, Zhou F, Wang J. Pediatric Palliative Care for Children with Cancer in a children's Tertiary Hospital in China: six-year experience of a Pediatric Palliative Care Service. *Palliat Med Rep*. 2021;2(1):1–8.
- Liu X, Wu X, Cheng Q, Ying W, Gong X, Lu D, Zhang Y, Liu Z. Meaning in life and its relationship with family cohesion: a survey of patients with palliative care in China. *Asia Pac J Oncol Nurs*. 2022;9(11):100118.

38. Cai S, Guo Q, Wang X, Wang R, Yang P, Zhou Y, Luo Y, Zhou X, Peng X. Development levels of pediatric palliative care teams and the extent of palliative care understanding and implementation among pediatric oncologists in China. *Pediatr Investig*. 2021;5(4):265–70.
39. Hu L, Chu Q, Fan Z, Chen Y. Discussion of advance care planning on end-of-life decisions with lung cancer patients in Wuhan, China: attitude, timing and future directions. *Intern Med J*. 2021;51(12):2111–8.
40. Ho KK, Cheng FW, Lin MK, Wai S, Lee V, Li CK. Palliative care service in patients with childhood cancer from a tertiary pediatric oncology center. *Pediatr Investig*. 2018;2(4):209–15.
41. Peng NH, Lee CH, Lee MC, Huang LC, Chang YC, DeSwarte-Wallace J. Effectiveness of Pediatric Palliative Care Education on Pediatric clinicians. *West J Nurs Res*. 2017;39(12):1624–38.
42. Chan WCH, Wong KLY, Leung MMM, Lin MKY. Perceived challenges in pediatric palliative care among doctors and nurses in Hong Kong. *Death Stud*. 2019;43(6):372–80.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.