

Multidrug-resistant *Mycobacterium abscessus* infection in an anophthalmic socket treated with bedaquiline on a compassionate use basis: A case report

David Galindo-Rodríguez ^{a,*}, Miguel Moreno Hijazo ^b, Celina Balint Ilie ^a, Daniel Rubio Castro ^a, Ignacio Vallés Tormo ^a, Eva Gloria Alias Alegre ^c

^a Department of Internal Medicine, Hospital Obispo Polanco, Teruel, Spain

^b Department of Medical Microbiology, Hospital Obispo Polanco, Teruel, Spain

^c Department of Ophthalmology, Hospital Obispo Polanco, Teruel, Spain

ARTICLE INFO

Keywords:

Mycobacterium abscessus

Bedaquiline

Endophthalmitis

Case report

systemic antibiotics.

After nine years of recurrent conjunctivitis, an infection in the anophthalmic cavity was diagnosed, characterized by marked inflammatory signs and necrosis. Despite local treatments, including intracavitary triamcinolone injections, successive washes with autologous serum, and dexamethasone and tobramycin eye drops, as well as multiple cycles of oral antibiotics (doxycycline, amoxicillin-clavulanate, cotrimoxazole, and levofloxacin), the clinical course worsened. Shortly thereafter, the patient experienced palpebral rupture and copious purulent discharge. (Fig. 1).

Following the isolation of *Achromobacter xylosoxidans* from the exudate culture and confirmation of its susceptibility to ceftazidime, directed treatment with intravenous ceftazidime was administered, with no clinical improvement. Subsequently, a rapidly growing mycobacterium was identified in a pus exudate, preliminarily identified as *Mycobacterium abscessus* complex (Fig. 2). Therefore, hospital admission was decided to initiate intensive treatment after surgical debridement of the anophthalmic cavity.

Comprehensive orbital debridement was performed, extending to the periosteum in selected areas to ensure the complete removal of all necrotic tissue, which was notably abundant. This procedure ensured the exposure of vascularized tissue throughout the orbital locations. After the surgical intervention, complementary tests were conducted, revealing normal levels of C-reactive protein and procalcitonin. Additionally, computed tomography of the orbits showed no evidence of bone involvement or regional dissemination. Distant involvement was also excluded based on chest radiography and repeated physical examinations.

Pending identification at the subspecies level, empirical combined intravenous treatment was initiated with amikacin at 12.5 mg/kg (750 mg), imipenem 1 g every 8 hours, and tigecycline 50 mg every 12 hours, along with oral azithromycin 500 mg.

During the admission, the patient had digestive intolerance attributable to the use of tigecycline, which led to its suspension due to the onset of hepatitis. Due to the lack of availability of the antibiogram of the isolate, tigecycline was replaced with intravenous cefoxitin 2 g every 8 hours. At the patient's request, she was discharged home after 19 days of intravenous treatment, without yet having the definitive microbiological identification. Pending the results of phenotypic resistance, a quadruple oral therapy composed of azithromycin 500 mg every 24 h, moxifloxacin 400 mg every 24 h, linezolid 600 mg every 12 h, and rifabutin 300 mg every 24 h was maintained.

The patient showed good initial progress during outpatient follow-up. However, after receiving the final results of the microbiological sensitivity test, which revealed a multi-resistant strain, bedaquiline was requested for off-label use. Table 1. Treatment was maintained for 6 months with linezolid at a dose of 600 mg every 24 h, azithromycin 500




Fig. 1. Infection of the anophthalmic cavity with marked presence of purulent eschar, upper eyelid rupture, and periorbital cellulitis.




Fig. 2. Growth of rapidly growing mycobacterial colonies on Chocolate Agar, indicated by an arrowhead. Acid-fast bacteria, stained with Ziehl-Neelsen staining and highlighted by an asterisk, exhibit clustering in cords—a characteristic pattern common in *M. abscessus*. Detritus is also visible in the background.

Table 1

Amikacin MIC >64 µg/mL.	Resistant (A1408G mutation in rrs gene).
Cefoxitin MIC 32 µg/mL.	Susceptible with Increased Exposure.
Ciprofloxacin MIC >4 µg/mL.	Resistant.
Clarithromycin MIC 0.5 µg/mL.	Inducible resistance (T28 genotype in erm gene).
Doxycycline MIC >16 µg/mL.	Resistant.
Tigecycline MIC 0.5 µg/mL.	No breakpoints have been established by the CLSI.
Imipenem MIC 32 µg/mL.	Susceptible with Increased Exposure.
Linezolid MIC 16 µg/mL.	Susceptible with Increased Exposure.
Moxifloxacin MIC >8 µg/mL.	Resistant.
Trimethoprim/Sulfamethoxazole MIC >8/152 µg/mL.	Resistant
Tobramycin MIC >16 µg/mL.	Resistant.

Antimicrobial Susceptibility Testing for our isolate of *Mycobacterium abscessus* subsp. *abscessus*. Susceptibility interpretation according to M24-A2, ‘Susceptibility Testing of Mycobacteria, Nocardiae, and Other Aerobic Actinomycetes,’ as per the Clinical and Laboratory Standards Institute (2011).

mg every 24 h, and bedaquiline according to the usual schedule (400 mg every 24 h for the first two weeks, followed by 200 mg three times a week). Rifabutin was withdrawn due to hepatic toxicity.

The patient showed satisfactory clinical improvement during follow-up (Fig. 3). Once the treatment was completed, no signs of local recurrence were observed, allowing the placement of an ocular prosthesis with a good aesthetic result.

3. Discussion

The primary challenge in managing ophthalmic infections caused by non-tuberculous mycobacteria lies in the misidentification of the causative agent. However, with the advent of advanced diagnostic tools, such misidentifications are becoming less frequent. In our case, the repeated isolation of *A. xylosoxidans* initially led to an inappropriate treatment with ceftazidime, resulting in a delay in adopting the correct

therapeutic strategy. Clinicians should consistently consider the possibility of non-tuberculous mycobacteria in the differential diagnosis following treatment failure, especially when the initial treatment seems appropriate. This is crucial due to the potential for faster-growing microorganisms to obscure the presence of non-tuberculous mycobacteria. In our case, it is impossible to determine the exact moment when the *M. abscessus* infection occurred due to the poor progression following multiple orbital interventions, as well as the prolonged latency period described in the literature between surgical contamination and the development of infection by this microorganism.

Unlike pulmonary infections caused by rapidly growing mycobacteria, there is currently no standard treatment for ophthalmic infections caused by these pathogens. The literature consulted offers a limited and heterogeneous number of cases, reflecting the variety of ocular and annexed tissues susceptible to infection by these mycobacteria. Generally, cases have been managed with a combination of local antimicrobial treatment, systemic therapy, and surgical debridement.^{3,5,7,11–14} However, this strategy faces challenges such as the need for prolonged intravenous treatment, high economic cost, limited availability of oral options, and low tolerability of the drugs used.^{15,16}

Among the agents with proven activity against *M. abscessus*, only macrolides and amikacin have unequivocally demonstrated a relationship between in vitro activity and clinical efficacy.¹⁰ Macrolides are considered the cornerstone of treatment; however, most isolates show activity in the *erm* gene, which causes inducible resistance to these drugs. Most authors argue that amikacin is the second most important drug in a combined regimen. Other first-line drugs include tigecycline, imipenem, ceftazidime, and linezolid. There is sufficient evidence of activity, and these can be used as part of the treatment regimen: clofazimine, tedizolid, moxifloxacin or gatifloxacin, doxycycline or omadacycline, cotrimoxazole, rifabutin, and bedaquiline. There is scarce clinical evidence for the recommendation of combining beta-lactams, but there is evidence of synergy between imipenem with ceftazidime, ceftaroline and ceftazidime, as well as imipenem with the beta-lactamase inhibitors avibactam and relebactam.^{9,10,17–23}

The treatment is divided into an induction phase and a maintenance phase,

there was no availability of home-based intravenous antimicrobial therapy, resulting in a suboptimal oral regimen. After confirming the pathogen's multi-resistance, compassionate use of bedaquiline was authorized, initially in combination with rifabutin, based on the synergy described in animal studies. However, rifabutin was quickly discontinued due to digestive intolerance, hepatitis, and concerns about pharmacological interactions with linezolid, whose effectiveness was already compromised by the high MIC of the microbiological isolate.

In our case, bedaquiline was chosen for its nearly universal in vitro activity against all rapidly growing mycobacteria and the lack of suitable oral alternatives for our patient: omadacycline was not considered due to a history of tigecycline-induced hepatitis, and clofazimine was discarded because of its lower likelihood of sensitivity, given the reference laboratory's inability to provide a sensitivity study for both antimicrobials.

Bedaquiline has a unique mechanism of action, targeting the inhibition of the c subunit of ATP synthase in the genus *Mycobacterium* spp., and possesses bactericidal activity at any stage of bacillus replication. Although it is FDA-approved exclusively for the treatment of pulmonary infections by multidrug-resistant tuberculosis as part of a treatment regimen in the absence of other valid alternatives, bedaquiline also demonstrates high in vitro activity against a broad group of nontuberculous mycobacteria, including *M. avium*, *M. leprae*, and, notably, rapidly growing mycobacteria, where other treatment options are lacking.

Nevertheless, the use of bedaquiline should be approached with caution and strictly supervised by experienced physicians, such as infectious disease specialists or internists. In addition to not being approved for this specific indication, the drug has been linked to an increased mortality rate compared to placebo and requires monitoring for cardiac toxicity due to potential QT interval prolongation.

4. Conclusions

This case illustrates the complexity and challenges in the management of ocular infections by rapidly growing mycobacteria, highlighting the need for the development of specific treatment guidelines for these infections.

CRedit authorship contribution statement

David Galindo-Rodríguez: Writing – review & editing, Writing – original draft. **Miguel Moreno Hijazo:** Writing – review & editing, Writing – original draft. **Celina Balint Ilie:** Writing – review & editing. **Daniel Rubio Castro:** Writing – review & editing. **Ignacio Vallés Tormo:** Writing – review & editing. **Eva Gloria Alias Alegre:** Writing – review & editing, Writing – original draft.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Roussel TJ, Stern WH, Goodman DF, Whitcher JP. Postoperative mycobacterial endophthalmitis. *Am J Ophthalmol*. 1989;107(4):403–406. [https://doi.org/10.1016/0002-9394\(89\)90664-8](https://doi.org/10.1016/0002-9394(89)90664-8).
- Pinitpuwadol W, Tesavibul N, Boonsopon S, Sakiyalak D, Sarunket S, Choopong P. Nontuberculous mycobacterial endophthalmitis: case series and review of literature. *BMC Infect Dis*. 2020;20(1):877. <https://doi.org/10.1186/s12879-020-05606-2>.
- Hsu CR, Chen JT, Yeh KM, et al. A cluster of nontuberculous mycobacterial endophthalmitis (NTME) cases after cataract surgery: clinical features and treatment outcomes. *Early Years Educat*. 2018;32(9):1504–1511. <https://doi.org/10.1038/s41433-018-0108-1>.
- Paulose RM, Joseph J, Narayanan R, Sharma S. Clinical and microbiological profile of non-tuberculous mycobacterial endophthalmitis-experience in a tertiary eye care centre in Southern India. *J Ophthalmic Inflamm Infect*. 2016;6(1):27. <https://doi.org/10.1186/s12348-016-0096-x>.
- Hung JH, Ko WC, Chen CY, et al. Postoperative Mycobacteroides abscessus subsp. abscessus endophthalmitis: clinical analysis of 12 clustered adults and a proposed therapeutic algorithm. *J Microbiol Immunol Infect*. 2020;53(5):766–777. <https://doi.org/10.1016/j.jmii.2018.12.011>.
- Kheir WJ, Sheheili H, Abdul Fattah M, Hamam RN. Nontuberculous Mycobacterial ocular infections: a systematic review of the literature. *BioMed Res Int*. 2015;2015, 164989. <https://doi.org/10.1155/2015/164989>.
- Klapper SR, Patrinely JR, Kaplan SL, Font RL. Atypical mycobacterial infection of the orbit. *Ophthalmology*. 1995;102(10):1536–1541. [https://doi.org/10.1016/s0161-6420\(95\)30834-2](https://doi.org/10.1016/s0161-6420(95)30834-2).
- Matieli LCV, De Freitas D, Sampaio J, Moraes NSB, Yu MCZ, Hofling-Lima AL. Mycobacterium abscessus endophthalmitis: treatment dilemma and review of the literature. *Retina*. 2006;26(7):826–829. <https://doi.org/10.1097/01.iae.0000244276.80716.96>.
- Brown-Elliott BA, Wallace Jr RJ. In vitro susceptibility testing of bedaquiline against Mycobacterium abscessus complex. *Antimicrob Agents Chemother*. 2019;63(2). <https://doi.org/10.1128/AAC.01919-18>.
- Griffith DE, Daley CL. Treatment of Mycobacterium abscessus pulmonary disease. *Chest*. 2022;161(1):64–75. <https://doi.org/10.1016/j.chest.2021.07.035>.
- Kim AY, Davis AS, Moreau A, Drevets DA, Melendez DP. Management of nontuberculous mycobacterial infections of the eye and orbit: a retrospective case series. *Am J Ophthalmol Case Rep*. 2020;20(100971), 100971. <https://doi.org/10.1016/j.ajoc.2020.100971>.
- Moorthy RS, Valluri S, Rao NA. Nontuberculous mycobacterial ocular and adnexal infections. *Surv Ophthalmol*. 2012;57(3):202–235. <https://doi.org/10.1016/j.survophthal.2011.10.006>.
- Kuo CT, Chen YT, Tsai IL, Tsai CY, Chen ZTY. Orbital and conjunctival nontuberculous mycobacteria infection. *Taiwan J Ophthalmol*. 2023;13(3):393–397. <https://doi.org/10.4103/2211-5056.354534>.
- Zuercher D, Malet T. Mycobacterium abscessus infection in a case of recurrent orbital implant exposure. *Orbit*. 2007;26(4):337–339. <https://doi.org/10.1080/01676830600987573>.
- Jeon K, Kwon OJ, Lee NY, et al. Antibiotic treatment of Mycobacterium abscessus lung disease: a retrospective analysis of 65 patients. *Am J Respir Crit Care Med*. 2009;180(9):896–902. <https://doi.org/10.1164/rccm.200905-0704oc>.
- Novosad SA, Beekmann SE, Polgreen PM, Mackey K, Winthrop KL, M. abscessus Study Team. Treatment of Mycobacterium abscessus infection. *Emerg Infect Dis*. 2016;22(3):511. <https://doi.org/10.3201/eid2203.150828>.
- Daley CL, Iaccarino JM, Lange C, et al. Treatment of nontuberculous Mycobacterial pulmonary disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Clin Infect Dis*. 2020;71(4):e1–e36. <https://doi.org/10.1093/cid/ciaa241>.
- Phillely JV, Wallace Jr RJ, Benwill JL, et al. Preliminary results of bedaquiline as salvage therapy for patients with nontuberculous Mycobacterial lung disease. *Chest*. 2015;148(2):499–506. <https://doi.org/10.1378/chest.14-2764>.
- Dupont C, Viljoen A, Thomas S, et al. Bedaquiline inhibits the ATP synthase in Mycobacterium abscessus and is effective in infected zebrafish. *Antimicrob Agents Chemother*. 2017;61(11). <https://doi.org/10.1128/AAC.01225-17>.
- Johansen MD, Daher W, Roquet-Banères F, et al. Rifabutin is bactericidal against intracellular and extracellular forms of Mycobacterium abscessus. *Antimicrob Agents Chemother*. 2020;64(11). <https://doi.org/10.1128/AAC.00363-20>.
- Lee J, Ammerman N, Agarwal A, Naji M, Li SY, Nuermberger E. Differential in vitro activities of individual drugs and bedaquiline-rifabutin combinations against actively multiplying and nutrient-starved Mycobacterium abscessus. *Antimicrob Agents Chemother*. 2021;65(2). <https://doi.org/10.1128/aac.02179-20>.
- Pandey R, Chen L, Manca C, et al. Dual β -lactam combinations highly active against Mycobacterium abscessus complex in vitro. *mBio*. 2019;10(1). <https://doi.org/10.1128/mBio.02895-18>.
- Story-Roller E, Maggioncalda EC, Lamichhane G. Select β -lactam combinations exhibit synergy against Mycobacterium abscessus in vitro. *Antimicrob Agents Chemother*. 2019;63(4). <https://doi.org/10.1128/AAC.02613-18>.
- Griffith DE, Aksamit T, Brown-Elliott BA, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med*. 2007;175(4):367–416. <https://doi.org/10.1164/rccm.200604-571ST>.