#### REVIEW

Special Topic - CALCIUM SIGNALING

# Thirty years of Ca<sup>2+</sup> spark research: digital principle of cell signaling unveiled

Fujian Lu<sup>1</sup>, Pengcheng Yang<sup>2</sup>, Donghui Zhang<sup>2</sup>, Xianhua Wang<sup>3</sup>, Heping Cheng<sup>3</sup><sup>⊠</sup>

- <sup>1</sup> Department of Cardiology, Zhongshan Hospital, Institutes of Biomedical Sciences, Fudan University, Shanghai Institute of Cardiovascular Diseases, Shanghai 200032, China
- <sup>2</sup> State Key Laboratory of Biocatalysis and Enzyme Engineering, School of Life Science, Hubei University, Wuhan 430062, China
- <sup>3</sup> National Biomedical Imaging Center, State Key Laboratory of Membrane Biology, Peking-Tsinghua Center for Life Sciences, College of Future Technology, Peking University, Beijing 100871, China

Received: 17 July 2024 / Accepted: 19 September 2024

**Abstract** Calcium ions (Ca<sup>2+</sup>) are an archetypical and most versatile second messenger in virtually all cell types. Inspired by the discovery of Ca<sup>2+</sup> sparks in the 1990s, vibrant research over the last three decades has unveiled a constellation of Ca<sup>2+</sup> microdomains as elementary events of Ca<sup>2+</sup> signaling and, more importantly, a digital-analog dualism as the system design principle of Ca<sup>2+</sup> signaling. In this brief review, we present a sketchy summary on advances in the field of sparkology, and discuss how the digital subsystem can fulfill physiological roles otherwise impossible for any analog system. In addition, we attempt to address how the digital-analog dualism endows the simple cation messenger with signaling speediness, specificity, efficiency, stability, and unparalleled versatility.

**Keywords** Ca<sup>2+</sup> signaling, Ca<sup>2+</sup> microdomain, Digital system design

#### INTRODUCTION

Intracellular  $Ca^{2+}$  signaling is paradoxically simple and complex. One divalent cation connecting thousands of target proteins (or  $Ca^{2+}$  signalome),  $Ca^{2+}$  emerges as the most versatile intracellular messenger orchestrating a myriad of physiological processes from muscle contraction to neuronal firing, and from cell proliferation to apoptosis (Berridge *et al.* 2000, 2003; Bootman 2012). Yet, its powerful information-coding ability derives almost entirely from the binding and unbinding of this cation to its proteinaceous effectors as well as the electrical currents it generates when moving across a biological membrane. The enigma arises as to how  $Ca^{2+}$ orchestrates molecular players to fulfill remarkably diverse and sometimes opposing physiological functions within a given cell.

First discovered in 1993 with the advent of confocal microscopy and fast, high-contrast fluorescent probes, such as Fluo3 (Kao *et al.* 1989), Ca<sup>2+</sup> sparks (Fig.1) represent discrete, local and brief events elemental to intracellular  $Ca^{2+}$  signaling in the heart (Cheng *et al.* 1993). Since then, vibrant research in muscles, neurons, and other excitable and non-excitable cells has unveiled a large family of spark-like elemental Ca<sup>2+</sup> signaling events, an exquisite constellation of their subcellular organization, and unique and unexpected signaling modalities enabled by such Ca<sup>2+</sup> microdomains. These advances have led to the appreciation of the digital-analog dualism of the Ca<sup>2+</sup> signaling system (Cheng and Lederer 2008), with the digital subsystem encompassing all discrete, all-or-none signaling microdomains exemplified by Ca<sup>2+</sup> sparks, gaining teleological insights into diverse supramolecular Ca<sup>2+</sup> signaling substructures found in virtually all types of cells. In this brief review, based on the 2023 Bei Shizhang

<sup>🖂</sup> Correspondence: Chengp@pku.edu.cn (H. Cheng)



**Fig. 1**  $Ca^{2+}$  sparks. **A** Two-dimensional confocal images of  $Ca^{2+}$  sparks in a quiescent cardiac myocyte (scan rate 1.0 s/frame). **B** Linescan confocal images of an action potential-elicited  $[Ca^{2+}]_i$  transient (top) and a spontaneous spark (bottom) (scan rate 2.0 ms/line). Time and space ordinates are displayed in the horizontal and vertical directions, respectively (Cheng *et al.* 1993)

award lecture delivered by the senior author, we attempt to elucidate the digital building principle of the  $Ca^{2+}$  signaling system. Through a few examples, we discuss how the digital-analog dualism may unify the simplicity and the versatility of  $Ca^{2+}$  signaling.

#### Ca<sup>2+</sup> SIGNALING MICRODOMAINS

In multicellular organisms, Ca<sup>2+</sup> concentrations in the extracellular space and membrane-bounded intracellular Ca<sup>2+</sup> stores, primarily the endoplasmic reticulum (ER), are in the millimolar (mmol/L) range. However, sustained high Ca<sup>2+</sup> levels in the cytosol are extremely toxic, so resting cytosolic Ca<sup>2+</sup> must be maintained at low levels, typically in the range of tens to a few hundreds of nmol/L, implicating a 10,000-fold gradient across the plasma membrane and the membrane of the Ca<sup>2+</sup> stores. Signaling involves change, and the steep gradients are exploited to produce rapid, high Ca<sup>2+</sup> signals by mobilizing Ca<sup>2+</sup> from these reservoirs to the cytosol, coupled with re-sequestration mechanisms to timely terminate the signal. Ca<sup>2+</sup> ions are impermeable to lipid biomembranes, and their fast-rate transmembrane mobilization is mediated by various Ca<sup>2+</sup> channels that are gated by physical factors such as membrane voltage, mechanical stretch and temperature; ligands including neurotransmitters, inositol 1,4,5-trisphosphate (IP<sub>3</sub>), ryanodine, and  $Ca^{2+}$  itself; and even ER store depletion. Transmembrane Ca2+ mobilizers also include Ca<sup>2+</sup> pumps and exchangers, which operate at a slower rate (on a molecule-tomolecule comparison). While channel-mediated Ca<sup>2+</sup> flux always moves down-hill the electrochemical potential gradients of Ca<sup>2+</sup>, transporters can facilitate up-hill Ca<sup>2+</sup> movement at the cost of free energy input. From a cytosol-centric perspective, each Ca<sup>2+</sup> mobilizer acts as a Ca<sup>2+</sup> source when it injects Ca<sup>2+</sup> flux into the cytosolic compartment; and as a Ca<sup>2+</sup> sink when it removes Ca<sup>2+</sup> ions to other membrane-partitioned compartment. In the cytosol milieu rich with Ca<sup>2+</sup> binding sites, the bound-to-free Ca<sup>2+</sup> ratio reaches about 100. Ca<sup>2+</sup> diffusion is slowed by an order of magnitude, and the spatial range of Ca<sup>2+</sup> signaling is also locally confined. This allows Ca<sup>2+</sup> signals at a few µm distances to operate quasi-independently. From the source-sink viewpoint, a  $Ca^{2+}$  binding site (or  $Ca^{2+}$  buffer) can function as both a source and a sink, depending on when and where it discharges or mops up free Ca<sup>2+</sup> ions. Furthermore, there exist many nanoscale and micro-scale membranous substructures, in the form of membrane-apposed cleft, button, vesicle, cistern, tubule, filopodium and lamellipodium, etc. In such subspaces, Ca<sup>2+</sup> handling is greatly influenced by physical constraints imposed by the membrane boundaries.

In the common-pool model of intracellular  $Ca^{2+}$  signaling, cytosolic  $Ca^{2+}$  is theorized to be spatially homogenous, and all effectors are governed by the same intracellular  $Ca^{2+}$  dynamics (Stern 1992). However, the very existence of discrete  $Ca^{2+}$  sparks that span only ~2 µm diameter and last about 20 ms (Cheng and Lederer 2008) has provided direct evidence that this is a significant oversimplification. In mammalian ventricular myocytes, cardiac dyads are supramolecular nanoscopic substructures formed by apposition of transverse (T) tubules (*i.e.*, invaginations of the plasma membrane) and cisterns of the ER/sarcoplasmic reticulum (SR) (Franzini-Armstrong *et al.* 1999; Lu and Pu 2020). Excitation-contraction (E-C) coupling links membrane depolarization to myocyte contraction, which

occurs primarily at cardiac dyads via the Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release (CICR) mechanism (Fabiato and Fabiato 1972; Fabiato et al. 1972). Specifically, membrane depolarization opens voltage-gated L-type Ca<sup>2+</sup> channels (LTCCs) on the T-tubules, and resultant  $Ca^{2+}$  influx rushes into the dyadic cleft and serves as the activator of an array of type 2 ryanodine receptor (RYR2) Ca<sup>2+</sup> release channels, liberating SR store Ca<sup>2+</sup> and giving rise to  $Ca^{2+}$  sparks. In a typical cardiac myocyte, a constellation of  $Ca^{2+}$  sparks arising from ~10,000 dyads summates to produce a cell-wide Ca<sup>2+</sup> transients that activates cell contraction. Essentially, cardiac E-C coupling reflects the behavior of a digital system in which Ca<sup>2+</sup> sparks serve as the elementary units. These unitary Ca<sup>2+</sup> sparks are locally controlled and digitally activated, providing a teleological explanation for salient architectural features such as the nanoscale dyadic cleft and microscale spatial separation of dyads, which align with the similarly sized length of a sarcomere. This alignment, possibly not coincidental, matches the dimension of sparks.

The digital mode of Ca<sup>2+</sup> signaling is universal to virtually all cell types examined, and a family of elementary Ca<sup>2+</sup> signaling events has been characterized in both excitable and non-excitable cells (Fig.2). In particular, Ca<sup>2+</sup> sparks are present in skeletal (Baylor *et al.* 2002; Hollingworth et al. 2001; Klein et al. 1996; Tsugorka et al. 1995) and smooth muscle myocytes (Burdyga and Wray 2005; Nelson *et al.* 1995; ZhuGe *et al.* 1998), neuroendocrine cells (e.g., chromaffin cells) (ZhuGe et al. 2006), and neurons containing different isoforms of RYRs (De Crescenzo et al. 2004; Koizumi et al. 1999; Ouyang et al. 2005). In the nanoscale dyadic cleft, a Ca<sup>2+</sup> nanospark with a higher Ca<sup>2+</sup> level has been visualized by using dyad-targeted Ca<sup>2+</sup> sensors GCaMP6f-Triadin1/ Junctin (Shang *et al.* 2014). Ca<sup>2+</sup> sparklet arising from a plasmalemmal LTCC (Wang et al. 2001) or TRPV4 channel (Sonkusare et al. 2012) has been optically recorded in cardiac or smooth muscle cells, respectively. Inside the Ca<sup>2+</sup> store, local depletion of Ca<sup>2+</sup> occurs concurrently during a Ca<sup>2+</sup> spark, and this phenomenon of Ca<sup>2+</sup> depletion has been visualized as a Ca<sup>2+</sup> blink (Lu et al. 2020), and is thought to provide a possible mechanism to signal spark termination (Brochet et al. 2005) and local activation of store-operated Ca<sup>2+</sup> entry (SOCE), manifested as Ca<sup>2+</sup> glows in the invadopodium of cancer cells during metastasis (Lu et al. 2019). In oocytes, discrete local Ca<sup>2+</sup> release of IP<sub>3</sub> receptor (IP<sub>3</sub>R) origin gives rise to Ca<sup>2+</sup> puffs (from clustered IP<sub>3</sub>Rs) and blips (from single IP<sub>3</sub>Rs) (Lock et al. 2019; Yao and Parker 1994). In human lung fibroblasts, discrete, local and short-lived high Ca<sup>2+</sup> microdomains, namely Ca<sup>2+</sup> flickers (Wei et al. 2009), are representative of a converging multimodal signaling. The source Ca<sup>2+</sup> of a flicker comes from both TRPM7, which senses plasmalemmal mechanical stress, and IP<sub>3</sub>R, which is co-activated by local Ca<sup>2+</sup> and IP<sub>3</sub>, the latter being linked to extracellular chemotactic signals. More recently, it has been shown that high Ca<sup>2+</sup> microdomains on the ER surface trigger liquid-liquid phase separation to specify autophagosome initiation sites (Zheng et al. 2022). Perinuclear  $Ca^{2+}$  waves in the nuclear envelope (Luo *et al.* 2008), with high  $Ca^{2+}$  passing through the nuclear pores, can preferentially access the matrix of the nucleus and regulate Ca<sup>2+</sup>-dependent biological processes within. The Golgi apparatus serves also as a reservoir for Ca<sup>2+</sup> in the cell.  $Ca^{2+}$  release from the Golgi  $Ca^{2+}$  channel can be triggered by various signaling pathways and stimuli, and create local high Ca<sup>2+</sup> signals influencing cellular responses such as secretion and cell growth (Pizzo et al. 2011). Overall, it is safe to conclude that each cell type harbors unique nanoscale supramolecular substructures that support dynamic microdomain Ca<sup>2+</sup> signaling. Their location and mode of action are tailored to the specific needs of vital physiological functions.

## DIGITAL PRINCIPLE OF Ca<sup>2+</sup> SIGNALING

The universality of Ca<sup>2+</sup> microdomains and their unique mode of action have led to the appreciation of a digitalanalog dualism for cell Ca2+ signaling (Cheng and Lederer 2008), in which discrete Ca<sup>2+</sup> signaling microdomains form a digital subsystem that is intertwined with an analog subsystem comprising  $Ca^{2+}$  signals in the bulk cytosol. Specifically, the digital subsystem is built for high-threshold Ca<sup>2+</sup>-dependent processes, such as activation of  $BK_{Ca}$  and  $Cl_{Ca}$ , activation of CaMKII and calcineurin, and vesicle secretion (Ouyang et al. 2005), all of which require  $Ca^{2+}$  levels to reach tens of  $\mu$ mol/L and beyond. This subsystem is discontinuous in space and individual microdomains operate intermittently in an all-or-none fashion. Its overall Ca<sup>2+</sup> signaling effect is thus depicted by a digital function,  $S(n, t_{event})$ , where *n* and tevent denote the number and timing of microdomain activation. Spatially and temporally patterned activation of the digital subsystem may enable intracellular Ca<sup>2+</sup> signaling to achieve roles that are otherwise impossible for any analog system. In the following we briefly discuss sub-principles of digital Ca<sup>2+</sup> signaling as illustrated by examples in different types of cells or physiological contexts.



**Fig. 2** Exemplary  $Ca^{2+}$  signaling microdomains. Cartoon illustration of various microdomains featured in different types of cells, all being artificially integrated in a hypothetical single cell. (1) Cardiac dyad. Single LTCC openings generate  $Ca^{2+}$  sparklets to activate a RYR2  $Ca^{2+}$  nanospark in the cleft and a  $Ca^{2+}$  spark of 2 µm diameter in the surrounding space, as well as a  $Ca^{2+}$  blink in the connected ER/SR cistern. (2) Skeletal muscle triad (half structure illustrated). Arrayed RYR1 is mechanically gated by DHPR acting as the voltage sensor to initiate SR  $Ca^{2+}$  release, which, in turn, may activate nearby RYR3/ $\beta$  array via the CICR mechanism. (3) Subsurface cistern, to support subsurface sparks seen in smooth muscle cells and DRG neurons. (4) Clustered Orai1 and STIM1 forming an elemental unit for SOCE. Its digital operation gives rise to  $Ca^{2+}$  glows which morphology and kinetics are shaped by membranous structures like invadopodia. (5) Multimodal  $Ca^{2+}$  signaling microdomain. In human lung fibroblasts,  $Ca^{2+}$  flickers arise from both TRPM7  $Ca^{2+}$  influx, sensing mechanical signals, and IP<sub>3</sub>R  $Ca^{2+}$  release, sensing chemotactic signal mediated by platelet-derived growth factor receptor (PDGFR) signaling pathway, and dynamically regulate the assembly and disassembly of cytoskeletal filaments, and thereby steer the direction of cell migration. (6) Mitochondria-associated membranes (MAMs), the contact sites between mitochondria and ER, are a central hub for  $Ca^{2+}$  signaling, apoptosis, autophagy and lipid biosynthesis. (7) ER surface  $Ca^{2+}$  microdomains, sites of phase separation of protein complex for specification of autophagosome initiation. (8) Perinuclear  $Ca^{2+}$  waves from the ER in the nuclear envelope, with high  $Ca^{2+}$  passing through the nuclear pores. (9) Golgi  $Ca^{2+}$  microdomain

## Linear and stable control of an inherently nonlinear unstable system

CICR implies positive feedback and an inherently strong non-linearity, best suited for signaling rapidity and high-gain amplification. Theoretically, when the gain of CICR is greater than one, a common-pool system would become unstable such that the system as a whole operates in the all-or-none regime (Stern 1992). In fact, as informed from the whole-cell Ca<sup>2+</sup> transient and contraction strength, cardiac E-C coupling driven by CICR is gradable and tightly controlled by the duration and the magnitude of the triggering LTCC  $Ca^{2+}$ current, creating the so-called CICR paradox. In the digital  $Ca^{2+}$  signaling regime, however, the paradox is readily resolved. At the system level, global  $Ca^{2+}$  transient is, to the first order approximation, a linear function of the number of the unitary  $Ca^{2+}$  sparks evoked. The positive feedback and high-gain amplification are largely confined within individual dyads. Nevertheless, instability does occur when a cardiac myocyte is challenged with  $Ca^{2+}$ -overloading conditions. In this scenario, E-C coupling is no longer graded by LTCC triggering  $Ca^{2+}$ ,  $Ca^{2+}$  sparks tend to ignite neighboring dyads, and regenerative inter-dyadic CICR would result in propagating  $Ca^{2+}$  waves that are highly arrhythmogenic (Cheng *et al.* 1996).

### Enhanced efficiency and specificity of Ca<sup>2+</sup> signaling

This sub-principle applies to all  $Ca^{2+}$  signaling processes whose effects depend hyper-linearly on  $Ca^{2+}$  concentration. Given the same total  $Ca^{2+}$  mobilized, spatial and temporal condensation would disproportionally enhance their signaling efficiency. For instance, in a process with a power of two  $Ca^{2+}$ -dependence, the overall effect would be augmented as much as 100 times, if its spatial and temporal dimensions are each compressed by a factor of 10 (assuming effectors are uniformly distributed spatially). In this regard, we have shown that about seven subsurface  $Ca^{2+}$  sparks suffice to trigger a vesicle release in DRG neurons (Ouyang *et al.* 2005), while the amount of  $Ca^{2+}$  involved could barely raise  $Ca^{2+}$  level by a few nmol/L if uniformly distributed.

An additional advantage of a digital organization is related to signaling specificity. This is because the signal is turned on only at precise locations with specific timing. Ample examples can testify to this point. During melanoma invasion, SOCE-mediated Ca<sup>2+</sup> glows promote CaM/Pyk2 aggregation and matrix metalloproteinase secretion in invadopodia (Lu *et al.* 2019). High Ca<sup>2+</sup> on the ER surface can selectively activate phase separation of FIP200 for specification of autophagosome initiation sites on the ER (Zheng *et al.* 2022). By confining the high-Ca<sup>2+</sup> process at sites of MAMs (Vance 2014), inter-organelle Ca<sup>2+</sup> signaling can be channeled without affecting effectors in the cytosol.

# Microdomain $Ca^{2+}$ fulfills roles distinct from that of global $Ca^{2+}$

An elegant example came from the study of  $Ca^{2+}$  spark regulation of vascular tone. In arterial smooth muscles, while global  $Ca^{2+}$  elevation signifies muscle contraction, subsurface  $Ca^{2+}$  sparks relax the blood vessels. Mechanistically, high subsurface  $Ca^{2+}$  activates highthreshold BK<sub>Ca</sub> to cause membrane hyperpolarization, shut-off of LTCC  $Ca^{2+}$  influx and eventual decrease of bulk cytosolic  $Ca^{2+}$  concentration (Nelson *et al.* 1995).

# Complex behavior emerged from coordinating microdomain Ca<sup>2+</sup> activity

In migrating cells,  $Ca^{2+}$  plays multifunctional roles in directional sensing, cytoskeleton redistribution, traction force generation, and relocation of focal adhesions. Multimodal  $Ca^{2+}$  signaling microdomains are present at the front of migrating fibroblasts, with TRPM7 sensing mechanical signals and PDGFR sensing gradients of chemotactic signals. These signals converge onto plasmalemmal TRPM7- subspace  $IP_3R$  clusters to generate  $Ca^{2+}$  flickers. Spatial and temporal coordination of these digital events by both mechanical and chemotaxis signals is thought to orchestrate the assembly and disassembly of cytoskeletal filaments and thus direct cell migration (Wei *et al.* 2009).

#### **NEW HORIZONS**

Unraveling Ca<sup>2+</sup> signaling microdomains and the digital-analog dualism represents a paradigm shift in our understanding of the design principle of cellular Ca<sup>2+</sup> signaling system as a whole. Deeper insights into Ca<sup>2+</sup> signaling in both physiology and pathophysiology have also been obtained through elucidating molecular mechanisms, physiological significance, and pathological implications of Ca<sup>2+</sup> microdomains. However, many important unmet challenges lie ahead. First, the assembly of Ca<sup>2+</sup> microdomains remains poorly understood. We are only beginning to appreciate the complex mechanism underlying the formation and positioning of the supramolecular assembly of Ca<sup>2+</sup> microdomains. In heart, how do dyads as the hallmark architecture establish and maintain their striking registration along with sarcomeric Z-lines? Our recent work on in situ and in vivo proteomic mapping of dyadic proteins revealed that CMYA5, a master protein localized to Z-lines, contributes to tethering junctional SR (jSR) adjacent to these structures. T-tubules subsequently form and colocalize with jSR, yielding organized, properly positioned dyads (Lu et al. 2022), in which junctophilin family proteins play important roles (Hall et al. 2024). Second, at the system level, matching a plethora of Ca<sup>2+</sup> signalome proteins of different Ca<sup>2+</sup> affinity and functionality with a constellation of Ca<sup>2+</sup> microdomains of distinctive characteristics must be a daunting task, letting alone the combinatorial interaction among them. Developing innovative approaches to systematically elucidate the sorting mechanism and its underlying cell logic would yield profound insights into the digitalanalog dualism of cell signaling. Third, there are urgent needs to discriminate dysregulation of microdomain Ca<sup>2+</sup> signaling in a wide range of diseases (Gomez *et al.* 1997; Song et al. 2006). Advanced imaging techniques, such as three-dimensional electron microscope reconstruction and super-resolution light microscopy, coupled with new generations of microdomain-targeted probes, offer powerful means to uncover new architectures, new roles and new modes of microdomain Ca<sup>2+</sup>

signaling *in vitro* and *in vivo*, as well as their subtle changes in pathological conditions. In addition, computational modeling approaches are also valuable in integrating quantitative cell  $Ca^{2+}$  data into a unifying framework.

Given the archetypical role of Ca<sup>2+</sup> as a second messenger, future research should systematically investigate whether the digital principle is applicable to other common biological messengers such as cAMP, IP<sub>3</sub>, nitric oxide, and reactive oxygen species (ROS). In this regard, previous studies have shown that cAMP signaling is highly compartmentalized during  $\beta_2$ -adrenergic stimulation (Kuschel et al. 1999; Mika et al. 2012) and that cAMP signaling microdomains selectively modulate intra-dvadic, but spare extra-dvadic targets including the Ca<sup>2+</sup> pump SERCA2a in cardiac myocytes (Zhou et al. 2009). As another example, single-mitochondrion ROS bursts during multi-faceted signaling events known as "mitoflashes" (Wang et al. 2008) operate in a frequency-dependent, digital mode, and play important roles in the regulation of cardiac ATP homeostasis (Wang et al. 2017), synaptic plasticity (Fu et al. 2017), and brown adipocyte thermogenesis (Chen et al. 2024). Altogether, we are witnessing the emergence of a fundamental principle of cell physiology: the digital organization of signaling microdomains is crucial for achieving rapid, specific, efficient, stable, and unparallelly diverse cell signaling.

Acknowledgements We are grateful to Drs. L.S. Song and S.Q. Wang for valuable discussions on the concepts presented in this work. We are also in debt to our collaborators, colleagues and students whose collective contribution has laid the foundation of the field of sparkology. This work is supported by the National Natural Science Foundation of China (32293211, 61972009, 92157105, 81827809) (to HC) and JIH1340085Y (to FL).

#### **Compliance with Ethical Standards**

**Conflict of interest** Fujian Lu, Pengcheng Yang, Donghui Zhang, Xianhua Wang, and Heping Cheng declare that they have no conflict of interest.

**Human and animal rights and informed consent** This article does not contain any studies with human or animal subjects performed by any of the authors.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

#### References

- Baylor SM, Hollingworth S, Chandler WK (2002) Comparison of simulated and measured calcium sparks in intact skeletal muscle fibers of the frog. J Gen Physiol 120(3): 349–368
- Berridge MJ, Bootman MD, Roderick HL (2003) Calcium signalling: dynamics, homeostasis and remodelling. Nat Rev Mol Cell Biol 4(7): 517–529
- Berridge MJ, Lipp P, Bootman MD (2000) The versatility and universality of calcium signalling. Nat Rev Mol Cell Biol 1(1): 11–21
- Bootman MD (2012) Calcium signaling. Cold Spring Harb Perspect Biol 4(7): a011171. https://doi.org/10.1101/cshperspect. a011171
- Brochet DX, Yang D, Di Maio A, Lederer WJ, Franzini-Armstrong C, Cheng H (2005) Ca<sup>2+</sup> blinks: rapid nanoscopic store calcium signaling. Proc Natl Acad Sci USA 102(8): 3099–3104
- Burdyga T, Wray S (2005) Action potential refractory period in ureter smooth muscle is set by Ca sparks and BK channels. Nature 436(7050): 559–562
- Chen X, Bahetiyaer H, Song X, Jiang Z, Qi W, Gao W, Zhang L, Zhang J, Cheng H, Wang X (2024) Mitochondrial flashes are interlinked with adaptive thermogenesis in brown adipocytes. hLife 2(4): 179–188
- Cheng H, Lederer MR, Lederer WJ, Cannell MB (1996) Calcium sparks and  $[Ca^{2+}]_i$  waves in cardiac myocytes. Am J Physiol 270(1 Pt 1): C148–159
- Cheng H, Lederer WJ (2008) Calcium sparks. Physiol Rev 88(4): 1491–1545
- Cheng H, Lederer WJ, Cannell MB (1993) Calcium sparks: elementary events underlying excitation-contraction coupling in heart muscle. Science 262(5134): 740–744
- De Crescenzo V, ZhuGe R, Velazquez-Marrero C, Lifshitz LM, Custer E, Carmichael J, Lai FA, Tuft RA, Fogarty KE, Lemos JR, Walsh JV, Jr. (2004) Ca<sup>2+</sup> syntillas, miniature Ca<sup>2+</sup> release events in terminals of hypothalamic neurons, are increased in frequency by depolarization in the absence of Ca<sup>2+</sup> influx. J Neurosci 24(5): 1226–1235
- Fabiato A, Fabiato F (1972) Excitation-contraction coupling of isolated cardiac fibers with disrupted or closed sarcolemmas. Calcium-dependent cyclic and tonic contractions. Circ Res 31(3): 293–307
- Fabiato F, Fabiato A, Sonnenblick EH (1972) Regenerative calcium liberation by the sarcoplasmic reticulum on skeletal and cardiac muscle cells. J Physiol (Paris) 65(Suppl 1): 118A
- Franzini-Armstrong C, Protasi F, Ramesh V (1999) Shape, size, and distribution of Ca(2+) release units and couplons in skeletal and cardiac muscles. Biophys J 77(3): 1528–1539
- Fu ZX, Tan X, Fang H, Lau PM, Wang X, Cheng H, Bi GQ (2017) Dendritic mitoflash as a putative signal for stabilizing longterm synaptic plasticity. Nat Commun 8(1): 31. https://doi.org/ 10.1038/s41467-017-00043-3
- Gomez AM, Valdivia HH, Cheng H, Lederer MR, Santana LF, Cannell MB, McCune SA, Altschuld RA, Lederer WJ (1997) Defective excitation-contraction coupling in experimental cardiac hypertrophy and heart failure. Science 276(5313): 800–806
- Hall DD, Takeshima H, Song LS (2024) Structure, function, and regulation of the junctophilin family. Annu Rev Physiol 86:

123-147

- Hollingworth S, Peet J, Chandler WK, Baylor SM (2001) Calcium sparks in intact skeletal muscle fibers of the frog. J Gen Physiol 118(6): 653–678
- Kao JP, Harootunian AT, Tsien RY (1989) Photochemically generated cytosolic calcium pulses and their detection by fluo-3. J Biol Chem 264(14): 8179–8184
- Klein MG, Cheng H, Santana LF, Jiang YH, Lederer WJ, Schneider MF (1996) Two mechanisms of quantized calcium release in skeletal muscle. Nature 379(6564): 455–458
- Koizumi S, Bootman MD, Bobanovic LK, Schell MJ, Berridge MJ, Lipp P (1999) Characterization of elementary Ca<sup>2+</sup> release signals in NGF-differentiated PC12 cells and hippocampal neurons. Neuron 22(1): 125–137
- Kuschel M, Zhou YY, Spurgeon HA, Bartel S, Karczewski P, Zhang SJ, Krause EG, Lakatta EG, Xiao RP (1999) beta2-adrenergic cAMP signaling is uncoupled from phosphorylation of cytoplasmic proteins in canine heart. Circulation 99(18): 2458–2465
- Lock JT, Smith IF, Parker I (2019) Spatial-temporal patterning of Ca(2+) signals by the subcellular distribution of IP(3) and IP(3) receptors. Semin Cell Dev Biol 94: 3–10
- Lu F, Ma Q, Xie W, Liou CL, Zhang D, Sweat ME, Jardin BD, Naya FJ, Guo Y, Cheng H, Pu WT (2022) CMYA5 establishes cardiac dyad architecture and positioning. Nat Commun 13(1): 2185. https://doi.org/10.1038/s41467-022-29902-4
- Lu F, Pu WT (2020) The architecture and function of cardiac dyads. Biophys Rev 12(4): 1007–1017
- Lu F, Sun J, Zheng Q, Li J, Hu Y, Yu P, He H, Zhao Y, Wang X, Yang S, Cheng H (2019) Imaging elemental events of store-operated Ca(2+) entry in invading cancer cells with plasmalemmal targeted sensors. J Cell Sci 132(6): jcs224923. https://doi.org/ 10.1242/jcs.224923
- Lu F, Zhao Y, Xie W, Guo Q, Wang SQ, Wang X, Cheng H (2020) Imaging sarcoplasmic reticulum Ca(2+) signaling in intact cardiac myocytes. Circulation 142(15): 1503–1505
- Luo D, Yang D, Lan X, Li K, Li X, Chen J, Zhang Y, Xiao RP, Han Q, Cheng H (2008) Nuclear Ca<sup>2+</sup> sparks and waves mediated by inositol 1, 4, 5-trisphosphate receptors in neonatal rat cardiomyocytes. Cell Calcium 43(2): 165–174
- Mika D, Leroy J, Vandecasteele G, Fischmeister R (2012) PDEs create local domains of cAMP signaling. J Mol Cell Cardiol 52(2): 323–329
- Nelson MT, Cheng H, Rubart M, Santana LF, Bonev AD, Knot HJ, Lederer WJ (1995) Relaxation of arterial smooth muscle by calcium sparks. Science 270(5236): 633–637
- Ouyang K, Zheng H, Qin X, Zhang C, Yang D, Wang X, Wu C, Zhou Z, Cheng H (2005) Ca<sup>2+</sup> sparks and secretion in dorsal root ganglion neurons. Proc Natl Acad Sci USA 102(34): 12259–12264
- Pizzo P, Lissandron V, Capitanio P, Pozzan T (2011) Ca(2+) signalling in the Golgi apparatus. Cell Calcium 50(2): 184–192
- Shang W, Lu F, Sun T, Xu J, Li LL, Wang Y, Wang G, Chen L, Wang X, Cannell MB, Wang SQ, Cheng H (2014) Imaging Ca<sup>2+</sup>

- Song LS, Sobie EA, McCulle S, Lederer WJ, Balke CW, Cheng H (2006) Orphaned ryanodine receptors in the failing heart. Proc Natl Acad Sci USA 103(11): 4305–4310
- Sonkusare SK, Bonev AD, Ledoux J, Liedtke W, Kotlikoff MI, Heppner TJ, Hill-Eubanks DC, Nelson MT (2012) Elementary Ca<sup>2+</sup> signals through endothelial TRPV4 channels regulate vascular function. Science 336(6081): 597–601
- Stern MD (1992) Theory of excitation-contraction coupling in cardiac muscle. Biophys J 63(2): 497–517
- Tsugorka A, Rios E, Blatter LA (1995) Imaging elementary events of calcium release in skeletal muscle cells. Science 269(5231): 1723–1726
- Vance JE (2014) MAM (mitochondria-associated membranes) in mammalian cells: lipids and beyond. Biochim Biophys Acta 1841(4): 595–609
- Wang SQ, Song LS, Lakatta EG, Cheng H (2001) Ca<sup>2+</sup> signalling between single L-type Ca<sup>2+</sup> channels and ryanodine receptors in heart cells. Nature 410(6828): 592–596
- Wang W, Fang H, Groom L, Cheng A, Zhang W, Liu J, Wang X, Li K, Han P, Zheng M, Yin J, Wang W, Mattson MP, Kao JP, Lakatta EG, Sheu SS, Ouyang K, Chen J, Dirksen RT, Cheng H (2008) Superoxide flashes in single mitochondria. Cell 134(2): 279–290
- Wang X, Zhang X, Wu D, Huang Z, Hou T, Jian C, Yu P, Lu F, Zhang R, Sun T, Li J, Qi W, Wang Y, Gao F, Cheng H (2017) Mitochondrial flashes regulate ATP homeostasis in the heart. eLife 6: e23908. https://doi.org/10.7554/eLife.23908
- Wei C, Wang X, Chen M, Ouyang K, Song LS, Cheng H (2009) Calcium flickers steer cell migration. Nature 457(7231): 901–905
- Yao Y, Parker I (1994) Ca<sup>2+</sup> influx modulation of temporal and spatial patterns of inositol trisphosphate-mediated Ca<sup>2+</sup> liberation in Xenopus oocytes. J Physiol 476(1): 17–28
- Zheng Q, Chen Y, Chen D, Zhao H, Feng Y, Meng Q, Zhao Y, Zhang H (2022) Calcium transients on the ER surface trigger liquidliquid phase separation of FIP200 to specify autophagosome initiation sites. Cell 185(22): 4082–4098
- Zhou P, Zhao YT, Guo YB, Xu SM, Bai SH, Lakatta EG, Cheng H, Hao XM, Wang SQ (2009) Beta-adrenergic signaling accelerates and synchronizes cardiac ryanodine receptor response to a single L-type Ca<sup>2+</sup> channel. Proc Natl Acad Sci USA 106(42): 18028–18033
- ZhuGe R, DeCrescenzo V, Sorrentino V, Lai FA, Tuft RA, Lifshitz LM, Lemos JR, Smith C, Fogarty KE, Walsh JV (2006) Syntillas release Ca<sup>2+</sup> at a site different from the microdomain where exocytosis occurs in mouse chromaffin cells. Biophys J 90(6): 2027–2037
- ZhuGe R, Sims SM, Tuft RA, Fogarty KE, Walsh JV, Jr. (1998) Ca<sup>2+</sup> sparks activate K<sup>+</sup> and Cl<sup>-</sup> channels, resulting in spontaneous transient currents in guinea-pig tracheal myocytes. J Physiol 513(Pt 3): 711–718