original article

# Prospective evaluation of serum and peritoneal fluid markers as indicators of postoperative complications in patients with enteric anastomosis

Mladen Kasalović,[a] B Surgical Clinic, University Clinical Center Kragujevac, Kra

[d]Laboratory Diagnostics Service, Clinical Hospital Center, Kosovska Mitrovica, Serbia; [e]Faculty of Medicine, University of Pristina, Medical Statistics and Informatics, Kosovska Mitrovica, Serbia

**Correspondence:** Dr. Mladen Novica Kasalović · Surgical Clinic, Clinical Hospital Center Kosovska Mitrovica, Kosovska Mitrovica, Serbia · kasalovicm2105@gmail.com · ORCID: ttps://orcid.org/0000-0002-6634-1380

**Citation:** Kasalović M, Odalović B, Mihajlović L, Jakovljević S, Elek Z, Igrutinović G, et al. Prospective evaluation of serum and peritoneal fluid markers as indicators of post-operative complications in patients with enteric anastomosis. Ann Saudi Med 2024; 44(6): 422-434. DOI: 10.5144/0256-4947.2024.422

**Received:** July 24, 2024

**Accepted:** October 27, 2024

**Published:** December 5, 2024

**Copyright:** Copyright © 2024, Annals of Saudi Medicine, Saudi Arabia. This is an open access article under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND). The details of which can be accessed at http://creativecommons. org/licenses/by-nc-nd/4.0/

**Funding:** None.

**BACKGROUND:** Postoperative complications of colorectal cancer surgery contribute to increased morbidity and mortality in patients.

**OBJECTIVES:** Investigate the role of biochemical markers in serum and peritoneal fluid in the development of postoperative complications in patients with enteral anastomosis.

**DESIGN:** Prospective.

**SETTING:** University hospitals.

**PATIENTS AND METHODS:** The studied population consisted of patients who underwent surgical treatment with created anastomosis or Hartmann's resection from April 2022 to April 2024, conducted at the Clinical-Hospital Center Kosovska Mitrovica and the University Clinical Center Kragujevac. Spearman's correlation coefficient ($r_s$) was used to test associations between categorical variables.

**MAIN OUTCOME MEASURES:** Lactate, albumin, lactate dehydrogenase, and IgA antibodies were monitored as predictors of anastomotic dehiscence and general postoperative complications.

**SAMPLE SIZE:** 52

**RESULTS:** The concentration of lactate in the drain fluid on the third postoperative day was statistically significantly higher in patients who did not develop anastomotic dehiscence ($P$=.006). The concentration of IgA antibodies in the drain fluid on the third and fifth days post-surgery showed a moderate negative correlation with lactate concentration ($r_s$=-.670, $P$=.012; $r_s$=-.577, $P$=.039), repectively. There was a significantly higher concentration of albumin in the drain fluid on the third day post-surgery in patients who developed dehiscence ($P$=.040), and on the seventh day post-surgery in those who did not develop dehiscence ($P$=.001). The concentration of LDH on the third day in the drain fluid after surgery was statistically significantly higher in patients who did not develop dehiscence ($P$=.020). There was a statistically significant difference in lactate concentration in the drain fluid on the third ($P$<.001) and fifth days ($P$=.041) post-surgery, as well as in albumin concentration on the third day post-surgery ($P$=.024) with respect to the development of general postoperative complications.

**CONCLUSION:** This study revealed significant differences in the concentrations of lactate, albumin, and LDH in the drain fluid on the third and fifth days post-surgery with respect to the development of complications. These results suggest that monitoring these markers may

help in the early identification of patients at risk of complications such as dehiscence.
**LIMITATIONS:** Limited literature on specific aspects of this study, including the absence of a control group, small sample size, and two-center study.
**CONFLICT OF INTEREST:** None.

Colorectal cancer exhibits various inheritance patterns, including sporadic, familial, and hereditary forms. Diet, physical activity, smoking, and alcohol consumption are factors that can influence the risk of colorectal cancer.[1] Characteristic symptoms of colorectal cancer include changes in bowel habits, blood in the stool, pain, weight loss, and anemia.[1] After a right hemicolectomy, an L-L anastomosis is recommended, particularly for colorectal adenocarcinoma.[2] However, primary resection and anastomosis in patients with colorectal cancer obstruction require complex surgical techniques and may be associated with high mortality and morbidity rates.[3] Recent years of research emphasize the importance of the increased systemic inflammatory response being associated with reduced survival in patients with colorectal cancer, regardless of tumor stage.[4]

Hartmann's procedure, which includes rectosigmoid resection, closure of the rectal stump, and formation of a colostomy, is applied in cases of obstruction or perforation of the left side of the colon.[5] Studies comparing postoperative outcomes of Hartmann's resection and resection with primary anastomosis suggest similar mortality and morbidity rates.[2] Treatment of Meckel's diverticulum usually involves surgical resection of the small intestine, often with end-to-end anastomosis.[3] Gastrointestinal stromal tumors (GISTs) require radical resection, especially if the tumor is larger than 2 cm.[6]

Albumin (g/L), as an important predictor of surgical risk, correlates with the degree of malnutrition and can be an indicator of poor outcomes in hospitalized patients.[7] Postoperative hyperlactatemia is associated with microcirculation damage and decreased tissue ability to extract oxygen. Elevated lactate (mol/L) levels arise due to increased aerobic glycolysis. Lactic acidosis occurs in two clinical scenarios: hypoxic and metabolic.[8] IgA antibodies provide the first line of defense in the mucosa of the respiratory, gastrointestinal, and genitourinary tracts against bacteria and viruses. Increased concentrations of immunoglobulins in the blood have been observed in various conditions, including infections and chronic diseases.[9] Lactate dehydrogenase (LDH) is an important enzyme in anaerobic metabolism and is used as an indicator of acute and chronic diseases. Elevated LDH levels are associated with various conditions, including muscular dystrophy and pathological fluid accumulation.[9]

Anastomotic dehiscence is a serious complication in surgery that can lead to systemic and local complications. The International Study Group of Rectal Cancer recommends grading the severity of this complication. The frequency of dehiscence varies depending on different factors, including the type of surgery and timing of resection.[10-14] Complications of surgical treatment for colorectal cancer contribute to increased morbidity and mortality among patients. Comparison of different anastomosis methods has not revealed significant differences in complication rates.[15] Anastomotic dehiscence is potentially the most dangerous complication of surgical treatment in patients. Risk factors for anastomotic dehiscence are classified into three categories: patient-related factors, operative factors, and disease-related factors.[16]

Nutritional support plays a crucial role in wound healing and postoperative recovery, and poor nutritional status is strongly associated with delayed wound healing and longer hospital stays after surgery.[17] Patients who experience anastomotic dehiscence undergo a "double blow," first as a result of the surgery and second due to sepsis.[18]

The main objective of this study was to examine the role of biochemical markers in serum and peritoneal fluid in the development of postoperative complications in patients with enteral anastomosis.

## PATIENTS AND METHODS

Our study was planned as a prospective research study, using a panel study design that combines a cohort study with a cross-sectional study. The research was conducted at the Clinical-Hospital Center Kosovska Mitrovica and the University Clinical Center Kragujevac. Approval for the implementation of this research was obtained from the Ethics Committee of the institution

where the research was conducted, as well as from the Ethics Committee of the Faculty of Medicine, University of Pristina, with temporary headquarters in Kosovska Mitrovica. The research was carried out from April 2022 to April 2024.

In this study, the first cross-sectional study included biochemical analyses on patients hospitalized with a diagnosis of colorectal cancer, intestinal diverticulum, or GIST. Patients were monitored during the entire treatment process, and in the postoperative period, a second cross-sectional study was conducted, during which certain parameters were monitored on the third, fifth, and seventh postoperative days. The study included patients with a diagnosis of colorectal cancer, intestinal diverticulum, or GIST, confirmed by radiological methods and histopathology, of both sexes, aged 18 years and older. After being informed about the purpose and conduct of the study, written consent was obtained from all participants for their participation in the study and the use of their blood samples and peritoneal drainage fluid for research purposes.

As part of the study, sociodemographic data and blood samples were collected in the preoperative period, as well as blood samples and peritoneal drainage fluid in the postoperative period on the third, fifth, and seventh days after surgery. Sociodemographic data analyzed include gender and age. Clinical parameters collected included: serum albumin, lactate, IgA antibody, and LDH levels before surgery, as well as on the third, fifth, and seventh postoperative days in serum and peritoneal drainage fluid; specific postoperative complications (anastomotic dehiscence); general postoperative complications (urinary tract infection, renal failure, deep vein thrombosis, pneumonia); day of postoperative occurrence of anastomotic dehiscence; localization of the pathological process; type of surgical intervention; type of operation (anastomosis/Hartmann resection); and mortality outcome. Processing and analysis of samples were performed in the laboratory of CHC Kosovska Mitrovica, according to standard protocols and using commercial reagents. The primary outcome variable was the presence of specific complications in patients undergoing surgical treatment.

For the analysis of these parameters, commercial reagents were used on the Beckman Coulter AU 480 automatic analyzer platform (Beckman Coulter, Brea, CA, USA). Commercial calibrators traceable to NIST (National Institute of Standards and Technology) standard reference material and Beckman Coulter Master calibrators were used for method calibration. Test tubes without additives (BD Vacutainer) were used to obtain serum. After free blood coagulation, the sera were separated by centrifugation (15 minutes, 3000 revolutions/minute), aliquoted, and stored at a temperature of -80 degrees Celsius until analysis. Peritoneal drainage fluid was taken in one tube of 1 mL from the abdominal drain. Plastic sterile test tubes without additives, centrifuged, aliquoted, and properly labeled were used. Samples of peritoneal drainage fluid were taken with a sterile syringe on the valve of the abdominal drainage bag, using sterile gloves.

The sample size was determined based on average albumin values as predictors of complication occurrence. According to the literature, 52 patients are required to achieve a study power of 80% and a significance level of .05.[16,19] Statistical analyses were carried out with IBM SPSS for Windows version 21 statistical software. Continuous variables are presented as median (range) and categorical variables as frequencies and percentages. Distributions of the numeric variables were examined by Kolmogorov-Smirnov test for normality. For numeric variables, differences between the groups were determined by the Mann-Whitney test. The chi-square test and the Fisher's exact test were used to determine associations between categorical variables. Spearman's correlation coefficient (rs) was used to test the association. A *P* value of less than .05 was considered statistically significant.

## RESULTS

In this study, data collected from 52 patients with a diagnosis of colorectal cancer, diverticulum of the bowel, or GIST were analyzed. Sociodemographic data revealed that the median (IQR) age of the participants was 65.5 [50.5-73.25]) years, with a gender distribution of 59.6% males and 40.4% females (**Tables 1 and 2**).

There was no statistically significant difference in the age of participants concerning the occurrence of dehiscence (*P*=.655). The average age of patients without dehiscence was (SD 65.9 [9.4]) years, while the median (IQR) age of patients with dehiscence was 65.5 (14) years. There was no statistically significant difference by gender regarding the occurrence of dehiscence (*P*=.870) (**Table 3**). There was no statistically significant difference in the incidence of dehiscence based on the type of surgical procedure (*P*=.196) (**Tables 4 and 5**). The highest number of dehiscences was recorded in patients with the pathological process localized to the sigmoid colon, while the fewest cases were noted in other parts of the colon (**Table 6**).

There was a statistically significant difference in lactate concentration in the drain on the third postoperative day concerning the occurrence of dehiscence (**Table 7**). Lactate concentration was significantly higher

**Table 3.** Frequency of dehiscence by sex.

| | Dehiscence | | *P* |
|---|---|---|---|
| | Yes | No | |
| Gender | | | |
| Male | 8 (61.5) | 23 (59) | .870 |
| Female | 5 (38.5) | 16 (41) | |

Data are number (percentage).

**Table 4.** Frequency of anastomotic dehiscence by surgical procedure.

| | Dehiscence | | *P* |
|---|---|---|---|
| | Yes | No | |
| Surgical procedure | | | |
| Entero-entero | 4 (30.8) | 0 (0) | |
| Ileo-colon | 3 (23.1) | 21 (53.8) | |
| Colon-colon | 0 (0) | 7 (17.9) | .0017 |
| Colon-recto | 1 (7.7) | 3 (7.7) | |
| Hartmann | 5 (38.5) | 8 (20.5) | |

Data are number (percentage).

**Table 5.** Frequency of anastomotic dehiscence by surgical technique.

| | Dehiscence | | *P* |
|---|---|---|---|
| | Yes | No | |
| Surgical technique | | | |
| Anastomosis | 8 (61.5) | 31 (79.5) | .196 |
| Hartmann | 5 (38.5) | 8 (20.5) | |

Data are number (percentage).

in the drainage fluid on the third day after surgery in patients who did not develop dehiscence. Additionally, the concentration of albumin in the drainage fluid on the third postoperative day was statistically significantly higher in patients who developed dehiscence, whereas albumin concentration on the seventh postoperative day was statistically significantly higher in patients who did not develop dehiscence. LDH concentration was statistically significantly higher on the third postoperative day in patients who did not develop dehiscence.

There was a statistically significant strong positive correlation between lactate concentration and IgA antibodies in serum on the fifth postoperative day (rs=.748, *P*=.003). An increase in lactate values was accompanied by an increase in IgA antibody levels (**Table 8**).

**Table 6.** Frequency of anastomotic dehiscence by localization of pathological process.

| | Dehiscence | | *P* |
|---|---|---|---|
| | Yes | No | |
| Localization of pathological process | | | |
| Ascendant colon | 3 (23.1) | 21 (53.8) | |
| Transvese colon | | | |
| Descendant colon | 0 (0) | 2 (5.1) | |
| Sigmoid colon | 5 (38.5) | 16 (41) | .003 |
| Rectal diverticulum | 1 (7.7) | 0 (0) | |
| Small intestine | 3 (23.1) | 0 (0) | |
| GIST | 1 (7.7) | 0 (0) | |

Data are number (percentage).

There was a statistically significant moderate negative correlation between lactate concentration and IgA antibodies in the drain on the third postoperative day (rs=-.670, *P*=.012). An increase in IgA antibody levels was accompanied by a decrease in lactate values. There was also a statistically significant moderate negative correlation between lactate concentration and IgA antibodies in the drain on the fifth postoperative day (rs=-.577, *P*=.039). An increase in IgA antibody levels was associated with a decrease in lactate values (**Table 9**).

There was a statistically significant strong positive correlation between serum IgA antibody concentration and IgA antibodies in the drain on the third, fifth, and seventh days after surgery in patients who developed dehiscence. Elevated serum IgA antibody levels were associated with elevated IgA antibody levels in the drain (**Table 10**).

There was a statistically significant difference in the age of patients concerning the occurrence of general postoperative complications (*P*=.006). The median (minimun-maximum) age of patients who did not develop complications was 65.5 (40.0-88.0) years. General postoperative complications were more common in older patients.

There was also a statistically significant difference in the frequency of general postoperative complications

**Table 7.** Biochemical parameters in serum and peritoneal drainage fluid in relation to the development of dehiscence.

| Biochemical parameters | Dehiscence | | *P* |
|---|---|---|---|
| | Yes | No | |
| Serum lactates | | | |
| Preoperatively | 1.9 (0.8-4.5) | 1.8 (0.8-11.8) | .479 |
| Third day after surgery | 2.1 (0.6-3.1) | 2.24 (0.7-11.4) | .281 |
| Fifth day after surgery | 1.7 (0.5-4.8) | 2.5 (0.7-19.2) | .171 |
| Seventh day after surgery | 1.5 (0.6-23.7) | 3.2 (0.7-20.8) | .145 |
| Lactates in drainage fluid | | | |
| Third day after surgery | 2.2 (0.01-15.1) | 9.8 (0.5-37.5) | **.006** |
| Fifth day after surgery | 6.2 (1.8-24.1) | 8.9 (0.5-35.6) | .353 |
| Seventh day after surgery | 5.4 (1.8-28.3) | 6.0 (0.9-20.2) | .545 |
| Serum albumin | | | |
| Preoperatively | 33 (20-49) | 37 (25-59) | .340 |
| Third day after surgery | 33 (27-49) | 33 (24-55) | .849 |
| Fifth day after surgery | 31 (28-44) | 32.5 (25-56) | .406 |
| Seventh day after surgery | 34 (20-55) | 34 (28-45) | .800 |
| Albumin in drainage fluid | | | |
| Third day after surgery | 30 (21-45) | 25 (15-54) | **.040** |
| Fifth day after surgery | 31 (16-39) | 25 (12-55) | .370 |
| Seventh day after surgery | 29 (21-39) | 38 (32-44) | **.001** |
| IgA in serum | | | |
| Preoperatively | 1.4 (0.4-3.1) | 1.5 (0.5-5.8) | .342 |
| Third day after surgery | 1.0 (0.4-3.9) | 1.4 (0.5-5.0) | .212 |
| Fifth day after surgery | 1.3 (0.3-4.7) | 1.3 (0.4-4.8) | .883 |
| Seventh day after surgery | 1.3 (0.3-4.5) | 1.7 (0.4-4.3) | .866 |
| IgA in drainage fluid | | | |
| Third day after surgery | 0.6 (0.2-4.2) | 0.8 (0.2-3.2) | .672 |
| Fifth day after surgery | 0.6 (0.4-5.8) | 0.9 (0.2-2.9) | .100 |
| Seventh day after surgery | 0.9 (0.2-4.1) | 1.0 (0.5-2.6) | .657 |
| LDH in serum | | | |
| Preoperatively | 251 (17-628) | 253 (86-938) | .866 |
| Third day after surgery | 301 (57-753) | 291 (135-789) | .642 |
| Fifth day after surgery | 276 (16-954) | 248 (94-1152) | .901 |
| Seventh day after surgery | 383 (51-1219) | 293 (45-1263) | .287 |
| LDH in drainage fluid | | | |
| Third day after surgery | 623 (251-1210) | 1622 (95-10770) | **.020** |
| Fifth day after surgery | 738 (86-19500) | 1186.5 (137-14980) | .821 |
| Seventh day after surgery | 1063 (128-30400) | 783 (93-3350) | .600 |

Data are median (minimum-maximum).

**Table 8.** Correlation of IgA antibody and lactate concentrations in preoperative and postoperative serum (Spearman's correlation coeffcient).

| | Lactate preoperatively | Lactate on the third day after surgery | Lactate on the fifth day after surgery | Lactate on the seventh day after surgery |
|---|---|---|---|---|
| IgA antibodies | | | | |

**Table 12.** Biochemical parameters in serum and peritoneal drainage fluid relative to the development of general complications.

| Biochemical parameters | General complications | | *P* |
|---|---|---|---|
| | Yes | No | |
| Serum lactate | | | |
| Preoperatively | 1.5 (0.8-2.9) | 1.9 (0.8-11.8) | .078 |
| Third day after surgery | 1.9 (0.6-3.7) | 2.4 (0.8-11.4) | .326 |
| Fifth day after surgery | 1.9 (0.5-5.1) | 2.3 (0.7-19.2) | .418 |
| Seventh day after surgery | 1.9 (0.6-23.7) | 2.8 (0.7-20.8) | .439 |
| Lactates in drainage fluid | | | |
| Third day after surgery | 2.2 (0.01-15.1) | 10.5 (0.5-37.5) | **.001** |
| Fifth day after surgery | 4.8(0.6-24.1) | 10.0 (0.5-35.6) | **.041** |
| Seventh day after surgery | 4.2 (0.9-28.3) | 8.2 (1.4-23.7) | .683 |
| Serum albumin | | | |
| Preoperatively | 38 (20-50) | 36 (24-59) | .662 |
| Third day after surgery | 34 (26-52) | 32 (24-55) | .138 |
| Fifth day after surgery | 31 (28-48) | 32 (25-56) | .460 |
| Seventh day after surgery | 33.5 (28-51) | 33.5 (20-55) | .755 |
| Albumin in drainage fluid | | | |
| Third day after surgery | 30 (16-48) | 25 (15-54) | **.024** |
| Fifth day after surgery | 34 (16-49) | 25 (12-55) | .065 |
| Seventh day after surgery | 30 (21-44) | 34 (28-43) | .151 |
| IgA in serum | | | |
| Preoperatively | 1.4 (0.4-3.2) | 1.5 (0.6-5.8) | .234 |
| Third day after surgery | 1.4 (0.4-3.9) | 1.4 (0.5-5.0) | .138 |
| Fifth day after surgery | 1.3 (0.4-4.8) | 1.2 (0.3-4.7) | .924 |
| Seventh day after surgery | 1.5 (0.3-4.5) | 1.7 (0.4-4.3) | .866 |
| IgA in drainage fluid | | | |
| Third day after surgery | 0.8 (0.2-4.2) | 0.7 (0.2-2.9) | .812 |
| Fifth day after surgery | 0.9 (0.3-5.8) | 0.9 (0.2-2.9) | .430 |
| Seventh day after surgery | 1.0 (0.2-4.1) | 0.9 (0.5-2.6) | .870 |
| LDH in serum | | | |
| Preoperatively | 282.5 (17-938) | 245 (86-724) | .378 |
| Third day after surgery | 355 (57-789) | 261.5 (135-730) | .194 |
| Fifth day after surgery | 291.5 (16-1152) | 246 (94-974) | .306 |
| Seventh day after surgery | 296 (51-1263) | 313.5 (45-1178) | .899 |
| LDH in drainage fluid | | | |
| Third day after surgery | 603 (155-1308) | 1673 (95-10770) | **.002** |
| Fifth day after surgery | 583.5 (86-3060) | 1224 (137-19500) | .193 |
| Seventh day after surgery | 1008 (128-2814) | 1858 (93-30400) | .369 |

Data are median (minimum-maximum).

based on the type of surgical procedure (*P*=.037). General complications were more frequent with the anastomosis technique compared to the Hartmann procedure (**Table 11**).

There was a statistically significant difference in lactate concentration in the drainage fluid on the third and fifth days after surgery with respect to the development of general postoperative complications. The average lactate value in the drainage fluid on the third and fifth days after surgery is significantly higher in patients who did not develop general postoperative complications. There is also a statistically significant difference in albumin concentration in the drainage fluid on the third day after surgery concerning the development of general postoperative complications. The average albumin value in the drainage fluid on the third day after surgery is significantly higher in patients who developed general postoperative complications.

A A A Y A The

tissue regeneration and the stabilization of postoperative metabolism, which reduce the risk of dehiscence. In other words, patients who show a rapid recovery of albumin after a temporary drop may have a better potential for wound healing.[23]

Lactate values were determined in serum on the third, fifth, and seventh postoperative days, but no association with anastomotic dehiscence was found. Also, lactate values in abdominal drainage fluid on the same days were measured, where only on the third postoperative day was a significantly higher concentration of lactate identified in patients with dehiscence. Overall, only 5 (9.6%) patients had anastomotic dehiscence on the third postoperative day. In contrast to these findings, Sparreboom et al[24] also measured lactate values in serum but did not find a significant difference on the third postoperative day. A high lactate level usually signals anaerobic glycolysis, indicating tissue hypoxia or ischemia. However, elevated lactate in patients who have not developed dehiscence may be indicative of a successful adaptive response to temporary ischemia. Short-term ischemia can trigger regenerative mechanisms in tissues, such as angiogenesis (formation of new blood vessels), which improves perfusion of the anastomosis and reduces the risk of dehiscence.[25]

The concentration of LDH in abdominal drainage fluid on the third postoperative day was statistically significantly higher in patients who did not develop dehiscence. Elevated LDH in drainage fluid may indicate adequate removal of damaged cells and tissues, contributing to better and faster wound healing. Conversely, in patients who developed dehiscence, lower LDH concentration may indicate inadequate metabolic response and the inability of tissue to recover properly after surgical intervention. These findings suggest that LDH may play an important role in the healing process and that its concentration in drainage fluid may be a useful indicator of recovery success. It should be noted that we did not find studies in the available literature directly investigating the concentration of LDH in drainage fluid in the context of anastomotic dehiscence. Therefore, our results may represent a new contribution to understanding this area and indicate the need for further research to better understand the role of LDH and its potential application as a biomarker in assessing the risk of postoperative complications.

This study demonstrates a statistically significant and strong positive correlation between lactate concentration and IgA antibodies in the serum on the fifth postoperative day ($r_s$=.748, *P*=.003). This result suggests that an increase in lactate levels, which is an indicator of metabolic stress or hypoxia, correlates with an increase in IgA antibody levels, which are part of the immune response to inflammatory and infectious agents. This correlation may indicate that both parameters are responses to postoperative inflammatory or stress conditions. The increase in lactate might stimulate the production of IgA antibodies as part of the immune response to inflammatory stimuli. As we did not find similar studies in the available literature. Our result represents a new insight into the relationship between these parameters in the postoperative period.

In this study, a statistically significant moderate negative correlation was identified between lactate concentration and IgA antibodies in the drain on the third and fifth postoperative days ($r_s$=-.670, *P*=.012 and $r_s$=-.577, *P*=.039). These results indicate that an increase in IgA antibody levels correlates with a decrease in lactate levels in the drain fluid. This negative correlation may suggest that elevated levels of IgA antibodies, which play a role in the immune response to inflammatory stimuli, could be associated with improved patient conditions, resulting in reduced lactate levels. Lactate is often an indicator of metabolic stress and hypoxia, so an increase in IgA antibodies might be part of a positive immune response leading to decreased metabolic burden. Since similar studies were not found in the available literature, our result provides a new insight into the dynamics between these parameters in the postoperative period.

Additionally, the study found a statistically significant strong positive correlation between serum IgA antibody levels and IgA antibody levels in the drain on the third, fifth, and seventh postoperative days in patients who developed anastomotic dehiscence. This positive correlation indicates that an increase in IgA antibodies in the serum is accompanied by an increase in the same antibodies in the drain fluid. Given that IgA antibodies are crucial in the local immune response to inflammation and infections, elevated levels of these antibodies may indicate an active immune response in the anastomosis area. This suggests that the immune response, represented by increased IgA levels, could be an important factor in the development of dehiscence, potentially as a reaction to inflammatory processes or infections affecting wound healing. Since we did not find similar studies in the literature exploring this aspect, our results may offer new and significant insights into the relationship between serum and drain IgA antibodies in the context of postoperative complications such as anastomotic dehiscence. A low level of IgA in this context may mean less inflammatory reaction in the abdominal cavity. Excessive inflammation can lead to tissue damage and impaired healing, which

increases the risk of dehiscence. Low IgA, in this case, could reflect a reduction in the inflammatory response, which may help preserve the integrity of the anastomosis, especially if the inflammation is controlled by other mechanisms of the immune system.[26]

The study showed statistically significant differences in the concentrations of certain biochemical parameters in drain fluid with respect to the development of general postoperative complications. The concentration of lactate in the drain fluid on the third and fifth postoperative days was significantly higher in patients who did not develop general postoperative complications. This finding suggests that elevated lactate levels might be associated with better stability and a lower inflammatory response in the immediate postoperative period in patients without complications.

Additionally, the concentration of LDH in the drain fluid on the third postoperative day was significantly higher in patients who did not develop general postoperative complications. This finding may indicate more active metabolism and recovery in the early phase of recovery among patients without complications, potentially reflecting better overall health or a more successful healing process. These results are intriguing and may highlight the complex role of these parameters in monitoring and managing postoperative conditions. Since similar studies were not found in the available literature, our results may offer new insights and encourage future research to explore how these biochemical markers could influence the prognosis and management of postoperative complications.

In this study, 38.5% of patients who underwent Hartmann's procedure experienced specific postoperative complications, including dehiscence of the rectal stump that was closed blindly. Similarly, but with a slightly higher complication rate, the study by Hut et al[27] found that 44% of patients who underwent the same procedure developed postoperative complications. These data highlight a significant risk of specific postoperative complications in patients undergoing Hartmann's procedure, and the differences in complication rates may be attributed to varying treatment protocols or patient sample variations.

Rectal diverticulum is a rare condition with an etiology that remains poorly understood.[28] An interesting angiomorphological study from Portugal, involving 80 human cadavers, proposed the hypothesis that the low prevalence of rectal diverticula may be related to the course of the vasa recta.[28] In cases of acute presentation, when conservative treatment fails and there is significant clinical urgency, emergency intervention in the form of rectal resection or Hartmann's procedure is considered.[28] During the observation period of our study, we identified only one case of rectal diverticulum perforation that required emergency surgical intervention via Hartmann's procedure. This indicates the rarity of this complication but also highlights the need for prompt and decisive surgical action in such cases. The patient who required emergency surgery due to perforation of a rectal diverticulum was a 66-year-old woman. Diverticular disease of the colon is a common condition affecting about 50% of individuals over the age of 80.[29] This high prevalence underscores the necessity for careful monitoring and timely response in older patients exhibiting symptoms of this disease, as complications, though rare, may necessitate urgent and serious surgical interventions.

GISTs account for approximately 1% of primary gastrointestinal tract cancers.[30] A study by Kassimi et al28 presented a case of a 79-year-old woman diagnosed with GIST. In our study, we included a 40-year-old woman with the same diagnosis during the observation period. This age difference highlights that GISTs can affect various age groups, which is significant for clinical practice in terms of early recognition and diagnosis of this rare but serious disease.

The strengths of this study include a comprehensive analysis of concentrations of various biochemical parameters in serum and drain fluid, which allows for a deeper understanding of their role in predicting and managing postoperative complications such as anastomotic dehiscence and general postoperative complications. This study is also one of the first to investigate the relationship between lactate and IgA antibody concentrations in both drain fluid and serum, providing new insights into their potential significance as indicators for postoperative monitoring. Additionally, the limited availability of literature specifically addressing certain aspects of this research highlights the need for further investigation to confirm and clarify the obtained results.

The lack of a control group in our study represents a limitation we consciously accepted, taking into account the specific characteristics of the patient population and the nature of postoperative complications associated with enteric anastomoses. The decision not to include a control group was based on several key factors:

First, ethical considerations played a significant role. A control group would have involved patients who were not subjected to surgical intervention, from whom peritoneal fluid would have been collected for comparison with operated patients. Collecting peritoneal fluid from patients who were not undergoing an anastomosis would have constituted an unnecessary invasive procedure without clear clinical benefit for those patients.

In accordance with the ethical principles of medical research, our priority was to minimize risks and discomfort for the patients.

Second, the nature of complications following enteric anastomoses is such that they occur exclusively in operated patients. A control group of non-operated patients would not have provided relevant data for analysis because these patients have no potential to develop the postoperative complications that were the focus of our research, such as anastomotic leakage, peritoneal infection, and other similar complications. Therefore, comparison with a non-operated control group would not have contributed to a deeper understanding of the markers we were investigating, as postoperative changes in serum and peritoneal fluid are directly related to the surgical procedure itself.

Our approach was to closely monitor serial samples of peritoneal fluid and serum in operated patients and thoroughly analyse changes in biological markers before and after the onset of complications. This strategy allowed for longitudinal tracking of each patient as a kind of internal control, which increased the strength of our design and enabled more precise identification of markers associated with complications.

In order to further investigate the relationships between biochemical markers and postoperative complications, future studies should include larger patient samples. This would allow a more detailed analysis and improve the understanding of these relationships.

The concentration of biochemical markers in drain fluid can serve as a valuable indicator for assessing postoperative complications, including anastomotic dehiscence. This research demonstrated significant differences in the concentrations of lactate, albumin, and LDH in drain fluid on the third and fifth days after surgery, in relation to the development of complications. These results suggest that monitoring these markers may aid in the early identification of patients at risk for complications, such as dehiscence. Specifically, significant variations in lactate and LDH concentrations point to potential mechanisms associated with postoperative outcomes.

## REFERENCES


**1.** Kostić Z. C-reaktivni protein i matriksna metaloproteinaza-9 u peritoneumskoj drenažnoj tečnosti kao rani pokazatelji postoperativnih komplikacija kod bolesnika operisanih zbog kolorektalnog karcinoma. Doktorska disertacija VMA, Beograd 2014

**2.** Goulder F. Bowel anastomoses: The theory, the practice and the evidence base. World J Gastrointest Surg. 2012;27;4(9):208-13

**3.** Nguyen DA, Phan TAM, Do PTT, Thai TT. Emergency surgery for obstructed colorectal cancer in Vietman. Asian J Surg. 2020;43(6):683-689

**4.** Dsouza R, Varghese G, Korula DR, Dutta AK. Crohn's disease associated adenocarcinoma of oleocaecal region: a miscalculated approach. BMJ Case Rep. 2020;16;13(4):e234512

**5.** Tan L, Liu ZL, Ran MN, Tang LH, Pu YJ, Liu YL, et al. Comparison of the prognosis of four different treatment strategies for acute left malignant colonic obstruction: a systematic review and network meta-analysis. World J Emerg Surg. 202;18;16(1):11

**6.** Mege D, Manceau G, Berjot LB, Bridoux V, Lakkis Z, Venara A, et al. Surgical management of obstructive right-sided colon cancer at a national level results of a multicenter study of the French Surgical Association in 776 patients. Eur J Surg Oncol. 2018;44(10):1522-1531

**7.** Kiziltan R, Yilmaz Ö, Aras A, Çelik S, Kotan Ç. Factors affecting mortality in emergency surgery in cases of complicated colorectal cancer. Med Glas (Zenica). 2016;13(1):62-7

**8.** Veličković J, Palibrk I, Miličić B, Veličković D, Jovanović B, Rakić G, et al. The assotiation of early postoperative lactate levels with morbidity after elective major abdominal surgery. Bosn J Basic Med Sci. 2019;19(1):72-80

**9.** Farhana A, Lappin SL. Biochemistry, lactate dehydrogenase. StatPearls Publishing; 2022. Bookshelf ID: NBK557536

**10.** Sciuto A, Merola G, Palma GDD, Sodo M, Pirozzi F, Bracale UM, et al. Predictive factors for anastomotic leakage after laparoscopic colorectal surgery. World J Gastroenterol. 2018;24(21):2247-2260

**11.** Deng S, Cao Y, Gu J, Wu K, Li J, Tao K, et al. Endoscopic diagnosis and treatment of complete anastomosis stenosis after colorectal resection without protective ileostomy: report of two cases and literature review. J Int Med Res. 2020;48(4):300060520914833

**12.** Arron MNN, Greijdanus NG, Breok RPGT, Dekker JWT, Workum F, Goor H, et al. Trends in risk factors of anastomotic leakage after colorectal cancer surgery (2011-2019): a dutch population-based study. Colorectal Dis. 202;23(12):3251-3261

**13.** Tinca AC, Ion D, Bolocan A, Muşat F, Andronic O, Păduraru DN. Low anterior resection – emergency versus elective surgery in rectal cancer treatment – comparative analysis. Chirurgia (Bucur). 2019;114(2):216-221

**14.** Walker KG, Bell SW, Rickard MJFX, Mehanna D, Dent OF, Chapuis PH, et al. Anastomotic leakage is predictive of diminished survival after potentially curative resection for colorectal cancer. Ann Surg 2004;240(2):255-259

**15.** Chassin JL, Rifkind KM, Sussman B, Kassel B, Fingaret A, Drager S, et al. The stapled gastrointestinal tract anastomosis: incidence of postoperative complications compared with the sutured anastomosis. Ann Surg. 1978;188(5):689-96

**16.** Telem DA, Chin EH, Nguyen SQ, Divino CM. Risk factors for anastomotic leak following colorectal surgery: a case-control study. Arch Surg. 2010;145(4):371-6

**17.** Lee HS, Shim H, Jang JY, Lee H, Lee JG. Early feeding is feasible after emergency gastrointestinal surgery. Yonsei Med J. 2014;55(2):395-400

**18.** McArdle CS, McMillan DC, Hole DJ. Impact of anastomotic leakage on long-term survival of patients undergoing curative resection for colorectal cancer. Br J Surg. 2005;92(9):1150-4

**19.** Faul F, Erdfelder E, Lang AG, Bucher A. G*Power 3: A flexibe statistical power analysis program for the social, behavioral, and biomedical sciences. Behavior Research Methods 2007;39:175-191

**20.** Su'a B, Milne T, Jaung R, Jin JZ, Svirskis D, Bissett IP, et al. Detection of Anastomotic Leakage Following Elective Colonic Surgery: Results of the Prospective Biomarkers and Anastomotic Leakage (BALL) Study. J Surg Res. 2022;273:85-92

**21.** Paliogiannis P, Deidda S, Maslyankov S, Paycheva T, Farag A, Mashhour A, et al. C reactive protein to albumin ratio (CAR) as predictor of anastomotic leakage in colorectal surgery. Surg Oncol. 2021;38:101621

**22.** Shimura T, Toiyama Y, Hiro J, Imaoka H, Fujikawa H, Kobayashi M, et al. Monitoring perioperative serum albumin can identify anastomotic leakage in colorectal cancer patients with curative intent. Asian J Surg. 2018;41(1):30-38

**23.** Vincent JL, Dubois MJ, Navickis RJ, Wilkes MM. Hypoalbuminemia in acute illness: is there a rationale for intervention? A meta-analysis of cohort studies and controlled trials. Ann Surg. 2003 Mar;237(3):319-34

**24.** Sparreboom CL, Komen N, Rizopoulos D, Verhaar AP, Dik WA, Wu Z, et al. A multicentre cohort study of serum and peritoneal biomarkers to predict anastomotic leakage after rectal cancer resection. Colorectal Dis. 2020;22(1):36-45

**25.** Brooks GA. Lactate as a fulcrum of metabolism. Redox Biol. 2020 Aug;35:101454

**26.** Cerutti A, Rescigno M. The biology of intestinal immunoglobulin A responses. Immunity. 2008 Jun;28(6):740-50

**27.** Hut A, Akturk OM, Cakir M. Classification of Hartmann's procedure reversal complications according to the Clavien-Dindo System. Ann Ital Chir. 2022;92:130-134

**28.** Gumber A, O'Kelly J, Shalli K. Rectal diverticulum: a rare surgical dilemma that manifest in different ways and manageable with new treatment options. Transl Cancer Res. 2023;12(7):1660-1664

**29.** Gorgoraptis S, Xenaki S, Athanasakis E, Daskalaki A, Lasithiotakis K, Chrysou E, et al. A case of solitary rectal diverticulum presenting with a large retrorectal abscess. Ann Med Surg (Lond). 2019;49:57-60

**30.** Kassimi M, Berrada S, Habi J, Guerroum H, Chihabeddine M, Mahi M. Gastrointestinal stromal tumor of the small intestine with lung metastasis. Radiol Case Rep. 2021;17(1):32-34