Letters to the Editor

COMMENTS ON "THE SCM TEST FOR CANCER. AN EVALUATION IN TERMS

OF LYMPHOCYTES FROM HEALTHY DONORS AND CANCER PATIENTS"

by MITCHELL. WOOD. PENTYCROSS AND BAGSHAWE

· · ·	(<u>Br. J. Cancer (1980)</u> 41.	772)
	SIR —In their paper Mitchell <i>et al.</i> give the ocytes	incubated with PHA or some
» <u></u>	one in the baber interior of the zire the outers	
i	impression that having established the SCM antigen	preparations, the rate of FDA
<u></u>		
6		
¥		
ja-		
-		

948 LETTERS TO THE EDITOR of SCM-responding lymphocytes are normally <u>contact with us and did not find it difficult to</u> higher than 0.185 (Cercek & Cercek, 1978b). Hence, their low control P values again indiassimilate the new findings. On the contrary, after establishing the SCM measurements. modifications of the test (Pritchard et al., It is not, therefore, surprising that they were

T

1

unable to obtain as good a differentiation 1978: Øriasaeter et al., 1979). It follows that

949

LETTERS TO THE EDITOR

I Deserve TI Deserve		<u>, ,, 1, ,1, 1.</u>	<u> </u>	
a. `				
	ĩ			
a a a a a a a a a a a a a a a a a a a				
۹ <u>ــــــــــــــــــــــــــــــــــــ</u>				
j				
				,
<u>-</u>				
				4
	2			
	и то станование и на			