

ORIGINAL RESEARCH

Assessing Retail Fruit and Vegetable Availability in Urban and Rural Underserved Communities

Akiko S. Hosler, PhD, Deepa T. Rajulu, MS, Bonnie L. Fredrick, MS, Adrienne E. Ronsani, MS

Suggested citation for this article: Hosler AS, Rajulu DT, Fredrick BL, Ronsani AE. Assessing retail fruit and vegetable availability in urban and rural underserved communities. *Prev Chronic Dis* 2008;5(4). http://www.cdc.gov/pcd/issues/2008/oct/07_0169.htm. Accessed [date].

PEER REVIEWED

Abstract

Introduction

Fruits and vegetables (F&Vs) are important parts of a healthy, balanced diet. Consumption of F&Vs is low among residents of socioeconomically disadvantaged communities. We investigated and compared retail F&V availability in urban and rural underserved communities in New York State.

Methods

All food retail stores and farmers' markets (N = 263) in downtown Albany and in Columbia and Greene counties in New York State were visited and surveyed. Food stores were classified as *F&V stores* if they stocked more than the minimum varieties of fresh F&Vs defined by this study and as *fruit-for-snack stores* if they had ready-to-eat fruits only. Store density per 10,000 residents was calculated as a standardized measure of F&V availability. Adjustment weights were created to incorporate store size and business hours into the analysis.

Results

The weight-adjusted density (per 10,000 residents) of F&V stores was 4.6 in Albany's minority neighborhood (reference category), 11.4 in Albany's racially mixed neigh-

borhood ($P = .01$), 7.8 in Columbia and Greene counties' rural community ($P = .10$), and 9.8 in Columbia and Greene counties' small-town community ($P = .02$). Significant differences were not found in fruit-for-snack stores, which ranged from 2.0 per 10,000 in the mixed neighborhood to 3.4 per 10,000 in the rural community.

Conclusion

The urban minority neighborhood had the most barriers to fresh F&Vs in retail outlets, even when compared with the rural community. The low availability of retail F&Vs in the minority neighborhood was attributed to the lack of supermarkets and not the absolute lack of food stores. Public health intervention strategies to increase retail F&V availability in urban minority neighborhoods may aid in mitigating these disparities.

Introduction

Fruits and vegetables (F&Vs) are important parts of a healthy, balanced diet. Many studies have documented the benefits of F&V consumption for weight control (1,2) and prevention of chronic diseases, including cardiovascular disease (3,4), diabetes (5,6), and certain types of cancer (7). On the basis of recommendations by the *Dietary Guidelines for Americans* (8), *Healthy People 2010* (HP 2010) has set national objectives for increasing proportions of Americans who consume at least 2 daily servings of fruit and 3 daily servings of vegetables, with at least 1 serving being a dark-green or orange-colored vegetable (9). Yet most Americans do not consume the daily recommended amount of F&Vs (10,11). Consumption of F&Vs is reportedly low among socioeconomically disadvantaged, rural, and poor residents (12).

daily servings of F&Vs are complex. Environmental barriers, notably the limited availability of fresh produce in local that individuals living in close proximity to supermarkets in neighborhoods without supermarkets or located farther away from supermarkets (15-17). The inverse

One study found that urban African American women who shopped at their neighborhood small-food stores consumed fewer F&Vs than did their higher-income peers who were

Studies suggest that consumption of F&Vs by disadvantaged individuals can be increased if they have access to a supermarket or food store that provides an adequate

to retail fresh F&Vs have increased considerably (21). Supermarkets and grocery stores with produce departments are much less likely to be found in economically

remote, rural communities also face barriers to obtaining retail fresh F&Vs (12).

store classification scheme; and a standardized, weight-adjusted method to calculate store density as a measure

of fresh F&Vs.

Study communities

County, and a downtown portion of the city of Albany, the Albany Prevention Research Center's Diabetes Prevention and Control Project (28). We defined downtown Albany

("Albany" hereafter) as 4 geographically contiguous zip of 2000 US census data showed that these areas formed a region with a population of low socioeconomic status

that residential racial composition was the sole predictor for availability of low-fat milk and high-fiber bread in these same communities (29). We divided Albany into a minority neighborhood, which was defined by census

Hispanic populations, and a mixed neighborhood, which Hispanic populations.

population is predominantly non-Hispanic white, into a community was composed of CBGs with populations that are 100% rural, as defined by the US Census Bureau two counties' remaining CBGs. This division contrasted

communities during July through September 2003. We defined a "food store" as a retail store that was open for

the survey and that stocked at least 1 of the following food

were included regardless of the number of days they were open each week. We used 3 sources to obtain a list of food

and Markets' (NYSDAM's) current inspected food store database (accessed using a Freedom of Information Act request), 2) the NYSDAM's online guide for farm-fresh products and farmers' markets (31), and 3) online local yel-

definition and whether they were located in the study area. After eliminating nonqualified, closed, and duplicate

process of identifying food stores is shown in Figure 1.

The 1-page survey assessed more than 10 types of food and nonfood items. Among other indicators, F&V pres-

PREVENTING CHRONIC DISEASE

PUBLIC HEALTH RESEARCH, PRACTICE, AND P

ence and variety were assessed. A team of 2 or 3 trained surveyors conducted the in-store surveys. If fewer than 10 types of fresh F&Vs were sold, surveyors wrote down each type of F&V; if more than 10 types of F&Vs were sold, surveyors noted that there was a large variety. Surveyors noted whether at least 1 dark-green or orange-colored vegetable was present, and they collected information about store hours and number of cash registers used. We used the number of cash registers as a measure for store size, as suggested by a previous study (25). To ensure interrater reliability, results of the assessment were recorded when all members of the survey team agreed on the assessment decisions. As a standardized protocol, surveyors asked permission to conduct a survey, and all stores granted permission. Written consent was obtained from an employee of each store who was authorized to make customer-related decisions.

The location of each store was measured at the front door by coordinates using a hand-held global positioning system (GPS) device (Thales Navigation, Inc, Alexandria, Virginia). Survey data were analyzed using SPSS version 12.0 (SPSS, Inc, Chicago, Illinois), and GPS information was processed using MapInfo software (Pitney Bowes Software, Inc, Troy, New York).

Measurement

We initially grouped surveyed food stores using a business-type classification system developed by Morland et al (22). However, we found that considerable variations in

fresh F&V presence and variety existed within the same business type. For instance, 46% of gas station stores had some fresh F&Vs, but 54% did not carry F&Vs at all. Among convenience stores in the same chain, some locations had produce sections but others did not. Furthermore, “supermarket” did not automatically mean that the store had a produce department; 1 large national chain’s local stores had no produce departments at all. Therefore, we devised a store classification system to properly categorize the stores by the availability and variety of fresh F&Vs sold (Figure 2).

We categorized the business as an *F&V store* if it stocked at least 2 types of fresh fruit, excluding lemons and limes, and at least 3 types of fresh vegetables, including at least 1 dark-green or orange-colored vegetable. We further grouped the F&V stores into 3 types: *super produce stores*, or supermarkets with a produce department; *year-round produce stores*, which were nonsupermarket F&V stores that operate year-round, including grocery stores, convenience stores, and specialty food stores; and *seasonal produce stores*. The last category included seasonal farm (ie, barn) stores, roadside huts and stands, and farmers’ markets. Stores that did not meet the F&V measure but did carry at least 1 type of ready-to-eat fresh fruit, such as apples, oranges, and bananas, were designated as *fruit-for-snack stores*. We determined that recognition of this type of store would be useful for promoting fruit consumption (ie, choosing a piece of fruit in lieu of candy or a snack food). The remaining stores were grouped into the *non-F&V store* category.

ability, we calculated store density per 10,000 residents and compared the densities of the 4 communities. Data

all stores equally, including large supermarkets, small corner stores, and farmers' markets that open only 4 to 7 hours per week. To adjust for store size and operating

was natural-log-transformed and added the constant of 1 to normalize the right-skewed distribution with the minimum value of 1. Store hours were measured by per-week

with 98 hours as both the mean and the median values. A 98-hours-per-week operation is equivalent to opening from 7:00 A. . to 9:00 P.M. daily, a time period we determined was adequate for most residents to shop for F&Vs. We

that was open for 98 hours per week. The weight increased if the store had more than 1 cash register and decreased if the store operated fewer than 98 hours per week.

from 1 plus the natural-log-transformed number of cash registers multiplied by the store hours per week divided by 98. For stores open 98 hours per week or longer, a constant of 98 was used:

w_i / 98) where w = adjustment weight, ci = number of cash registers, and hi = business hours per week if $hi < 98$, $hi = 98$ for all other stores.

Results

Characteristics of the study communities are presented in Table 1. The minority neighborhood had the highest percentage of racial and ethnic minorities (74.1%) of the 4 communities, and the rural community had the lowest percentage (4.6%). One in 3 residents in the minority neighborhood and 1 in 4 residents in the mixed neighborhood were living below poverty. In both the mixed and minority neighborhoods, three-quarters of housing units were renter-occupied. At least one-third of households in both the minority and mixed neighborhoods did not have a vehicle. The rural community, with a total population of nearly 62,000, had the lowest rates among the 4 study communities of racial/ethnic minorities (4.6%), individuals living in poverty (8.9%), renter-occupied housing (24.6%), and households without a vehicle (5.9%), and it had the highest rate of citizens aged 65 years or older (16.3%).

A total of 263 eligible food stores and farmers' markets were categorized with our classification system. Of stores designated as F&V stores, which made up 36.5% of all food stores, 16 were super produce stores, 54 were year-round produce stores, and 26 were seasonal produce stores. Most F&V stores stocked far more than the minimum fresh F&Vs defined by this study. Additionally, 37 stores (14%) were fruit-for-snack stores, and 130 stores (49%) were non-F&V stores. Seventeen stores classified as non-F&V stores carried some lemons, limes, and/or vegetables, but none were dark green or orange-colored. A summary of statistics for the numbers of cash registers and store hours, as well as adjustment weights, is shown in Table 2.

The availability of fresh F&Vs was measured by store density (Table 3). Unadjusted population density of F&V stores was 4.2 per 10,000 residents for the minority neighborhood, 7.3 per 10,000 residents for the mixed neighborhood, 6.5 per 10,000 residents for the rural community, and 6.8 per 10,000 residents for the small-town community. No statistically significant differences were found among the communities.

Population densities of all stores were weight-adjusted to factor in store size and business hours. For every 10,000 residents the communities had the following number of F&V stores: 4.6 for the minority neighborhood, 11.4 for the mixed neighborhood, 7.8 for the rural community, and 9.8 for the small-town community. The store densities were significantly higher for the mixed neighborhood ($P = .01$) and

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors' affiliated institutions. Use of trade names is for identification only and does not imply endorsement by any of the groups named above.

the small-town community ($P = .02$) than for the minority neighborhood. This was largely because the adjustment weight upwardly corrected the contribution of super produce stores, which were present in the mixed neighborhood (7.3 per 10,000 residents), rural community (2.4 per 10,000 residents), and small-town community (4.0 per 10,000 residents) but not in the minority neighborhood. The contribution of seasonal stores was downwardly corrected by the weight, primarily because of their shorter business hours, but the small-town community still had a significantly higher density of seasonal stores (1.6 per 10,000 residents, $P = .04$) than did the minority neighborhood, where there were no seasonal stores. No significant differences were found among year-round produce stores (ranged from 4.0 per 10,000 residents in the mixed neighborhood to 4.6 in the minority neighborhood), fruit-for-snack stores (ranged from 2.0 per 10,000 residents in the mixed neighborhood to 3.4 in the rural community), and all food stores combined (ranged from 17.7 per 10,000 residents in the rural community to 25.4 per in the mixed community).

Discussion

Our study demonstrated that the urban minority neighborhood was the most disadvantaged in terms of retail F&V availability, as measured by the population density of F&V stores. The low retail availability of fresh F&Vs in this community appears to be largely because of the lack of super produce stores and not the absolute lack of food stores. In fact, the urban minority community had a large number of food stores, and, of the 4 communities, the highest population density of year-round produce stores and non-F&V stores and the second-highest population density of fruit-for-snack stores in the weight-adjusted model. This is a case of relative deficiency, in which high-impact super produce stores were conspicuously missing. In contrast, the urban mixed neighborhood had 3 large super produce stores. The higher density by population of supermarkets and large grocery stores in racially mixed areas compared with predominantly minority areas in the urban setting has been reported by other studies conducted elsewhere in the United States (22,26,27).

Although the results were not statistically significant, the rural community had higher overall fresh F&V availability compared with the urban minority neighborhood. The rural community had the lowest density by population of total food stores but had the highest density of fruit-for-

snack stores and the second highest density of year-round produce stores. The high availability of ready-to-eat fruits can be explained by the existence of many family-owned fruit orchards and berry farms in this community. The small-town community had the second highest overall F&V availability, with a balanced representation of super produce, year-round produce, and seasonal produce stores. In the winter when seasonal produce stores are not in business, both communities in Columbia and Greene counties probably see some decline in F&V availability. However, the decline probably does not reach the level of the urban minority neighborhood because these seasonal stores make a small contribution to the overall F&V availability in the rural and small-town communities.

Our study has several limitations. We used a cross-sectional survey of food stores and did not adjust for the variability caused by harvest, delivery, or reshelving schedule. We did not directly assess quantity or quality of fresh F&Vs because of a lack of standard measurements suited for field studies. The designation of neighborhoods and communities was made on the basis of CBG data and may differ from actual areas designated as communities. The rural community encompasses a large tract of land, and most remote portions of the community may have more environmental barriers to retail F&V availability. The store classification system and the adjustment weight were new tools and have not been tested in other communities. This study, by design, did not assess the availability of non-retail fresh F&Vs such as those grown in private or community gardens. Availability of canned, jarred, and frozen F&Vs was not assessed because of time constraints, the lack of standard measurements, and the presence of salt, sugar, and nutrient loss that can occur in the process of canning, jarring, and freezing F&Vs.

A growing need exists to develop public health policies and innovative intervention strategies to increase retail

PREVENTING CHRONIC DISEAS

PREVENTING CHRONIC DIS

14. Seymore JD, Yaroch AL, Serdula M, Blanck HM, Khan LK. Impact of nutrition environmental interventions on point-of-purchase behavior in adults: a review. *Prev Med* 2004;39(Suppl 2):S108-136.
15. Morland K, Wing S, Diez Roux A. The contextual effect of the local food environment on residents' diets: the atherosclerosis risk in communities study. *Am J Public Health* 2002;92(11):1761-7.
16. Rose D, Richards R. Food store access and household fruit and vegetable use among participants in the US Food Stamp Program. *Public Health Nutr* 2004;7(8):1081-8.
17. Laraia BA, Siega-Riz AM, Kaufman JS, Jones SJ. Proximity of supermarkets is positively associated with diet quality index for pregnancy. *Prev Med* 2004;39(5):869-75.
18. Zenk SN, Schulz AJ, Hollis-Neely T, Campbell RT, Holmes N, Watkins G, et al. Fruit and vegetable intake in African Americans: income and store characteristics. *Am J Prev Med* 2005;29(1):1-9.
19. Kristal AR, Goldenhar L, Muldoon J, Morton RF. Evaluation of a supermarket intervention to increase consumption of fruits and vegetables. *Am J Health Promot* 1997;11(6):422-5.
20. Sloane DC, Diamant AL, Lewis LB, Yancey AK, Flynn G, Nascimento LM, et al. Improving the nutritional resource environment for healthy living through community-based participatory research. *J Gen Intern Med* 2003;18(7):568-75.
21. Curtis KA, McClellan S. Falling through the safety net: poverty, food assistance and shopping constraints in an American City. *Urban Anthropol* 1995;24:93-135.
22. Morland K, Wing S, Diez Roux A, Poole C. Neighborhood characteristics associated with the location of food stores and food service places. *Am J Prev Med* 2002;22(1):23-9.
23. Shaffer A. The persistence of L.A.'s grocery gap: the need for a new food policy and approach to market development. Los Angeles (CA): Center for Food and Justice, Urban and Environmental Policy Institute, Occidental College; 2002.
24. Zenk SN, Schulz AJ, Israel BA, James SA, Bao S, Wilson ML. Neighborhood racial composition, neighborhood poverty, and the spatial accessibility of supermarkets in metropolitan Detroit. *Am J Public Health* 2005;95(4):660-7.
25. Horowitz C, Colson KA, Hebert PL, Lancaster K. Barriers to buying healthy foods for people with diabetes: evidence of environmental disparities. *Am J Public Health* 2004;94(9):1549-54.
26. Baker EA, Schootman M, Barnidge E, Kelly C. The role of race and poverty in access to foods that enable individuals to adhere to dietary guidelines. *Prev Chronic Dis* 2006;3(3). http://www.cdc.gov/pcd/issues/2006/jul/05_0217.htm
27. Powell LM, Slater S, Mirtcheva D, Bao Y, Chaloupka FJ. Food store availability and neighborhood characteristics in the United States. *Prev Med* 2007;44(3):189-95.
28. Shaw BA, Gallant MP, Riley-Jacome M, Spokane LS. Assessing sources of support for diabetes self-care in urban and rural underserved communities. *J Community Health* 2006;31(5):393-412.
29. Hosler AS, Varadarajulu D, Ronsani AE, Fredrick BL, Fisher BD. Low-fat milk and high-fiber bread availability in food stores in urban and rural communities. *J Public Health Manage Pract* 2006;12(6):556-62.
30. Question and answer center. Washington (DC): U.S. Census Bureau. https://ask.census.gov/cgi-bin/askcensus.cfg/php/enduser/std_alp.php. Accessed September 10, 2007.
31. Farm and market search. Albany (NY): New York State Department of Agriculture and Markets. <http://www.agmkt.state.ny.us>. Accessed September 10, 2007.
32. Cummins S, Macintyre S. "Food deserts" — evidence and assumption in health policy making. *BMJ* 2002;325(7361):436-8.

PREVENTING CHRONIC DISEAS

Table 3. Unadjusted and Weight-Adjusted Fruit and Vegetable Availability, by Store Density Per 10,000 Residents, Albany, Columbia County, and Greene County, New York, 2003

Store Type	Minority Neighborhood ^a	Mixed Neighborhood ^b		Rural Community ^c		Small-Town Community ^d	
	Density	Density	P Value	Density	P Value	Density	P Value
Unadjusted							
F&V store ^e , total	4.2	7.3	.19	6.5	.20	6.8	.16
Super produce ^f	0.0	2.0	.02	1.0	.11	1.4	.06
Year-round produce ^g	4.2	3.3	.66	3.7	.73	3.0	.40
Seasonal produce ^h	0.0	2.0	.02	1.8	.03	2.4	.01
Fruit-for-snack store ⁱ	2.3	2.7	.80	2.9	.62	1.8	.64
Non-F&V store	13.8	9.4	.22	5.8	<.001	8.9	.05
Total food store	20.3	19.4	.84	15.2	.09	17.5	.39
Weight-adjusted							
F&V store ^e , total	4.6	11.4	.01	7.8	.10	9.8	.02
Super produce ^f	0.0	7.3	<.001	2.4	.01	4.0	.001
Year-round produce ^g	4.6	4.0	.78	4.4	.90	4.2	.80
Seasonal produce ^h	0.0	0.0	<.99	1.0	.11	1.6	.04
Fruit-for-snack ⁱ	2.7	2.0	.66	3.4	.60	2.4	.80
Non-F&V store	15.4	12.0	.38	6.5	<.001	10.7	.08
Total food store	22.7	25.4	.59	17.7	.12	22.9	.96

Abbreviations: F&V, fruit and vegetable; CBGs, census block groups.

^a Region within the city of Albany containing CBGs composed of ≥50% nonwhite and/or Hispanic populations. Reference category for comparison.

^b Region within the city of Albany containing CBGs composed of <50% nonwhite and/or Hispanic populations.

^c Region within Columbia and Greene counties containing CBGs composed of 100% rural populations, as defined by the US Census Bureau (25).

^d Region within Columbia and Greene counties containing CBGs not included in the rural community.

^e Food store categorized as an F&V store if it stocked at least 2 types of fresh fruit, excluding lemons and limes, and at least 3 types of fresh vegetables, including at least 1 dark-green or orange-colored vegetable.

^f An F&V store defined as a supermarket with a produce department.

^g An F&V store defined as a nonsupermarket store that operates year-round and includes grocery stores, convenience stores, and specialty food stores.

^h An F&V store that includes seasonal farm (ie, barn) stores, roadside huts and stands, and farmers' markets.

ⁱ Defined as stores carrying at least 1 type of ready-to-eat fresh fruit, such as apples, oranges, and bananas. Although these stores did not meet the F&V measure, they were not categorized as non-F&V stores.

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the U.S. Department of Health and Human Services, the Public Health Service, the Centers for Disease Control and Prevention, or the authors' affiliated institutions. Use of trade names is for identification only and does not imply endorsement by any of the groups named above.