Non-critical healthcare tools as a potential source of healthcare-acquired bacterial infections in eastern Ethiopia: A hospital-based cross-sectional study

SAGE Open Medicine Volume 7: 1–10 © The Author(s) 2019 Article reuse guidelines: sagepub.com/journals-permissions DOI: 10.1177/2050312118822627 journals.sagepub.com/home/smo



Fitsum Weldegebreal¹^(D), Desalegn Admassu¹, Dereje Meaza² and Mulatu Asfaw³

Abstract

Background: Outbreaks of healthcare-acquired infections have been linked to contaminated medical devices such as electronic thermometers, sphygmomanometers, stethoscopes, latex gloves, masks, neckties, white coats and other.

Objective: The aim of this study was to assess non-critical healthcare tools as a potential source of healthcare-acquired bacterial infections and associated factors in public health hospitals of Harar, eastern Ethiopia from March 2016 to February 2017.

Methods: A hospital-based cross-sectional study was conducted on 212 non-critical healthcare tools owned by different health professionals. The data were collected from each owner using self-administered questionnaire. Swab specimens were collected from 187 stethoscopes and 25 sphygmomanometers using sterile cotton tips. Bacterial investigation and antimicrobial susceptibility tests were performed using standard culture tests. The data were double entered into EPI-Data version 3.1 and exported into the Statistical Package for Social Sciences version 16 for analysis.

Result: The overall prevalence of non-critical healthcare tool contamination was 53.8%. A total of 137 bacterial strains were isolated. *Staphylococcus aureus* was the most frequent isolate (35%). Resistance to two or more different classes of antimicrobial was found to be 19.3%. The proportion of stethoscopes or sphygmomanometers contamination owned by the health professionals who were not cleaned regularly before and after examining each patient was found to be high (77%). The majority of non-critical healthcare tools used by health professionals working in the intensive care unit were contaminated (75%) followed by medical wards (73.5%).

Conclusion: This study confirmed that the majority of the stethoscopes and sphygmomanometers were contaminated with pathogenic bacteria known to be associated with healthcare-acquired infections. Most of the healthcare workers did not practice stethoscope and sphygmomanometers disinfection. Strict and careful decontamination of stethoscopes and sphygmomanometers need to be in place before use.

Keywords

Stethoscopes, sphygmomanometers, healthcare-acquired bacterial infections, Harar, Ethiopia

Date received: 17 July 2018; accepted: 11 December 2018

Background

Healthcare-acquired infections (HCAIs) are a major public health problem worldwide and remain a significant hazard for hospitalized patients and healthcare workers, particularly in developing countries like Ethiopia, where resources and the awareness of infection prevention and control are limited. It has long been recognized as crucial factors bedeviling the quality and outcomes of healthcare delivery.^{1,2} Healthcare workers can transmit pathogenic microorganism including ¹Department of Medical Laboratory Sciences, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia ²School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia ³School of Medicine, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

Corresponding author:

Fitsum Weldegebreal, Department of Medical Laboratory Sciences, College of Health and Medical Sciences, Haramaya University, P.O. Box 235, Harar, Ethiopia. Email: fwmlab2000@gmail.com

Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (http://www.creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (https://us.sagepub.com/en-us/nam/open-access-at-sage). antibiotic-resistant pathogens through their hands and contaminated medical devices such as electronic thermometers, sphygmomanometers, stethoscopes, gloves, masks, neckties, pens, badges and white coats.^{2–6}

Medical equipment used in the non-critical care setting is less likely to have standard disinfection and cleaning protocols than equipment in the critical care setting. Thus, medical care equipment is more likely to carry a considerable number of pathogenic bacterial strains.7 The contamination of stethoscope particularly the diaphragm is reported mainly due to lack of regular disinfection (before and after examining each patient).⁸ Furthermore, antibiotic-resistant bacteria may be transmitted from one patient to another through medical devices.9 Stethoscope is an important instrument; if used without disinfection, it might bring the risk of infection to the patient and may continuously impose the risk serially to all patients. Stethoscopes used by medical practitioners, students and health workers have been shown to be a potential vector in the transmission of HCAIs. The disinfection of devices is not done as a routine by most health professionals.^{10,11} Draping of stethoscopes around the neck is commonly seen as a practice, resulting in the risk of recontamination of the diaphragm of the stethoscope from unclean earpieces, with normal flora and pathogenic bacterial strains harboring the ears of the healthcare workers. A single stethoscope often used for all inpatients and outpatients. The universal and unavoidable use of the stethoscope and its direct contact with multiple patients makes it an important potential factor in the dissemination of microorganisms from one patient to another.12,13

In resource-poor settings such as most developing countries, rates of HCAI exceed 20%,¹⁴ but available data are scanty and more research is urgently needed in developing and transitional countries.

Healthcare workers often overlook non-critical healthcare tools including stethoscopes and sphygmomanometers as sources of infection. Studies have, however, demonstrated a significant risk of transmission of HCAI with these devices.¹⁵

Patients who acquire HCAIs require additional diagnostic tests, therapeutic treatments and prolonged hospital stay. The Centers for Disease Control and Prevention (CDC) states that cleaning and disinfecting environmental surfaces are fundamental in reducing their potential contribution to the incidence of HCAIs.¹⁶

Identifying the bacterial pathogens causing HCAIs from non-critical healthcare tools and their antimicrobial susceptibility pattern would help to make awareness of the magnitude of the problem and inherent dangers associated with HCAIs. It would also help to reduce the exposure of the hospitalized patient to multidrug-resistant pathogens in the hospital, and thus improve the clinical condition of the patients. There was a scarcity of data on non-critical healthcare tools used by staffs and students as a potential source of healthcare-acquired bacterial infections in eastern Ethiopia. This study was aimed to assess non-critical healthcare tools (stethoscopes and sphygmomanometers) used by staffs and students as a potential source of healthcare-acquired bacterial infections and associated factors in public hospitals of Harar, eastern Ethiopia.

Materials and methods

Study area and period

The study was conducted from March 2016 to February 2017 at two public hospitals (Hiwot Fana Specialized University Hospital and Jugal Hospital). Hiwot Fana Specialized University Hospital is a large and complex teaching hospital found in Harar town. The hospital is expected to serve about 5.8 million people in the eastern part of Ethiopia. A total of 300 students from different departments in Haramaya University, College of Health and Medical Sciences, Jigjiga University, Harar Nursing College and other private institutions attached to the hospital for practical training. Jugal Hospital is found in the Harari Regional State. It also serves as a practice site for Haramaya University students. It has a total number of 120 beds.¹⁷

Study design and population

A cross-sectional study was conducted among stethoscopes and sphygmomanometers used by students and staffs in the outpatient department (OPD), pediatric, surgery, gynecology and obstetrics and internal medicine wards.

Sample size determination

The sample size for stethoscope contamination was determined using a single-population proportion formula considering the prevalence of contamination reported in Jimma University Specialized Hospital (85.8%),¹⁸ 95% confidence level and a 5% margin of error. The final sample size for stethoscope contamination was 187. While all sphygmomanometers used in selected wards of the two hospitals during study were included in the study.

Sampling technique

Health professionals (specialist, general practitioners, health officer and nurses) and students along with their respective stethoscopes were allocated proportionally to Hiwot Fana Specialized University Hospital and Jugal Hospital based on their total number. Accordingly, 149 from Hiwot Fan Specialized University Hospital and 38 from Jugal Hospital were selected using simple random sampling (lottery) method by using the attendance sheet of the respective wards as a sampling frame.

Method of data collection

A structured questionnaire developed from different kinds of literature^{2,16,18–21} was used as a data collection tool. Before

the actual data collection was pursued, the questionnaire was pretested on 5% of the sample size out of the study area to ensure its validity. The questionnaire was amended based on the pretest feedback. A questionnaire consisted of two parts: socio-demographic characteristics and infection prevention/ disinfection practice

The data were collected using a self-administered questionnaire after written consent was obtained from each participant. Then, a swab sample was collected from each stethoscope or sphygmomanometers at a convenient time for health professionals when they were free of disturbance in the hospital. The participants were not informed the date of sample collection.

Before taking a swab from the noncritical tool, both hands of data collectors were cleaned with an alcohol-based instant hand sanitizer and powder-free, disposable, gloves were worn throughout the work per sample to prevent contamination. Sterilized cotton swab moisten by sterile normal saline was rotated and swiped from overall parts of the stethoscope and sphygmomanometers' area. The swab was placed immediately in Amie's transporting media. The collected samples were coded by a unique identification number and transferred to the Bacteriology Laboratory of the Department of the Medical Laboratory Sciences, Haramaya University within 30 min of collection.^{22,23}

Culture isolation and identification of bacteria

Aseptically collected sample was streaked onto blood agar and MacConkey agar plates (Oxoid, LTD, UK) following the standard techniques described.²² The inoculated plates were incubated aerobically at 37°C for 24–48 h. Primary isolation of bacteria was made based on their colony characteristics and Gram reaction. Further identification of the bacterial organism to species level was carried using biochemical tests.^{24,25} In brief, Gram-negative bacteria were identified by doing a series of biochemical tests like triple sugar iron agar, indole, Simmon's citrate agar, oxidase, urease test, Motility– Indole–Urea (MIU) and methylene red. Mannitol salt agar, catalase and coagulase tests were used for identification of Gram-positive bacteria. Colony count \geq 20 CFU/diaphragm was considered as significant contamination.¹⁸

Antimicrobial susceptibility test

The antimicrobial susceptibility test was done using the Kirby–Bauer disk diffusion method based on the Clinical Laboratory Standards Institute (CLSI) guideline.²⁴ In brief, pure culture (4–5 colonies) were added to a sterile tube containing 5 mL of normal saline (0.85% NaCl) and mixed gently until it forms a homogeneous suspension equivalent to 0.5 McFarland standards. Sterile cotton swabs were dipped into the suspension, excess fluid was removed by gentle rotation of the swab against the inner surface of the tube, and the bacterial suspension over the entire surface of Mueller

Hinton agar (Oxoid Ltd, UK) was inoculated and left at room temperature for 3-5 min to dry. Antimicrobial disks such as cefoxitin $(30 \mu g)$, ciprofloxacin $(5 \mu g)$, amoxicillin $(30 \mu g)$, erythromycin (15 µg), norfloxacin (10 µg), cefotaxime $(30 \,\mu g)$, tetracycline $(30 \,\mu g)$, trimethoprim-sulfamethoxazole $(1.25/23.75 \,\mu\text{g})$ and vancomycin $(30 \,\mu\text{g})$ were placed at least 24 mm away from each other and 15 mm from the edge to avoid the overlapping zone of inhibition. It was gently pressed to ensure that the disk attached to the agar surface of the plate. Inoculated plates were incubated at 37°C. After overnight incubation, the diameter zone of inhibition was measured using a digital caliper. The results were read to the nearest millimeter and interpreted as resistance (R), intermediate (I) and sensitive (S) based on CLSI criteria. Bacterial isolates resistant to two or greater than two from a different class of antimicrobial group were classified as resistance to two or more antimicrobial.24

Data processing and analysis

Data were double entered into EPI-Data version 3.1 and transferred to the Statistical Package for Social Science (SPSS) software version 16 for analyzing. Descriptive statistics were used to summarize different variables and presented in the form of texts and tables. Bivariate and multivariate logistic regressions were carried out to identify factors associated with a bacterial contamination of stethoscope or sphygmomanometers. Those variables in the bivariate logistic regression analysis with a p value less than or equal to 0.25 were considered as a candidate for multivariate logistic regression analysis. A variable with p value less than 0.05 at 95% confidence interval (CI) in the multivariate logistic model was considered as statistically significant.

Quality assurance

All culture media were prepared following the manufacturer's instructions, and their sterility was checked by incubating 3%–5% of the batch at 37°C overnight and observed for growth. Culture media, which showed any growth, was rejected and replaced by a new sterile batch. The reference strains such as *Staphylococcus aureus* (ATCC25923) for blood agar and *Escherichia coli* (ATCC-25922) and *Pseudomonas aeruginosa* (ATCC-27853) for MacConkey agar were used for quality control of culture. Depending on the bacterial species, culture media was incubated aerobically at 35°C–37°C and observed for the degree of growth, size of colonies and other characteristics.

The quality of antimicrobial agents was checked against *S. aureus* (ATCC 25923) and *E. coli* (ATCC 25922). The inhibition zone was felt within the pre-set standard zone of inhibition.²⁴ The questionnaire was pretested on 5% of the sample size out of the study area to ensure its validity. Training on data collection and samples was given to data collectors by investigators before commencement of actual

data collection. The data collection was closely supervised by supervisors and the investigators. Completeness of each questionnaire was checked daily during the data collection period. Double data entry was done by two data clerks into EpiData. The consistency of the entered data was crosschecked by comparing the two separately entered data.

Table 1. Sociodemographic characteristics of study participantsin Hiwot Fana Specialized University Hospital and Jugal Hospital,Harar, Ethiopia, 2017 (n=212).

Characteristics of participants		No.	%
Gender	Male	95	44.8
	Female	117	55.2
Age group	20–29	159	75
(in years)	>30	53	25
Profession	Anesthetists	10	4.7
	Clinical specialist, general	23	10.9
	practitioner and health officers		
	Nurse	45	21.2
	Medical students	134	63.2
Department	Out-patient department	42	19.8
	Emergency	49	23.I
	Medical ward	49	23.I
	Gynecology and maternity ward	10	4.7
	Pediatrics ward	16	7.5
	Surgical ward	26	12.3
	Intensive care unit	20	9.4

Result

Sociodemographic characteristics

A total of 212 study participants along with their stethoscope or sphygmomanometer were participated in the study. The response rate was 100%; 117 (55.2%) were females. The professionals include medical students (134), clinical specialists, general practitioners and health officer (23), anesthetists (10) and nurses (45). These health professionals were working in different wards, namely, OPD (42), pediatrics ward (16), emergency ward (49), medical ward (49), surgical ward (26), gynecology and maternity wards (10) and intensive care unit (20) (Table 1).

Infection prevention/disinfection practice

A total of 187 stethoscopes and 25 sphygmomanometers owned by different health professionals were examined for bacterial contamination. The majority (76.4%) of the stethoscope was used by the professionals for more than a year. About two-third of (65.6%) healthcare workers did not practice stethoscope and sphygmomanometers disinfection before and after examining each patient; 186 (87.7%) and 136 (64.2%) of the study participants have no habit of hand washing between patient examination and with soap after using stethoscope or sphygmomanometer, respectively (Table 2).

 Table 2.
 Information on infection prevention among the study participants in Hiwot Fana Specialized University Hospital and Jugal Hospital, Harar, Ethiopia, 2017 (n=212).

Characteristics		No.	%
Types of non-critical healthcare tools	Stethoscope	187	88.2
	BP cuff	25	11.8
Length of stethoscope or sphygmomanometers	Weeks	20	9.4
used	Months (<i td="" year)<=""><td>30</td><td>14.2</td></i>	30	14.2
	Years	162	76.4
Cleaned stethoscopes or sphygmomanometers	Yes	73	34.4
before and after examining each patient	No	139	65.6
Frequency of stethoscopes or	Monthly	41	19.3
sphygmomanometers cleaning	Weekly	112	52.8
	Daily	59	27.9
Type of agent used for cleaning	Soap and water	2	0.9
	Alcohol	151	71.2
	Dry cotton	59	27.9
On duty usually stethoscope or	Around neck	60	28.3
sphygmomanometers placed	In gown's pocket	77	36.3
	In trousers pocket	29	13.7
	In bag	46	21.7
Between patient examination habit of hand	Yes	26	12.3
washing	No	186	87.7
Wash hands with soap after using stethoscopes	Yes	76	35.8
or sphygmomanometers in the hospital	No	136	64.2

The frequency of bacterial isolates

The overall prevalence of non-critical healthcare tool contamination was 53.8% (114/212). Of these, 101 (54.0%) and 13 (52%) were stethoscopes and sphygmomanometers, respectively. A total of 137 bacterial strains were isolated. *S. aureus* was the most frequent isolate (35%) among Grampositive bacteria followed by coagulase-negative staphylococcus (CoNS; 16.8%) and *Bacillus* species (8.8%). Regarding Gram-negative bacteria, *Klebsiella pneumonia* (12.4%) were the most common isolates followed by *P. aeruginosa* (7.3%), *E. coli* (7.3%), *Proteus* spp. (6.6%) and *Salmonella* spp. (5.8%).

Factors associated with the contamination of noncritical healthcare tools

The proportion of stethoscopes or sphygmomanometers contamination used by health professionals who were not cleaned regularly their stethoscopes or sphygmomanometers before and after examining each patient was high (77%). The contamination of stethoscopes or sphygmomanometer used by health professionals working in the intensive care unit (75%) and medical ward (73.5%) was high compared with other wards. In multivariate analysis, gender, occupation, professional working department, cleaning of stethoscopes or sphygmomanometers before and after examining each patient and the frequency of stethoscopes or sphygmomanometers cleaning were found to be statistically significant (p < 0.05). Stethoscopes or sphygmomanometers used by males were 1.5 times more likely contaminated compared to females (adjusted odds ratio (AOR): 1.51, 95% CI: 1.81-12.14). Stethoscopes or sphygmomanometers which were not disinfected before and after the use for examining patient was six times more likely contaminated compared to those cleaned regularly (AOR: 5.95, 95% CI: 3.09–19.04). Stethoscopes or sphygmomanometers used in the intensive care unit (AOR: 1.97, 95% CI: 1.02, 9.63) and medical ward (AOR: 1.89, 95% CI: 1.06, 8.52) were two times more likely contaminated compared to their counterparts. Stethoscopes or sphygmomanometers used by medical students were about 1.5 times more likely contaminated compared with those used by the staffs (AOR: 1.49, 95% CI: 1.42, 8.02). Stethoscopes or sphygmomanometers disinfected monthly and weekly were four times (AOR: 4.01, 95% CI: 2.00, 11.58) and two times (AOR: 2.02, 95% CI: 1.81, 8.1) more likely to be contaminated with bacteria compared to those cleaned daily, respectively (Table 3).

Antimicrobial susceptibility pattern of the isolates

Overall, 137 bacterial isolates were tested against nine different commonly used antimicrobial for the treatment of bacterial infections. All *P. aeruginosa* isolates were 100% resistant to Amoxicillin and Cefotaxime. All *P. aeruginosa* and *E. coli* isolates were 100% sensitive to Norfloxacin; 25 (25%) of the *Salmonella* spp. showed resistance to Tetracycline and Vancomycin. All *Salmonella* isolates were susceptible to Ciprofloxacin.

More than 78% of *K. pneumoniae* and coagulase-negative Staphylococcus isolates were susceptible to Cefoxitin and Ciprofloxacin. Majority of *Proteus* species, *Bacillus* species and *S. aureus* isolates were susceptible to Norfloxacin. Relatively, *S. aureus* (8.3%) showed the least resistance against Norfloxacin (Table 4).

Resistance to two or more different classes of antimicrobial groups

The overall prevalence of resistance to two or more antimicrobial was 16.8%. *S. aureus* (66.3%) was most predominantly showed resistance to two or more antimicrobial followed by coagulase-negative staphylococcus (39.1%) and *P. aeruginosa* (30%) (Table 5).

Discussion

The introduction of medical devices for the management and treatment of diseases has contributed to the emergence of HCAIs with the consequence that put the patient into poor prognosis. The introduction of such devices is not wrong by itself, instead facilitates the medical procedures, but lack of commitment by the medical personnel to the infection prevention protocols was significant.¹⁸ The implication of the findings is that the stethoscope and sphygmomanometers might be a vector playing an important role in the transmission of potentially pathogenic bacteria, as well as in the spread of antimicrobial-resistant strains in the hospital environment.

In this study, 55.6% of the stethoscopes and 60% of sphygmomanometers were contaminated with various types of bacteria which is almost similar with previous studies conducted by Kuhu Pal et al.²⁰ in Tertiary Care Hospital of Rural Bengal (52%) and Africa-Purino and his colleagues²⁶ in the Santo Tomas University Hospital (57%). The incidence of stethoscopes contamination was also in line with other observations.^{19,20,27–29} But lower compared to the previous studies reported by Marinella et al.³⁰ (100%), Wood et al.³¹ (100%), Zuliani-Maluf et al.³² (87%), Youngster et al.³³ (85.7%) and Uneke et al.²⁸ (80.1%). The variation could be due to the differences in the time of exposure, a number of patients contacted and adherence level of infection prevention protocol implementation.

In this study, the contamination of stethoscopes or sphygmomanometers used by health professionals who were not cleaned regularly before and after examining each patient was high (77%). The higher contamination was recorded among non-critical healthcare tools used by health professionals working in the intensive care unit (75%) and medical wards (73.5%). This finding was in line with Jimma

		Stethoscopes	s or BP cuff H	Crude OR (95% CI)	p value	Adjusted OR (95% CI)	p value
		Yes (%)	No (%)				
Gender	Male	63 (66.3)	32 (33.7)	1.86 (1.12–11.02)	0.021	1.51 (1.31–12.14)	0.031
Ľ	Female	51 (43.6)	66 (56.4)	_		_	
Age group (in years) 2	20–29	97 (61)	62 (39)	0.64 (0.130–2.004)	0.841		
	≥30	17 (32.1)	36 (67.9)	_			
Professions A	Anesthetists	5 (50.0)	5 (50.0)	1.01 (0.310–5.319)	0.409		
0	Clinical specialist,	12 (52.2)	II (47.8)	0.50 (0.234–2.821)	0.513		
P0 00	general practitioner and health officers						
~	Nurse	17 (38.6)	27 (61.4)	2.01 (0.97–8.45)	0.401		
2	Medical students	80 (59.7)	54 (40.3)	_			
Occupation	Student	80 (59.7)	54 (40.3)	2.75 (1.20–9.31)	0.010	1.49 (1.42–8.02)	0.011
S	Staff	34 (43.6)	44 (56.4)				
Department	ICU	15 (75)	5 (25)	2.40 (1.91–10.71)	0.012	1.97 (1.02–9.63)	0.003
ш.	Emergency	19 (38.8)	30 (61.2)	1.23 (1.01–2.79)	0.315	0.218 (0.123–1.81)	0.429
~	Medical ward	36 (73.5)	13 (26.5)	2.76 (1.16–10.60)	0.019	1.89 (1.06–8.52)	0.021
0	Gynecology and	5 (50)	5 (50)	1.21 (1.08–3.45)	0.510	0.46 (0.21–1.04)	0.621
c	maternity ward						
e.	Pediatrics ward	7 (43.8)	9 (56.2)	1.91 (1.48–4.36)	0.302	0.71 (0.28–1.86)	0.351
S	Surgical ward	15 (57.7)	II (42.3)	1.41 (1.63–3.01)	0.282	0.92 (0.54–2.37)	0.302
0	Out-patient	17 (40.5)	25 (59.5)	_		_	
O.	department						
Clean or disinfect stethoscopes or Y	Yes	7 (9.9)	66 (90.1)	_	0.001	_	0.001
sphygmomanometers before and after examining each patient	No	107 (77)	32 (23)	6.32 (4.37–19.56)		5.95 (3.09–19.04)	
Frequency of stethoscopes or	Monthly	27 (65.9)	14 (34.1)	5.31 (2.07–14.03)	0.001	4.01 (2.00–11.58)	0.002
sphygmomanometers cleaning V	Weekly	46 (41.1)	66 (58.9)	3.04 (2.54–9.02)	0.016	2.02 (1.81–8.19)	0.014
	Daily	9 (15.3)	50 (84.7)				
Between patient examination habit Y	Yes	6 (23.1)	20 (76.9)	_	0.021	_	0.013
of hand washing	No	98 (52.7)	88 (47.3)	3.12 (1.67–6.87)		2.95 (1.41–6.08)	
Wash hands with soap	Yes	34 (44.7)	42 (55.3)	1.48 (0.78–5.02)	0.09		
after using stethoscopes or	No	71 (52.2)	65 (47.8)	_			
sphygmomanometers in the hospital							

6

Isolated bacteria	Multiple antimicrobial resistance			
	Drugs for two No. (%)	Drugs for three No. (%)	Total No. (%)	
Staphylococcus aureus (n = 48)	8 (60)	3 (6.3)	(66.3)	
Coagulase-negative	7 (30.4)	2 (8.7)	9 (39.1)	
Staphylococcus (CoNS) (n = 23)			· · · · · · · · · · · · · · · · · · ·	
Klebsiella pneumonia (n = 17)	_	-	_	
Bacillus species $(n = 12)$	_	_		
Pseudomonas aeruginosa (n = 10)	3 (30)	-	3 (30)	
Proteus species (n = 9)	_	_		
Escherichia coli (n = 10)	_	_		
Salmonella species (n=8)	_	_		
Total isolates (N = 137)	18 (13.1)	5 (3.6)	23 (16.8)	

Table 5. Multiple antimicrobial resistance of bacterial isolates from Stethoscope or Sphygmomanometers of health professionals to commonly used antibiotic disks in HFSUH and Jugal Hospital, Harar, Ethiopia, 2017 (n = 137).

a high contamination rate of stethoscopes and sphygmomanometers with potential pathogens that may cause a variety of diseases. These bacterial strains were also resistant to commonly used antimicrobial agents.

This study has several limitations. The sample size of sphygmomanometers was too small. This might reduce the power of analysis to show whether there is a difference in the sphygmomanometers utilized and contamination. Only nine antimicrobial disks were used to determine antimicrobial susceptibility of the isolates that may affect the generalizability of the finding. The antimicrobial susceptibility testing was performed in vitro, which might not indicate in vivo sensitivity. In addition, the cross-sectional nature of the study limits the investigation to the level of the association between determinants and outcomes of interest. Hence, it was impossible to get information about a causal relationship with the majority of associated factors.

Conclusion

This study confirmed that the majority of the stethoscopes and sphygmomanometers were contaminated with pathogenic bacteria known to be associated with HCAIs. Furthermore, a large proportion of isolates are resistant to multiple classes of antimicrobial agents commonly prescribed in the hospital. Being male, failure to disinfect (clean) before and after examining each patient, working in the intensive care unit and medical ward, being a student, disinfecting (cleaning) monthly or weekly were factors contributed to the contamination of noncritical healthcare tools. An intervention on the actual factors contributed to the contamination of non-critical healthcare tool should be implemented to reduce the bacterial load and HCAIs. Most of the healthcare workers did not practice about stethoscope and sphygmomanometers disinfection before and after examining each patient. Providing continuous on-the-job training for healthcare workers on infection prevention, patient safety and regular disinfection of non-critical healthcare tools contributes for minimizing contamination. Strict adherence to the stethoscope and sphygmomanometers' disinfection can minimize contamination and ensure improved patient safety in the hospital environment.

Acknowledgements

We would like to thank Haramaya University for financing this research work. Our thanks also go the study participants, individuals and organization that supports us in conducting this study. F.W. designed the study; participated in data collection, analysis, interpretation and write-up; drafted the manuscript and critically revised the manuscript. D.A. designed the study; participated in data collection, analysis, interpretation and write-up; drafted the manuscript and critically revised the manuscript. D.M. designed the study; participated in data collection, analysis, interpretation and write-up; drafted the manuscript and critically revised the manuscript. D.M. designed the study; participated in data collection, analysis, interpretation and write-up; drafted the manuscript and critically revised the manuscript. M.A. designed the study; participated in data collection, analysis, interpretation and write-up; drafted the manuscript and critically revised the manuscript. M.A. designed the study; participated in data collection, analysis, interpretation and write-up; drafted the manuscript. All authors read and approved the final manuscript.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval

Ethical approval for this study was obtained from Haramaya University Colleges of Health and Medical Sciences Institutional Research and Ethical Review Committee (IHRERC23/2016). Based on the objective of the study, an official letter was sent to Hiwot Fana Specialized University Hospital and Jugal Hospitals. Confidentiality was maintained and all respondents' questionnaire anonymously prepared.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded by Haramaya University.

Informed consent

The objective of the study was explained to each study participants. Those volunteered and signed the consent form were involved in the study. Written informed consent was obtained from all subjects before the study.

ORCID iD

Fitsum Weldegebreal (D) https://orcid.org/0000-0002-7867-1483

References

- Horan TC and Gaynes RP. Surveillance of nosocomial infections. In: Mayall CG (ed.) *Hospital epidemiology and infection control*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins, 2004, pp. 1659–1702.
- Uneke CJ, Ogbonna A, Oyibo PA, et al. Bacteriological assessment of stethoscopes used by medical students in Nigeria: implications for nosocomial infection control. *World Health Popul* 2008; 10(4): 53–61.
- 3. World Health Organization. *WHO guidelines for hand hygiene in health care: first global patient safety challenge clean care is safer care.* Geneva: WHO, 2009, p. 270.
- Steinlechner C, Wilding G and Cumberland N. Microbes on ties: do they correlate with wound infection. *Ann R Coll Surg Engl* 2002; 84: 307–309.
- Kotsanas D, Scott C, Gillespie EE, et al. What's hanging around your neck. *Med J Aust* 2008; 188(1): 5–8.
- Treakle AM, Thom KA, Furuno JP, et al. Bacterial contamination of health care workers' white coats. *Am J Infect Contr* 2009; 37: 101–105.
- Cohen H, Amir J, Matalon A, et al. Stethoscopes and otoscopes: a potential vector of infection. *Fam Pract* 1997; 14(6): 446–449.
- Schabrun S and Chipchase L. Health care equipment as a source of nosocomial infection: a systematic review. *J Hosp Infect* 2006; 63(2): 239–245.
- Fenelon L, Holcroft L and Waters N. Contamination of stethoscopes with MRSA and current disinfection practices. *J Hosp Infect* 2009; 71(4): 376–378.
- Whittington AM, Whitlow G, Hewson D, et al. Bacterial contamination of stethoscopes in the intensive care unit. *Anaesthesia* 2009; 64: 620–624.
- Gopinath KG, Stanley S, Mathai E, et al. Pagers and stethoscopes as vehicles of potential nosocomial pathogens in a tertiary care hospital in a developing country. *Trop Doct* 2011; 41(1): 43–45.
- Parmar RC, Valvi CC, Sira P, et al. A prospective, randomized, double-blind study of the comparative efficacy of immediate versus daily cleaning of stethoscope using 66% ethyl alcohol. *Indian J Med Sci* 2004; 58(10): 423–430.
- Hayden KM, Bonten MM, Blom WD, et al. Reduction in the acquisition of vancomycin-resistant enterococcus after enforcement of routine environmental cleaning measures. *Clin Infect Dis* 2006; 42(13): 1552–1560.
- Pittet D, Allegranzi B and Storr J. The WHO clean care is safer care programme: field-testing to enhance sustainability and spread of hand hygiene improvements. *J Infect Public Health* 2008; 1(1): 4–10.
- Schabrun S and Chipchase L. Healthcare equipment as a source of nosocomial infection: a systematic review. *J Hosp Infect* 2006; 63(2): 239–245.

- Sehulster L and Chinn RY. Guidelines for environmental infection control in health-care facilities. *MMWR Recomm Rep* 2003; 52(RR-10): 1–42.
- 17. Hiwot Fana specialized teaching hospital and Jugal Hospital annual report, 2015 (unpublished).
- Shiferaw T, Beyene G, Kassa T, et al. Bacterial contamination, bacterial profile and antimicrobial susceptibility pattern of isolates from stethoscopes at Jimma University Specialized Hospital. *Ann Clin Microbiol Antimicrob* 2013; 12: 39.
- Wilkins RL, Restrepo RD, Bourne KC, et al. Contamination level of stethoscopes used by physicians and physician assistants. *J Physician Assist Educ* 2007; 18(2): 41–43.
- Pal K, Chatterjee R, Biswas A, et al. Bacterial contamination and disinfection of stethoscopes: a knowledge gap among health care personnel of a tertiary care hospital of Rural Bengal. *IOSR-JDMS* 2015; 14(7): 44–49.
- Uneke CJ, Ogbonna A, Oyibo PG, et al. Bacterial assessment of stethoscope used by medical students in Nigeria: implication for nosocomial infection control. *Health Care Quar* 2009; 12(3): 132–138.
- Cheesbrough M. District laboratory practice in tropical countries, part 2. 2nd ed. Cambridge: Cambridge University Press, 2006, pp. 105–114.
- Shooriabi M, Chabi A, Satvati R, et al. Investigating the ratio and type of bacterial contamination of dentists' mobile phones in dentistry unit of Sina hospital in Ahvaz. *IJMHS* 2016; 5(8): 317–325.
- Clinical Laboratory Standard Institute (CLSI). Performance standards for antimicrobial disk susceptibility tests (CLSI Document M100-S25). 25th ed. Wayne, IL: Clinical and Laboratory Standards Institute, 2015.
- Verma K, Barasa A, Dara D, et al. Isolation and characterization of bacteria from mobile phones of students and employees at the University of Gondar, Ethiopia. *Bullet Pharmaceut Res* 2015; 5(3): 96–100.
- Africa-Purino FC, Dy ER and Coronel R. Stethoscopes: a potential source of nosocomial infections. *Phil J Microbiol Infect Dis* 2000; 29(2): 9–13.
- Pandey A, Asthana AS, Tiwari R, et al. Physicians accessories: doctor what you carry is every patient's worry? *Indian J Pathol Microbiol* 2010; 53(4): 711–713.
- Uneke CJ, Ogbonna A, Oyibo PG, et al. Bacterial contamination of stethoscopes used by health workers: public health implications. *J Infect Dev Ctries* 2010; 4(7): 436–441.
- 29. Singh G, Urhekar AD, Hodiwala AV, et al. Bacterial contamination of stethoscopes used by health care workers in a tertiary care hospital in Navi Mumbai. *Int J Pharma Bio Sci* 2013; 3(1): 186–193.
- Marinella M, Pierson C and Chenoweth C. The stethoscope, a potential source of nosocomial infection. *Arch Intern Med* 1997; 157(7): 786–790.
- Wood MW, Lund RC and Stevenson K. Bacterial contamination of stethoscopes with antimicrobial diaphragm covers. *Am J Infect Control* 2007; 35(4): 263–266.
- 32. Zuliani Maluf ME, Maldonado AF, Bercial ME, et al. Stethoscope: a friend or an enemy? *Sao Paulo Med J* 2002; 120: 13–15.
- 33. Youngster I, Berkovitch M, Heyman E, et al. The stethoscope as a vector of infectious diseases in the paediatric division. *Acta Paediatr* 2008; 97(9): 1253–1255.

- Parmar RC, Valvi CC, Sira P, et al. A prospective, randomized, double-blind study of comparative efficacy of immediate versus daily cleaning of stethoscope using 66% ethyl alcohol. *Indian J Med Sci* 2004; 58(10): 423–430.
- 35. Nunez S, Moreno A, Green K, et al. The stethoscope in the emergency department: a vector of infection. *Epidemiol Infect* 2000; 124(2): 233–237.
- 36. Bernard L, Kereveur A, Durand D, et al. Bacterial contamination of hospital physicians' stethoscopes. *Infect Control Hosp Epidemiol* 1999; 20(9): 626–628.
- Kilic IH, Ozaslan M, Karagoz I, et al. The role of stethoscopes in the transmission of hospital infections. *Afr J Biotechnol* 2011; 10(30): 5769–5772.
- Whittington AM, Whitlow G, Hewson D, et al. Bacterial contamination of stethoscopes in the intensive care unit. *Anaesthesia* 2009; 64: 620–624.
- Gebre-Sealsssie S. Antimicrobial resistance patterns of clinical bacterial isolates in Southwestern Ethiopia. *Ethiop Med J* 2007; 45(4): 363–370.