

Association between leisure-time physical activity and sedentary behavior with cardiometabolic health in the ELSA-Brasil participants

Francisco José Gondim Pitanga¹ , Sheila Maria Alvim Matos²,
Maria da Conceição C. Almeida³, Ana Luísa Patrão²,
Maria del Carmen Bisi Molina⁴ and Estela ML Aquino²

Abstract

Objectives: To assess associations, both individually and in combination, between leisure-time physical activity and sedentary behavior, and cardiometabolic health.

Methods: This cross-sectional study included 13,931 civil servants participating in the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). Leisure-time physical activity was analyzed using the leisure-time domain of the long-form International Physical Activity Questionnaire, while questions related to cumulative sitting time and leisure-based screen time on a weekday and on one day on the weekend were used to establish sedentary behavior. Data analysis was performed using multivariate logistic regression.

Results: Following adjustment for confounding variables, high levels of leisure-time physical activity and low levels of sedentary behavior were both associated with favorable cardiometabolic health markers in both genders. When these two factors were analyzed in conjunction, taking the combination of low levels of leisure-time physical activity and high levels of sedentary behavior as the reference, the inverse associations with cardiometabolic variables became even more significant.

Conclusion: High levels of leisure-time physical activity and low levels of sedentary behavior were both inversely associated with the cardiometabolic variables analyzed; however, the two variables when evaluated in conjunction appear to produce more consistent associations, particularly when sedentary behavior is evaluated according to leisure-based screen time.

Keywords

Leisure-time physical activity, sedentary behavior, cardiometabolic health

Date received: 7 September 2018; accepted: 7 January 2019

Introduction

Chronic non-communicable diseases, including cardiometabolic disorders, constitute a public health issue worldwide and represent one of the principal sources of disease burden.¹ The increasing prevalence rates of obesity and other cardiometabolic disorders have led to an expressive volume of studies worldwide focusing on the behavioral risk factors involved in the genesis and progression of these diseases. Most of these studies deal with issues related to lifestyle, diet, and physical activity, with few having focused on factors associated with sedentary behavior, principally in a combined or comparative analysis with leisure-time physical activity.²

¹Department of Physical Education, School of Education, Federal University of Bahia (UFBA), Salvador, Brazil

²Institute of Collective Health, Federal University of Bahia (UFBA), Salvador, Brazil

³Instituto Gonçalo-Moniz Research Center, Oswaldo Cruz Foundation (FIOCRUZ), Salvador, Brazil

⁴Federal University of Espírito Santo (UFES), Vitória, Brazil

Corresponding author:

Francisco José Gondim Pitanga, Department of Physical Education, School of Education, Federal University of Bahia (UFBA), Salvador, Brazil Av. Reitor Miguel Calmon, s/n, Vale do Canela, 40.000, Salvador, Bahia, Brazil. Email: pitanga@lognet.com.br



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Sedentary behavior refers to any activity characterized by very low energy expenditure, that is, not exceeding 1.5 metabolic equivalents, and includes specific behaviors such as sitting for reading, studying, and watching television. Generally, these habits are considered different from practicing small amounts of physical activity, a behavior in which the individual also fails to engage in moderate or vigorous physical activity that would require energy expenditure above 3 metabolic equivalents.³

More recently, the definition of sedentary behavior was broadened after a group of investigators reached a consensus defining sedentary behavior as any waking behavior characterized by energy expenditure ≤ 1.5 metabolic equivalents while in a sitting, reclining, or lying position.⁴

Various studies have shown an association between more time spent in sedentary activities and a greater occurrence of cardiovascular disease,⁵ type 2 diabetes,⁶ obesity,⁷ and metabolic syndrome,⁸ as well as a greater risk of death from cardiovascular disorders irrespective of physical activity level.^{9,10} There is evidence that excess sedentary behavior such as television viewing and computer use represents a relevant risk factor and merits as much investigation as that conducted on physical activity levels. Sitting for prolonged periods of time constitutes an important risk factor for all causes of mortality irrespective of the regular practice of physical activity.¹¹

On the other hand, various studies have shown that physical activity, particularly when performed within the domain of leisure time, is inversely associated with cardiometabolic diseases including diabetes, dyslipidemias, hypertension, and inflammatory markers.^{12–17}

Despite the evidence of inverse associations between the variables physical activity , $\text{sedentary behavior}$, and $\text{cardiometabolic health}$, evaluated individually, and cardiometabolic variables, studies aimed at evaluating which of these behaviors plays a more significant role in the control of these diseases are sparse. It has already been documented that individuals with low levels of physical activity and high levels of sedentary behavior are more likely to develop cardiovascular disease compared to individuals with low levels of both physical activity and sedentary behavior.¹⁸

Establishing a quantitative association between leisure-time physical activity, sedentary behavior, and cardiometabolic health may encourage public health managers to review educational messages that could encourage both an increase in physical activity and a reduction in sedentary behavior.

Therefore, the objective of this study was to assess the individual and combined associations of leisure-time physical activity and sedentary behavior, and cardiometabolic health.

Methods

Design and sample

The ELSA-Brasil is a cohort study involving 15,105 active or retired civil servants of 35–74 years of age from six different institutions of higher learning located in the cities of

Salvador, Vitória, Belo Horizonte, Rio de Janeiro, São Paulo, and Porto Alegre. Sample size estimation was based on the main study outcomes of type 2 diabetes and myocardial infarction. Considering an alpha value of 5%, statistical power of 80%, exposure prevalence of 20%, and a relative risk of 2.0, the required sample size was estimated at approximately 6400 individuals. In order to perform gender-specific analyses and allow for possible losses to follow-up, the desired sample size was approximately 15,000 individuals.¹⁹ Details of the methodology of this study have already been published elsewhere.^{19,20} All the participants who took part in the second wave of the study (2012–2014) and whose data on leisure-time physical activity and sedentary behavior were complete were selected for the present analysis: a total of 13,931 participants of both sexes.

The internal review boards of the six research centers involved in the ELSA-Brasil approved the study protocol. All the participants signed an informed consent form. The use of anonymized data guaranteed the confidentiality of the participants.

Measures

The data were produced by a team of interviewers and then verified by personnel trained and certified by a quality control committee.²⁰ The supervisory personnel were authorized to apply the study protocol in any of the ELSA-Brasil study centers. The interviews were conducted face-to-face, with blocks of questionnaires being applied.

Evaluation of physical activity

The International Physical Activity Questionnaire (IPAQ) was used to identify and quantify physical activity. This instrument is made up of questions on the frequency and duration of physical activities (i.e. moderate and vigorous physical activity and walking) performed during occupational activities and in household chores or for transport and during leisure time.²¹ In the ELSA-Brasil, only the domains of leisure-time physical activity and physical activity for transport were evaluated. To measure physical activity, the weekly frequency was multiplied by the duration of each one of the activities performed, with the results obtained being expressed in minutes per week. For the effects of this analysis, only the leisure-time domain was used, with leisure-time physical activity being classified as follows: 0 = insufficiently active (< 150 min/week of moderate physical activity or walking and/or < 60 min/week of vigorous physical activity or < 150 min/week of any combination of walking, moderate or vigorous physical activity); and 1 = physically active (≥ 150 min/week of moderate physical activity or walking and/or ≥ 60 min/week of vigorous physical activity or ≥ 150 min/week of any combination of walking, moderate or vigorous physical activity).

Evaluation of sedentary behavior

Sedentary behavior was documented for the first time in the second wave of the study. Participants were asked questions on the number of hours they spent sitting down (cumulative sitting time) and the number of hours they spend watching television, playing video games, and using a mobile phone or computer (leisure-based screen time) on a weekday and on one day on the weekend. A low level of sedentary behavior was classified as ≤ 2 h/day of leisure-based screen time¹⁰ and ≤ 8 h/day of cumulative sitting time.¹¹

Evaluation of cardiometabolic variables

The following cardiometabolic variables were analyzed: obesity, low levels of high-density lipoprotein cholesterol (HDL-C), hypertriglyceridemia, and hypertension. The data on medical assessments and clinical and subclinical parameters included in this analysis were obtained in the second wave of the ELSA-Brasil visits.²² Body mass index (BMI) was defined as weight in kilograms divided by height in square meters. Blood pressure was taken using a validated oscillometric device (Omron HEM 705CPINT) after a 5-min resting period, with the subject in a seated position in a quiet, temperature-controlled room (20°C–24°C). Three measurements were taken at 1-min intervals. The mean of the last two measurements was used for the analyses. Triglyceride and HDL-C levels were determined using enzymatic colorimetric methods.

A participant was classified as hypertensive if systolic blood pressure was ≥ 140 mmHg, if diastolic blood pressure was ≥ 90 mmHg, or if he or she had taken any medication to treat hypertension in the preceding 2 weeks. Obesity was defined as BMI ≥ 30.0 kg/m². Hypertriglyceridemia was defined as values > 150 mg/dL and low HDL-C as values < 35 mg/dL.

Data analysis procedures

Descriptive measures (proportions) were calculated for all the categorical variables. All the analyses were stratified by sex. This decision is supported by the scientific literature in which evidence has accumulated on gender differences in the use of time in work and at leisure.²³ The associations between the dependent variables (the cardiometabolic variables) and the independent variables (leisure-time physical activity, sedentary behavior, and their combinations), either individually or in conjunction, were analyzed using multivariate logistic regression. The following variables were considered potential confounding factors: age, BMI, education level, beer consumption, and smoking. Variables evaluated simultaneously (tetrachoric correlation) with correlation coefficient $\rho < 0.60$ and $p \leq 0.05$ at bivariate analysis were selected for the model.

Associations were sought between leisure-time physical activity alone and the cardiometabolic variables (obesity, low HDL-C, hypertriglyceridemia, and arterial hypertension) and between sedentary behavior alone and the same

cardiometabolic variables. Next, logistic regression models were proposed using combinations of leisure-time physical activity and low levels of sedentary behavior (cumulative sitting time and leisure-based screen time on a weekday and on a Saturday or Sunday), with the cardiometabolic variables as the endpoint. In all the analysis models, the combination of low levels of physical activity and high levels of sedentary behavior was defined as the reference. A confidence interval of 95% was established. The STATA software program, version 12.0 was used for the statistical analysis, which was performed in 2016.

Results

Overall, 6324 men and 7607 women were included in the analysis. The characteristics of the sample are shown in Table 1. The men were more likely to be smokers and consumed more beer on weekdays than the women. In addition, they were more likely to have hypertriglyceridemia and hypertension. Furthermore, although they were more active in their leisure time, the men tended to sit for longer periods of time on weekends and tended to watch more television, both during the week and on weekends. Women had better education levels and were more likely to be obese, with low HDL-C levels. There were no statistically significant differences between men and women in relation to their age or to the amount of weekday sitting time.

The associations between leisure-time physical activity and sedentary behavior, and cardiovascular variables are shown in Table 2 for both men and women. In men, associations were found between leisure-time physical activity and all the cardiometabolic endpoints analyzed except for hypertension. While in men sedentary behavior was strongly associated with obesity and hypertriglyceridemia, in women it was strongly associated with obesity alone. It should be noted that weekday leisure-based screen time was associated with all the cardiometabolic variables analyzed, in both men and women.

Tables 3 and 4 show the associations between the combined factors (leisure-time physical activity and sedentary behavior) and the cardiometabolic variables in men and women, respectively. Inverse associations were found for all the cardiometabolic variables, particularly when increased leisure-time physical activity and decreased weekday leisure-based screen time were analyzed together. These inverse associations were stronger in the combined analyses than when each variable was analyzed individually.

Discussion

This study sought to assess the individual and combined associations of leisure-time physical activity and sedentary behavior, and cardiometabolic health. Although the associations found when each variable was analyzed alone were statistically significant, the combinations of high

Table 1. Characteristics of the sample according to the study variables (Longitudinal Study of Adult Health (ELSA-Brasil, 2012–2014).

	Men (n=6.324)	Women (n=7.607)	p-value*
Age (years), n (%)			
34–50	2060 (32.4)	2350 (30.7)	0.06
51–60	2193 (34.5)	2762 (36.1)	
>60	2104 (33.1)	2545 (33.2)	
Education, n (%)			
Failed to complete elementary school	452 (7.1)	268 (3.5)	<0.01
Elementary school	488 (7.7)	390 (5.1)	
High school	1941 (30.6)	2456 (32.1)	
University/postgraduate	3467 (54.6)	4540 (59.2)	
Smoking, n (%)			
Never-smoker	3297 (51.9)	4890 (63.9)	<0.01
Smoker/former smoker	3049 (48.1)	2761 (36.1)	
Weekly beer consumption, n (%)			
<750 mL	4018 (63.2)	6638 (86.7)	<0.01
≥750 mL	2339 (36.8)	1019 (13.3)	
Leisure-time physical activity (M ± SD)	181.0 ± 242.3	130.2 ± 187.0	<0.01
n (%)			
Insufficiently active	3307 (52.3)	4742 (62.4)	<0.01
Active	3017 (47.7)	2865 (37.6)	
Sedentary behavior			
Weekday sitting time (M ± SD)	5.9 ± 3.4	5.8 ± 3.5	0.05
n (%)			
>8 h	1340 (21.1)	1642 (21.4)	0.60
≤8 h	5017 (78.9)	6015 (78.6)	
Weekend sitting time (M ± SD)	4.9 ± 3.1	4.4 ± 2.9	<0.01
n (%)			
>8 h	737 (11.6)	645 (8.4)	<0.01
≤8 h	5620 (88.4)	7012 (91.6)	
Weekday screen time at leisure (M ± SD)	2.3 ± 1.9	2.2 ± 1.7	<0.01
n (%)			
>2 h	2355 (37.0)	2627 (34.3)	<0.01
≤2 h	4002 (63.0)	5030 (65.7)	
Weekend screen time at leisure (M ± SD)	3.2 ± 2.2	2.9 ± 2.1	<0.01
n (%)			
>2 h	3669 (57.7)	3987 (52.1)	<0.01
≤2 h	2688 (42.3)	3670 (47.9)	
Obesity			
No	4842 (76.6)	5385 (71.0)	<0.01
Yes	1477 (23.4)	2200 (29.0)	
Hypertriglyceridemia			
No	4082 (64.6)	5998 (78.8)	<0.01
Yes	2237 (35.4)	1611 (21.2)	
Hypertension			
No	3475 (54.9)	4775 (62.6)	<0.01
Yes	2855 (45.1)	2849 (37.4)	
Low levels of HDL-C			
No	4931 (78.0)	5569 (73.2)	<0.01
Yes	1388 (22.0)	2040 (26.8)	

HDL-C: high-density lipoprotein cholesterol.

*Values for men and women were compared using the chi-square test for categorical variables and Student t-test for continuous variables.

levels of leisure-time physical activity and less sitting time and of high levels of leisure-time physical activity and less leisure-based screen time, both on weekdays and

on the weekend, resulted in more significant associations with the cardiometabolic endpoints analyzed, both in men and in women.

Table 2. Associations independent between leisure-time physical activity and sedentary behavior with cardiometabolic disorders in men and women (ELSA-Brasil, 2012–2014).

	Obesity ^a	Low levels of HDL-C ^b	Hypertriglyceridemia ^b	Hypertension ^b
<i>Men</i>				
Leisure-time physical activity				
Insufficiently active	1.00	1.00	1.00	1.00
Active	0.71 (0.62–0.82)*	0.76 (0.67–0.86)*	0.77 (0.69–0.85)*	0.90 (0.81–1.00)
Sedentary behavior				
Weekday sitting time				
>8h	1.00	1.00	1.00	1.00
≤8h	0.76 (0.64–0.91)*	0.92 (0.80–1.07)	0.84 (0.74–0.96)*	1.02 (0.89–1.17)
Weekend sitting time				
>8h	1.00	1.00	1.00	1.00
≤8h	0.72 (0.58–0.90)*	0.93 (0.77–1.12)	0.77 (0.66–0.91)*	0.92 (0.78–1.09)
Weekday screen time at leisure				
>2h	1.00	1.00	1.00	1.00
≤2h	0.68 (0.59–0.78)*	0.87 (0.77–0.99)*	0.68 (0.61–0.76)*	0.82 (0.74–0.92)*
Weekend screen time at leisure				
>2h	1.00	1.00	1.00	1.00
≤2h	0.66 (0.57–0.76)*	0.89 (0.79–1.01)	0.78 (0.70–0.87)*	0.86 (0.77–0.95)*
<i>Women</i>				
Leisure-time physical activity				
Insufficiently active	1.00	1.00	1.00	1.00
Active	0.57 (0.50–0.65)*	0.79 (0.70–0.88)*	0.79 (0.70–0.89)*	0.86 (0.77–0.96)*
Sedentary behavior				
Weekday sitting time				
>8h	1.00	1.00	1.00	1.00
≤8h	0.72 (0.71–0.94)*	1.09 (0.96–1.25)	1.06 (0.92–1.23)	1.11 (0.98–1.27)
Weekend sitting time				
>8h	1.00	1.00	1.00	1.00
≤8h	0.87 (0.62–0.95)*	1.04 (0.86–1.27)	1.01 (0.83–1.24)	1.05 (0.87–1.26)
Weekday screen time at leisure				
>2h	1.00	1.00	1.00	1.00
≤2h	0.76 (0.67–0.85)*	0.88 (0.79–0.98)*	0.75 (0.67–0.85)*	0.85 (0.77–0.95)*
Weekend screen time at leisure				
>2h	1.00	1.00	1.00	1.00
≤2h	0.76 (0.70–0.88)*	0.98 (0.88–1.08)	1.03 (0.92–1.15)	1.00 (0.90–1.10)

HDL-C: high-density lipoprotein cholesterol.

^aAdjusted for age, education level, beer consumption, and smoking.^bAdjusted for age, obesity, education level, beer consumption, and smoking.

*Statistically significant.

In this study, inverse associations were found both between obesity and leisure-time physical activity and between obesity and lower levels of sedentary behavior; however, these associations were stronger when the combined effect of the two variables was analyzed, both for men and women.

The present findings are in agreement with the results of other studies conducted recently that analyzed the combined effect of physical activity and sedentary behavior on cardiometabolic endpoints. For example, a cohort study carried out in London with 3670 participants reported a lesser likelihood

of obesity in individuals reporting a high level of physical activity and less leisure-time sitting.¹⁸

The mechanisms that contribute toward making this combination more effective in preventing obesity remain unclear. Lower levels of sedentary behavior may reinforce the protective effects of a greater amount of physical activity, probably through independent mechanisms. The combination of higher levels of physical activity and lower levels of sedentary behavior may also represent an increase in energy expenditure, with the combination of less physical activity

Table 3. Associations between the combined factors *leisure-time physical activity* and *sedentary behavior* with cardiometabolic disorders in men (ELSA-Brasil, 2012–2014).

Combinations of LTPA and sedentary behavior	Obesity ^a	Low levels of HDL-C ^b	Hypertriglyceridemia ^b	Hypertension ^b
Weekday sitting time				
Little LTPA—Much sitting time (n = 705)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 2412)	0.55 (0.44–0.70)*	0.70 (0.57–0.86)*	0.66 (0.55–0.80)*	0.89 (0.74–1.07)
Little LTPA—Little sitting time (n = 2602)	0.76 (0.61–0.95)*	0.91 (0.74–1.10)	0.89 (0.75–1.07)	0.99 (0.83–1.20)
Active—Much sitting time (n = 605)	0.70 (0.51–0.98)*	0.73 (0.56–0.96)*	0.83 (0.66–1.04)	0.86 (0.67–1.10)
Weekend sitting time				
Little LTPA—Much sitting time (n = 411)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 2721)	0.53 (0.40–0.71)*	0.77 (0.60–0.98)*	0.64 (0.51–0.79)*	0.79 (0.63–0.99)*
Little LTPA—Little sitting time (n = 2896)	0.72 (0.55–0.95)*	1.01 (0.79–1.29)	0.83 (0.67–1.03)	0.86 (0.69–1.08)
Active—Much sitting time (n = 296)	0.70 (0.45–1.10)	0.94 (0.64–1.36)	0.92 (0.67–1.27)	0.78 (0.56–1.08)
Weekday screen time at leisure				
Little LTPA—Much sitting time (n = 1264)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 1957)	0.49 (0.40–0.59)*	0.69 (0.58–0.81)*	0.52 (0.45–0.61)*	0.76 (0.65–0.89)*
Little LTPA—Little sitting time (n = 2043)	0.66 (0.55–0.79)*	0.87 (0.73–1.02)	0.66 (0.57–0.77)*	0.82 (0.71–0.96)*
Active—Much sitting time (n = 1060)	0.70 (0.57–0.87)*	0.75 (0.61–0.92)*	0.75 (0.66–0.89)*	0.89 (0.75–1.07)
Weekend screen time at leisure				
Little LTPA—Much sitting time (n = 1989)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 1369)	0.48 (0.39–0.60)*	0.71 (0.60–0.85)*	0.62 (0.53–0.73)*	0.80 (0.69–0.94)*
Little LTPA—Little sitting time (n = 1318)	0.66 (0.55–0.80)*	0.87 (0.72–1.00)	0.81 (0.70–0.94)*	0.83 (0.71–0.96)*
Active—Much sitting time (n = 1648)	0.74 (0.62–0.89)*	0.71 (0.60–0.83)*	0.79 (0.69–0.91)*	0.87 (0.76–1.01)

LTPA: leisure-time physical activity; HDL-C: high-density lipoprotein cholesterol.

^aAdjusted for age, education level, beer consumption, and smoking.^bAdjusted for age, obesity, education level, beer consumption, and smoking.

*Statistically significant.

Table 4. Associations between the combined factors *leisure-time physical activity* and *sedentary behavior* with cardiometabolic disorders in women (ELSA-Brasil, 2012–2014).

Combinations of LTPA and sedentary behavior	Obesity ^a	Low levels of HDL-C ^b	Hypertriglyceridemia ^b	Hypertension ^b
Weekday sitting time				
Little LTPA—Much sitting time (n = 1041)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 2314)	0.47 (0.39–0.57)*	0.88 (0.74–1.05)	0.83 (0.69–1.01)	0.94 (0.79–1.12)
Little LTPA—Little sitting time (n = 3701)	0.80 (0.68–0.95)*	1.13 (0.96–1.33)	1.08 (0.91–1.29)	1.08 (0.92–1.27)
Active—Much sitting time (n = 551)	0.50 (0.37–0.67)*	0.76 (0.60–1.00)	0.83 (0.64–1.10)	0.78 (0.61–1.00)
Weekend sitting time				
Little LTPA—Much sitting time (n = 389)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 2659)	0.44 (0.36–0.57)*	0.87 (0.68–1.11)	0.79 (0.61–1.02)	0.85 (0.67–1.08)
Little LTPA—Little sitting time (n = 4353)	0.76 (0.59–0.97)*	1.14 (0.90–1.45)	0.99 (0.77–1.27)	0.96 (0.76–1.20)
Active—Much sitting time (n = 206)	0.55 (0.35–0.87)*	0.83 (0.55–1.26)	0.73 (0.47–1.14)	0.68 (0.45–1.02)
Weekday screen time at leisure				
Little LTPA—Much sitting time (n = 1617)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 1905)	0.42 (0.36–0.51)*	0.69 (0.58–0.81)*	0.56 (0.47–0.67)*	0.74 (0.63–0.86)*
Little LTPA—Little sitting time (n = 3125)	0.73 (0.64–0.84)*	0.95 (0.83–1.09)	0.79 (0.67–0.90)*	0.81 (0.71–0.93)*
Active—Much sitting time (n = 960)	0.55 (0.45–0.67)*	0.87 (0.72–1.05)	0.82 (0.68–0.99)*	0.81 (0.68–0.97)*
Weekend screen time at leisure				
Little LTPA—Much sitting time (n = 2490)	1.00	1.00	1.00	1.00
Active—Little sitting time (n = 1418)	0.44 (0.37–0.53)*	0.76 (0.64–0.89)*	0.80 (0.68–0.96)*	0.87 (0.75–1.02)
Little LTPA—Little sitting time (n = 2252)	0.80 (0.70–0.91)*	1.02 (0.90–1.16)	1.02 (0.89–1.17)	0.99 (0.87–1.12)
Active—Much sitting time (n = 1447)	0.58 (0.49–0.68)*	0.82 (0.70–0.96)*	0.78 (0.66–0.92)*	0.86 (0.74–1.00)

LTPA: leisure-time physical activity; HDL-C: high-density lipoprotein cholesterol.

^aAdjusted for age, education level, beer consumption, and smoking.^bAdjusted for age, obesity, education level, beer consumption, and smoking.

*Statistically significant.

75% reduction in the capacity to absorb fat from the bloodstream, particularly by the skeletal muscles.³³

In this context, a recent study conducted with humans has shown that short breaks in sitting time at work can result in mild to moderate reductions in total cholesterol, triglycerides, and fasting blood glucose.³⁴

Thus, the combination of an increase in physical activity and a reduction in sedentary behavior may maximize positive effects on dyslipidemias, arterial hypertension, and obesity, resulting in more consistent inverse associations with cardiometabolic disorders.

One of the strong points of this study is that this is a cohort of volunteer civil servants who, although not representative of the general population, make up a significant number of participants from six different Brazilian state capitals. Nevertheless, these results must be interpreted with caution when extrapolating the findings to the entire adult population of Brazil, since these data refer to a population of workers who share similar characteristics, but that differ from those of the general population of the country. In addition, the possibility of bias in this sample of healthy workers cannot be ruled out.

In addition, the fact that the study is cross-sectional does not allow any cause-effect relationship between the variables to be established. Another possible limitation of the study refers to the data collected on physical activity and sedentary behavior, which was obtained using questionnaires. Nevertheless, this same instrument has been widely used in studies conducted in Brazil and abroad. It is also important to mention that the ELSA-Brasil is a longitudinal study and the incorporation of a more objective measure, accelerometry, is planned. This would increase the validity of the data, both on physical activity and on sedentary behavior.

Conclusion

The results of this study may represent an important contribution to public health in that the management of public policies aimed at promoting health may be improved by making further information available on the importance both of increasing leisure-time physical activity and decreasing sedentary behavior. The information that the combined effect of these two behaviors maximizes the associations with the variables that reflect cardiometabolic health should be taken into consideration by public health managers so that actions to encourage the practice of physical activity, particularly in leisure time, can be implemented, in addition to incorporating actions to encourage individuals to reduce sedentary behavior, particularly the amount of leisure-based screen time.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethical approval

The ELSA-Brasil was approved by internal review boards from all the six research centers involved. The study was approved by the ethics committee of the Institute of Public Health of the Federal University of Bahia under number 027-06/CEP-ISC.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by Project Attraction Young Talents, Science without Borders, CAPES, project A061/2013. The ELSA-Brasil baseline study was supported by the Brazilian Ministry of Health (Science and Technology Department) and the Brazilian Ministry of Science and Technology (Financiadora de Estudos e Projetos and CNPq National Research Council; grants 01 06 0010.00 RS, 01 06 0212.00BA, 01 06 0300.00 ES, 01 06 0278.00 MG, 01 06 0115.00SP, and 01 06 0071.00 RJ).

Informed consent

All the participants signed an informed consent form. The use of anonymized data guaranteed the confidentiality of the participant.

ORCID iD

Francisco José Gondim Pitanga  <https://orcid.org/0000-0002-1033-8684>

References

- Schmidt MI, Duncan BB, Silva GA, et al. Doenças Crônicas Não-Transmissíveis no Brasil: carga e desafios atuais. *S* 2011; 377(9781): 1949–1962.
- García-Hermoso A, Martínez-Vizcaino V, Sánchez-López M, et al. Moderate-to-vigorous physical activity as a mediator between sedentary behavior and cardiometabolic risk in Spanish healthy adults: a mediation analysis. *B* 2015; 12: 78.
- Pate RR, O'Neill JR and Lobelo F. The evolving definition of "sedentary." *E* 2008; 36(4): 173–178.
- Tremblay MS, Aubert S, Barnes JD, et al. Sedentary Behavior Research Network (SBRN)—terminology consensus project process and outcome. *B* 2017; 14(1): 75.
- Jakes RW, Day NE, Khaw KT, et al. Television viewing and low participation in vigorous recreation are independently associated with obesity and markers of cardiovascular disease risk: EPIC-Norfolk population-based study. *E* 2003; 57(9): 1089–1096.
- Hu FB, Li TY, Colditz GA, et al. Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *A* 2003; 289(14): 1785–1791.
- Banks E, Jorm L, Rogers K, et al. Screen-time, obesity, ageing and disability: findings from 91 266 participants in the 45 and Up Study. *A* 2011; 14: 34–43.
- Chang PC, Li TC, Wu MT, et al. Association between television viewing and the risk of metabolic syndrome in a community-based population. *B* 2008; 8: 193.
- Stamatakis E, Hamer M and Dunstan DW. Screen-based entertainment time, all-cause mortality, and cardiovascular

- events: population-based study with ongoing mortality and hospital events follow-up. *Ar CrM CS*, 2011; 57(3): 292–299.
10. Dunstan DW, Barr EL, Healy GN, et al. Television viewing time and mortality: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *C i AS*, 2010; 121(3): 384–391.
 11. Van der Ploeg HP, Chey T, Korda RJ, et al. Sitting time and all-cause mortality risk in 222 497 Australian adults. *A*, 2012; 172(6): 494–500.
 12. Hu FB, Leitzmann MF, Stampfer MJ, et al. Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *A*, 2001; 161: 1542–1548.
 13. Ellison RC, Zhang Y, Qureshi MM, et al. Lifestyle determinants of high-density lipoprotein cholesterol: the National Heart, Lung, and Blood Institute Family Heart Study. *Ar*, 2004; 147: 529–535.
 14. Pitanga FJG and Lessa I. Associação entre Atividade Física no Tempo Livre e Proteína C-reativa em adultos na cidade de Salvador-Brasil. *A p B S-C*, 2009; 92: 302–306.
 15. Pitanga FJ and Lessa I. Relationship between leisure-time physical activity and blood pressure in adults. *A p B S-C*, 2010; 95(4): 480–484.
 16. Pitanga FJG, Matos SMA, Almeida MDC, et al. Leisure-time physical activity, but not commuting physical activity, is associated with cardiovascular risk among ELSA-Brasil participants. *A p B S-C*, 2018; 110(1): 36–43.
 17. Lin X, Alvim SM, Simoes EJ, et al. Leisure time physical activity and cardio-metabolic health: results from the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Ar*, 2016; 5(6): e003337.
 18. Bell JA, Hamer M, Batty GD, et al. Combined effect of physical activity and leisure time sitting on long-term risk of incident obesity and metabolic risk factor clustering. *D S*, 2014; 57(10): 2048–2056.
 19. Aquino EM, Barreto SM, Bensenor IM, et al. Brazilian Longitudinal Study of Adult Health (ELSA-Brasil): objectives and design. *Ar*, 2012; 175(4): 315–324.
 20. Schmidt MI, Griep RH, Passos VM, et al. Strategies and development of quality assurance and control in the ELSA-Brasil. *A S*, 2013; 47(Suppl. 2): 105–112.
 21. Matsudo S, Araújo T, Matsudo V, et al. Questionário internacional de atividade física (IPAQ): Estudo de validade e reprodutibilidade no Brasil. *A B S-A F S*, 2001; 6(2): 5–18.
 22. Mill JG, Pinto K, Griep RH, et al. Medical assessments and measurements in ELSA-Brasil. *A S*, 2013; 47(Suppl. 2): 54–62.
 23. Vari R, Scazzocchio B, D'Amore A, et al. Gender-related differences in lifestyle may affect health status. *A S*, 2016; 52(2): 158–166.
 24. Crichton GE and Alkerwi A. Physical activity, sedentary behavior time and lipid levels in the observation of cardiovascular risk factors in Luxembourg study. *A*, 2015; 14: 87.
 25. de Moraes AC, Carvalho HB, Rey-Lopez JP, et al. Independent and combined effects of physical activity and sedentary behavior on blood pressure in adolescents: gender differences in two cross-sectional studies. *A*, 2013; 8(5): e62006.
 26. Chau JY, Grunseit A, Midthjell K, et al. Cross-sectional associations of total sitting and leisure screen time with cardiometabolic risk in adults: results from the HUNT Study, Norway. *A*, 2014; 17(1): 78–84.
 27. Hamer M, Stamatakis E and Steptoe A. Effects of substituting sedentary time with physical activity on metabolic risk. *A*, 2014; 46(10): 1946–1950.
 28. Grgic J, Dumuid D, Bengoechea EG, et al. Health outcomes associated with reallocations of time between sleep, sedentary behaviour, and physical activity: a systematic scoping review of isotemporal substitution studies. *BAS*, 2018; 15(1): 69.
 29. ACSM—American College of Sports Medicine: physical activity, physical fitness, and hypertension. *A*, 1993; 25(10): i–x.
 30. Gordon PM, Goss FL, Visich PS, et al. The acute effects of exercise intensity on HDL-C metabolism. *A*, 1994; 26(6): 671–677.
 31. Edwardson CL, Gorely T, Davies MJ, et al. Association of sedentary behaviour with metabolic syndrome: a meta-analysis. *A*, 2012; 7(4): e34916.
 32. Bey L and Hamilton MT. Suppression of skeletal muscle lipoprotein lipase activity during physical inactivity: a molecular reason to maintain daily low intensity activity. *A*, 2003; 551(Pt 2): 673–682.
 33. Bey L, Akunuri N, Zhao P, et al. Patterns of global gene expression in rat skeletal muscle during unloading and low intensity ambulatory activity. *A*, 2003; 13(2): 157–167.
 34. Mailey EL, Rosenkranz SK, Casey K, et al. Comparing the effects of two different break strategies on occupational sedentary behavior in a real world setting: a randomized trial. *A*, 2016; 4: 423–428.