

AIMS Public Health, 6(4): 355–369. DOI: 10.3934/publichealth.2019.4.355 Received: 09 May 2019 Accepted: 23 September 2019 Published: 29 September 2019

http://www.aimspress.com/journal/aimsph

Research article

Associations between breastfeeding duration and overweight/obese among children aged 5–10: a focus on racial/ethnic disparities in California

Christian E. Vazquez* and Catherine Cubbin

Steve Hicks School of Social Work, The University of Texas at Austin, Austin, Texas, USA

* Correspondence: Email: christian.vazquez@utexas.edu; Tel: +8052596963.

Abstract: Research on the association between breastfeeding and childhood obesity and research on racial/ethnic differences in breastfeeding both show inconsistencies. The current study examines: 1) whether immigrant Hispanic women have higher rates of breastfeeding compared to non-Hispanic (three separate groups: African-American, Asian/Pacific Islander, and White) and U.S.-born Hispanic women; 2) whether children who were breastfed are less likely to be overweight/obese compared to children who were not breastfed; and 3) whether associations between breastfeeding and weight status vary by race/ethnicity/nativity. The study builds on prior literature using representative data from the Geographic Research On Wellbeing study (GROW, 2012 2013) and focusing on ages 5 10 years, an age group that has not been well studied (N = 2675 mother/child dyads). Logistic regression was used to investigate the odds of child obesity ($95^{th}\%$) and child overweight ($85^{th}\%$) in a series of models: unadjusted (each variable individually), demographic

race/eth), socioeconomic statusand familyincome), and full model; with breastfeeding included in all models. Interactionsbetween race/ethnicity and breastfeeding duration were also examined. African-American (9.54%) andwhite (32.8%) women had the lowest and highest rates of ever breastfeeding, respectively. Whitewomen breastfed the longest (M = 10.52 months, SE = 0.028) and U.S.-born Hispanic women breastfedthe shortest (M = 7.05 months, SE = 0.41), on average. Children of African-American and U.S.-bornHispanic mothers had higher odds of being overweight/obese (74 75%) compared with children ofwhite mothers. No associationsadjusted models

and explained the racial/ethnic disparities. These results provide evidence in favor of there being no association between breastfeeding and childhood obesity.

Keywords: Hispanic; children; health disparities; breastfeeding; obesity

1. Introduction

Recent studies reporting national childhood obesity rates in the United States suggest rates continue to be high with a steady increase in many populations [1]. One of the most prominent studies, using data from the National Health and Nutrition Examination Survey (NHANES), provided evidence that 18.5% of children were identified as obese in 2015 2016, the highest rates ever documented [1]. These rates are disproportionately higher for children of color, with recent reports showing rates as high as 25.8% and 22.0% for Hispanic and Black children, respectively, compared to White (14.0%) and Asian children (11.0%) [1,2]. High rates of childhood obesity are a major concern considering the known health risks associated with being overweight/obese, such as developing type 2 diabetes, high blood pressure, heart disease, sleep apnea, liver disease, kidney disease, and certain types of cancer [1]. Children who are overweight/obese are also more likely to have negative non-health related outcomes, including poor educational outcomes and increased risk of being bullied [3]. The negative health risks can also be detrimental economically, with one study suggesting that being overweight/obese can lead to medical costs that are 42% higher than healthy-weight individuals [4].

Research on the association between breastfeeding and childhood obesity has resulted in inconclusive findings. Several recent studies and meta-analyses in the past decade have found evidence suggesting breastfeeding is a protective factor for childhood obesity [5 7]. Further, some studies suggest this is only true for certain groups, such as less acculturated Hispanic women or WIC participants [8 10]. There are also numerous studies that found no significant relationship between the two [11 13].

The lack of standardization of the way studies are being conducted may explain the variation of findings. The meta-analyses suffer from limitations due to small sample size and having to pool different types of breastfeeding measures (e.g., exclusive breastfeeding, mixed feeding, and exclusive formula feeding). Studies range from using ever breastfed to a specific number of weeks or months breastfed, which makes it difficult to compare studies. Studies are also difficult to compare because they use samples from prospective cohort, historical cohort, or cross sectional datasets. Another issue that affects all studies, including the meta-analyses, is the lack of standard covariates included across each study, with maternal BMI being a key covariate that is commonly missing. Further, only a few studies use representative samples. It is often the case that the study has no mention of race/ethnicity,

analysis acknowledging that publication bias could not be ruled out, explaining that studies showing significant results are more likely to be published.

The American Academy of Pediatrics recommends breastfeeding as the optimal feeding practice during at least the first 6 months of

both the mother and child, including developmental, economic, health, nutritional, and social benefits [15 17]. Despite these benefits, nearly 30% of infants in the U.S. are never breastfed [18,19]. Based on the Healthy People 2020 objectives, Asian women are the only racial/ethnic group reaching the goal of breastfeeding initiation of 81.9%, although Hispanic women are close, with certain subgroups of Hispanic women (e.g. less acculturated, Mexican) meeting this goal [20]. Mothers with lower rates of breastfeeding tend to be young, with low incomes, unmarried, less educated, and overweight or obese before pregnancy [16,20]. However, studies have found that immigrant Hispanic women with high breastfeeding rates tend to be anomalies with respect to some of the factors associated with not breastfeeding, as they tend to have lower incomes, be less educated, and have healthier pre-pregnancy weight, compared to all US-born mothers with similar breastfeeding rates [16,21]. Alternatively, Wouk et al. [22] suggest Hispanic women tend to have low rates of breastfeeding, but did not examine differences among subgroups. Some studies have demonstrated that examining this relationship by subgroups of Hispanic populations is important because certain groups have different behaviors based on immigrant status or country of origin [16,23].

Although various studies have examined the effects of breastfeeding as a protective factor for childhood obesity, few studies have investigated how this relationship may be different for racial/ethnic groups, especially minority families, using representative samples. Further, less attention has focused on children 5 to 10 years of age. The guiding question asks, if breastfeeding protects against childhood overweight/obesity, and immigrant Hispanic mothers have higher rates of breastfeeding (compared to non-Hispanics and US-born Hispanics), then are children born to immigrant Hispanic mothers less likely to be overweight/obese, compared to non-Hispanics and U.S.-born Hispanics? The purpose of the current study is to provide evidence on three fronts: 1) whether immigrant Hispanic women have higher rates of breastfeeding compared to non-Hispanic and U.S.-born Hispanic women; 2) whether children who were breastfed are less likely to be overweight/obese compared to children who were not breastfed; and 3) whether associations between breastfeeding and weight status varies by race/ethnicity/nativity.

2. Methods

2.1. Data source

Data for this study came from the Geographic Research On Wellbeing (GROW, 2012 2013) study. GROW is a population-based follow-up study of mothers interviewed at baseline for the California Maternal and Infant Health Assessment (MIHA), 2003 2007. MIHA, which is very similar t annual, statewide-representative

mail/telephone survey of mothers delivering live infants in California, linked with birth certificate data (about 3500 respondents each year). Response rates for MIHA exceeded 70% each year from 2003 to 2007. Surveys were completed in English (71%) or Spanish (29%). Additional details about the MIHA survey are available [24 27].

357

Because of budget limitations, it was not feasible to follow-up all women for GROW who were interviewed at baseline. Therefore, a decision was made to follow-up MIHA respondents from six largely urbanized counties with the highest number of respondents. Respondents in these six counties represented 55% of all respondents in MIHA from 2003 to 2007. Women were eligible for GROW if they lived in one of these six counties and had agreed to be re-contacted for potential future studies (N = 9256). The GROW survey consisted of approximately 80 questions regarding demographic, socioeconomic, neighborhood, psychosocial, and health-related characteristics pertaining to herself and her index child aged 5 to 10 (her infant from the MIHA survey). The response rate was 75% among the sample of eligible women who were located (N = 3016 out of 4026 located). Over half completed the survey by phone, and nearly three-quarters completed it in English. Missing income values were imputed using hot-deck methodology and the following variables: age, race/ethnicity, education, employment status, marital status and neighborhood poverty. Weights were created to produce data that were representative of the birth file and original MIHA sample in the six GROW counties, and a sampling fraction file was created to make a minor finite population correction to the standard errors for analyses. Additional details about the GROW study are available [26].

The analytic dataset excluded women whose race/ethnicity was reported as American = 69). Other exclusions were children who did not live with the respondent at least half the time (N = 41) or whose current percentile of weight for their age was either missing (N = 204) or considered an outlier (N = 18) according to CDC standards. In addition, respondents were excluded if information was missing on breastfeeding (N = 59) resulting in 2675 records (89% of the GROW sample) for analysis.

The GROW study was approved by the Institutional Review Boards at the University of Texas at Austin, the University of California, San Francisco, and the California Department of Public Health; all participants gave informed consent.

2.2. Variables

The dependent variable was Mothers reported their either pounds or kilos, which were converted to weight percentiles according to age, per CDC guidelines. Weight status was categorized as less than 85th percent, 85th to less than 95th percent, and 95th percent or higher. Above 85th percent is considered overweight or obese, and above 95th percent is considered obese. Due to height measurements being difficult for mothers to report, weight percentile rather than body mass index (BMI) was used. In a study evaluating the accuracy of parent reported height and weight, it was determined that parents more significantly underestimated height, particularly among younger children [28]. Additionally, the literature suggests that the use of weight percentiles as a proxy for BMI is acceptable if accurate height measurements are not available [29].

The primary independent variable was breastfeeding duration, measured in two ways. First, based on the literature indicating that greater than six months of breastfeeding is beneficial for healthy weight [5,30] a categorical variable was created (none, less than seven months, or seven months or longer). Second, among women who ever breastfed (N = 2418), a continuous variable of the number of months of breastfeeding was created.

A number of sociodemographic factors were included as controls. These include

(non-Hispanic African-American, non-Hispanic Asian/Pacific Islander, immigrant Hispanic, US-born Hispanic, and non-Hispanic White),

-reported height and

weight). Annual family income was measured as the total pretax income in 2011 from all sources. In combination with the number of people supported on that income, income was converted into increments of the federal poverty level.

2.3. Analysis

First, the distributions of covariates were examined. Next, bivariate relationships between breastfeeding duration or child weight status, and race/ethnicity were examined. Chi-square or t-tests were computed to compare proportions or means. Finally, logistic regression models were used to investigate the odds of child obesity (95th percent) in a series of models: (a) unadjusted models (each variable individually); (b) a demographic model

; (c) a socioeconomic status model (demographic model plus

; (d) and a full model

BMI). All models were calculated with the categorical and continuous measure of breastfeeding, as well as with overweight or obese as the dependent variable (85th percent), for a total of 4 iterations of the four models. Interactions between race/ethnicity and breastfeeding duration were also examined. All analyses were weighted, accounted for the complex sample design, and were conducted using SAS version 9.4.

3. Results

Table 1 presents descriptive characteristics of the sample. About half of the children were female age was about seven years old.

old, and 84% were married or living together. Over half of the mothers were Hispanic, followed by white, Asian/Pacific Islander, and African-American. About 60% of women had attended at least some college, and nearly half had family incomes that were at or below 200% of the federal poverty level. Twenty-three percent of mothers had BMIs that were in the obese range. While less than 10% of mothers did not breastfeed at all, nearly half breastfed for seven months or longer. Among women who ever breastfed, the average duration was nine months. Over three quarters of children had a normal weight status, compared with 13% who were overweight and nearly 11% who were obese.

Breastfeeding duration and child weight status varied significantly by race/ethnicity (Table 2). African-American women had the highest rate of no breastfeeding (21%), and white women had the lowest rate (6%). Among women who ever breastfed, U.S.-born Hispanic women breastfed the shortest amount of time (7 months) and white women breastfed the longest (10.5 months). Children of African-American or Hispanic mothers had the highest rates of being overweight/obese (26 27%), followed by children of white mothers (20%), and Asian/Pacific Islander mothers (17%).

Table 3 presents logistic regression models for obese children, using the categorical measure of breastfeeding duration.

example, children of mothers with less than high school education had 2.4 higher odds of reporting child obesity compared with children of mothers who were college graduates. After adjusting for child

duration,

additional adjustment for education and income, older children and children of African-American or

AIMS Public Health

U.S.-born Hispanic mothers had higher odds of reporting obesity compared with their reference groups.

significant effect was for children of obese mothers, whom had nearly 2.5 higher odds of obesity compared with children of normal weight mothers. No statistically significant interaction was found

	n (M)	% (SE)
Female	1292	49.1
Male	1383	50.9
	(6.91)	(0.02)
	(28.78)	(0.13)
African-American	315	6.6
Asian/Pacific Islander	273	15.5
Hispanic Immigrant	723	34.5
Hispanic U.S. Born	425	16.5
White	939	26.9
Marital Status		
Married/living together	2228	83.9
Unmarried	435	16.1
< High School	402	18.4
High School Graduate/GED	477	22.3
Some College	654	23.6
College Graduate	1131	35.7
Family Income		
0 100 % FPL	581	28.5
101 200 % FPL	448	20.3
201 400 % FPL	509	20.5
>400 % FPL	898	30.7
0 24.99	1213	45.7
25 29.99	774	31.0
30+	571	23.3
Breastfeeding Duration		
None	234	9.2
<7 months/none	1070	42.2
7 months +	1348	48.6
Breastfeeding Duration ^a (months)	(9.33)	(0.18)
Child Weight Status		
$<\!85^{ m th}\%$	2064	76.9
$^{\rm th}$ <95 $^{\rm th}$ %	327	12.6
95 th %	284	10.5

Table 1. Descriptive Characteristics, Geographic Research on Wellbeing Study (2012 2013), N = 2675 mother/child dyads.

Note: ^a Among ever breastfed (n = 2418). n(M) = sample size and mean. % (SE) = percent and standard error.

360

Nearly identical findings were observed using the continuous measure of breastfeeding duration among women who ever breastfed (Table 4). In models of child overweight/obese, similar results were also found (Appendix Tables 1 2). The main difference was that females had lower odds of being overweight/obese compared with male children and children of mothers who were either overweight or obese both had higher odds of being overweight/obese compared with children to mothers of normal weight.

	Race/ethnicity					
	African-	Asian/Pacific	Hispanic-	Hispanic-	White	
	American	Islander	Immigrant	U.S. Born		
	% or M	% or M (SE)	% or M (SE)	% or M (SE)	% or M	² or t-
	(SE)				(SE)	test
Breastfeeding						
Duration						
None	21.4	6.4	7.0	14.8	5.9	
<7 months	42.6	43.6	43.0	53.4	35.5	67.5***
	36.0	50.0	50.0	31.7	58.6	
Breastfeeding	8.46 (0.57)	9.96 (0.57)	9.19 (0.30)	7.05 (0.41)	10.51	301.8***
Duration ^a (months)					(0.28)	
Child Weight Status						
$<\!\!85^{th}\%$	72.6	83.4	73.1	73.9	80.3	
85^{th} <95 th %	13.2	9.5	14.0	12.1	12.4	29.5***
95 th %	14.2	7.1	12.9	14.0	7.3	

Table 2. Breastfeeding Duration and Child Weight Status by Race/ethnicity, Geographic Research on Wellbeing Study (2012) 2013, N = 2675 mother/child dyads.

Note: ^{*a*} Among ever breastfed (n = 2418). % = percent. M = mean. SE = standard error. ***p < 0.001.

	Unadjusted	Demographics	SES	Full
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Female	0.79 (0.59 1.07)	0.77 (0.57 1.04)	0.76 (0.56 1.02)	0.76 (0.56 1.03)
Male	1.00	1.00	1.00	1.00
	1.12 (1.02 1.23)*	1.11 (1.01 1.22)*	1.11 (1.01 1.23)*	1.10 (1.00 1.22)
Age	0.98 (0.96 1.01)	1.01 (0.98 1.03)	1.01 (0.99 1.04)	1.01 (0.99 1.04)
African-American	1.44 (0.96 2.16)	1.87 (1.14 3.08)*	1.75 (1.05 2.92)*	1.49 (0.89 2.51)
Asian/Pacific Islander	0.55 (0.32 0.93)*	0.87 (0.49 1.56)	0.88 (0.49 1.59)	0.92 (0.51 1.67)
Hispanic-Immigrant	1.43 (1.05 1.95)*	1.88 (1.29 2.73)**	1.40 (0.86 2.27)	1.34 (0.82 2.20)
Hispanic-U.S. Born	1.45 (1.02 2.05)*	1.97 (1.27 3.06)**	1.74 (1.11 2.72)*	1.58 (1.00 2.50)
White	1.00	1.00	1.00	1.00
Marital Status				
Married/living together	1.00	1.00	1.00	1.00
Unmarried	1.39 (0.99 1.97)	1.18 (0.81 1.71)	1.14 (0.77 1.68)	1.15 (0.78 1.70)
<high school<="" td=""><td>2.37 (1.57 3.58)***</td><td></td><td>1.77 (0.99 3.17)</td><td>1.54 (0.85 2.79)</td></high>	2.37 (1.57 3.58)***		1.77 (0.99 3.17)	1.54 (0.85 2.79)

Table 3. Lo	95 th %	by	Categorical	Breastfeeding,
Geographic Research on Wellbeing Study (2012	2013),	N =	2675 mother	child dyads.

	Unadjusted Demographics		SES	Full	
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	
Female	0.80 (0.60 1.08)	0.73 (0.53 1.01)	0.72 (0.52 1.00)*	0.73 (0.52 1.01)	
Male	1.00	1.00	1.00	1.00	
	1.12 (1.02 1.23)*	1.13 (1.02 1.25)*	1.14 (1.02 1.26)*	1.12 (1.01 1.24)*	
	0.98 (0.96 1.01)	1.00 (0.98 1.03)	1.01 (0.98 1.04)	1.01 (0.98 1.04)	
African-American	1.48 (0.99 2.22)	2.00 (1.15 3.48)*	1.83 (1.04 3.21)*	1.57 (0.88 2.80)	
Asian/Pacific Islander	0.55 (0.32 0.93)*	1.02 (0.56 1.85)	1.05 (0.58 1.91)	1.09 (0.59 2.01)	
Hispanic-Immigrant	1.41 (1.04 1.92)*	2.04 (1.37	1.51 (0.90 2.54)	1.48 (0.87 2.51)	
		3.02)***			
Hispanic-U.S. Born	1.45 (1.03 2.05)*	2.03 (1.26	1.78 (1.09 2.91)*	1.58 (0.94 2.66)	
		3.26)**			
White	1.00	1.00	1.00	1.00	
Marital Status					
Married/living together	1.00	1.00	1.00	1.00	
Unmarried	1.40 (0.99 1.97)	1.31 (0.89 1.95)	1.27 (0.85 1.91)	1.28 (0.85 1.94)	
<high school<="" td=""><td>2.38 (1.58</td><td></td><td>1.83 (0.98 3.39)</td><td>1.56 (0.82 2.95)</td></high>	2.38 (1.58		1.83 (0.98 3.39)	1.56 (0.82 2.95)	
	3.58)***				
High School Graduate/GED	1.75 (1.17				
	2.61)**				

Table 4. Lo $95^{th}\%$ by Continuous Breastfeeding,Geographic Research on Wellbeing Study (2012 2013), N = 2418 mother/child dyads.

4.

Discussion

longest, with immigrant Hispanic women and Asian/Pacific Islander women close behind. In contrast, African-American women and U.S.-born Hispanic women had the highest rates of never breastfeeding and breastfed the shortest amount of time. The finding that African-American women and U.S.-born Hispanic women tend to have the worst outcomes is consistent throughout the literature [18,19]. Research has document that high-acculturated Hispanic women are less likely to intend to or breastfeed their newborn, compared with low-acculturated Hispanic women [23]. One possible explanation for this is that African-American, U.S.-born Hispanic, and immigrant Hispanic women tend to share low-income status; however, immigrant Hispanic women also tend to stay at home instead of joining the labor force, which makes it easier to have time to breastfeed. Another explanation may be the normality of using formula in the United States. Through programs, such as Women, Infants, and Children (WIC), formula is readily available and is commonly used, whereas postnatal resources in South and Central America are not as available and use of formula is not common practice [16]. Additionally, the literature provides evidence that the low prevalence of exclusive breastfeeding in low-resourced countries may be due to caregivers introducing solid food at earlier stages of infancy [31]. Further, in a review examining the relationship between childhood obesity and breastfeeding among African-Americans, barriers to breastfeeding included impact on personal and family life, including going back to work, a lack of social support, fear of impact on sexuality, fear of pain, and lack of education [32]. One of the biggest barriers to increasing breastfeeding among African-American and U.S.-born Hispanic women continues to be education opportunities, attributed to both cultural relevance of information and convenience of location to receive the information [20,33]. Concerted efforts have been made to address these issues; yet the barriers persist. This leaves a gap for innovative approaches. One such policy that is attempting to fill this gap is the increase in Baby-Friendly hospitals [34]. Baby-Friendly hospitals are designed to optimize mother-baby bonding and to protect, promote and support breastfeeding in the first few

The purpose of the current study is to provide evidence on three fronts: 1) whether immigrant Hispanic women have higher rates of breastfeeding compared to non-Hispanic and U.S.-born Hispanic women; 2) whether children who were breastfed are less likely to be overweight/obese compared to children who were not breastfed; and 3) whether associations between breastfeeding and weight status varies by race/ethnicity/nativity. As discussed above, the literature on the topics of breastfeeding rates and breastfeeding as a protective factor against obesity has rendered mixed results. The results from

In this sample, white women had the lowest rates of having never breastfed, and breastfed the

this study provide evidence in favor of one side of the debate related to each topic.

[34]. There are currently around 500 hospitals in the United States with this designation [34]. However, the societal impact of this policy has yet to be evaluated.

d that children of immigrant Hispanic women had higher odds of being overweight/obese compared to children of non-Hispanic white women; however, that difference was no longer statistically significant after controlling for education and income. Moreover, breastfeeding was not found to be a protective factor against overweight/obesity for any race/ethnicity (i.e., no significant interaction effect between race/ethnicity and breastfeeding). After adjustment for demographic and socioeconomic factors, African-American children and those of U.S.-born Hispanic women have about a 75% higher odds of obesity. The mechanism explaining the remaining racia Literature spanning over two

hood

overweight/obesity status [10,35]. Not surprisingly, research in the area of childhood obesity interventions has moved toward studying the engagement of parents as part of the weight reduction program, as this has demonstrated successful outcomes for both the parent and child [36]. As the debate on whether or not breastfeeding is a protective factor against childhood overweight/obesity continues, the current study provides evidence in favor of there being no association between the two. This also provides evidence for possible publication bias, discussed in previous studies [13,14].

The Hispanic Paradox is the phenomenon that immigrant Hispanics are healthier than U.S.-born Hispanics and non-Hispanics, despite very low socioeconomic status [37]. Several studies, including Baker et al. [21] set out to understand if the Hispanic Paradox extends from caregivers to Hispanic children [38]. They found that after a certain amount of years any health benefits children were receiving from having a healthy immigrant parent had disappeared [38]. Previous studies suggest obesity, in particular, is a health issue that appears quicker than other health issues because of the decline in healthy eating at a young age [39,40]. The results of the current study are supportive of the Paradox not protecting children of immigrants, given that children of immigrant Hispanic women have higher odds of obesity, compared to children of non-Hispanic white women.

Children born to

parks and playgrounds), and presence or lack of neighborhood resources (e.g., grocery stores), which may be a factor in child weight status.

Despite the limitations, the current study has several strengths. The GROW study is a representative sample from a large and diverse state. This study helps to fill a gap in the literature by examining the relationship between breastfeeding and obesity outcomes for children between 5 to 10 years of age. Sub-groups of Hispanic women according to immigrant status were also examined.

4.2. Implications and future research

Based on the findings of the present study, it is suggested that interventions to increase breastfeeding provide tailored supports to U.S.-born Hispanic and African-American mothers, as they tend to experience disproportionately lower levels of breastfeeding. Given high levels of overweight and obesity overall, interventions promoting healthier weight for young children of all race/ethnicities in general may benefit from including components to enhance family involvement to simultaneously reduce parental BMI. Future studies should examine cultural, occupational, and social support factors that may be barriers to breastfeeding. Future research should also expand to explore effects of neighborhood level factors. The answers to these questions have implications for all persons attempting to increase breastfeeding rates and reduce racial/ethnic disparities in weight status among mothers and children by aiding the development of interventions that target familial or neighborhood level factors. Additionally, it is important to investigate the effect of changes in laws and policies, health promotion, the Special Supplemental Program for Women, Infants, and Children, and employer support [52].

5. Conclusion

The present study sought out to provide evidence to clarify previous mixed findings on the No associations were found between justed models, nor was there a significant

the racial/ethnic disparities. These results provide evidence in favor of there being no association between breastfeeding and childhood weight status.

Conflicts of interest

All authors declare no conflict of interest.

References

- 1.
 : Issue report: the state of obesity: better policies for a healthier

 America.
 (2018)

 Available
 from:

 https://stateofobesity.org/wp

 content/uploads/2018/09/stateofobesity2018.pdf.
- 2. Levine JA (2011) Poverty and obesity in the U.S. *Diabetes* 60: 2667–2668.

366

- 3. Carey FR, Singh GK, Brown III HS, et al. (2015) Educational outcomes associated with childhood obesity in the United States: Cross-sectional results from the 2011 2012 National *Int J Behav Nutr Phys Act* 12: S3.
- 4. Finkelstein EA, Trogdon JG, Cohen JW, et al. (2009) Annual medical spending attributable to obesity: Payer- and service-specific estimates. *Health Aff (Millwood)*. 28: w822 w 831.
- 5. Harder T, Bergmann R, Kallischnigg G, et al. (2005) Duration of breastfeeding and risk of overweight: A meta-analysis. *Am J Epidemiol* 162: 397–403.
- 6. Yan J, Liu L, Zhu Y, et al. (2014) The association between breastfeeding and childhood obesity: A meta-analysis. *BMC Public Health* 14: 1267–1267.
- 7. Wang L, Collins C, Ratliff M, et al. (2017) Breastfeeding reduces childhood obesity risks. *Child Obes* 13: 197–204.
- 8. Verstraete SG, Heyman MB, Wojcicki JM (2014) Breastfeeding offers protection against obesity in children of recently immigrated Latina women. *J Community Health* 39: 480–486.
- 9. Ehrenthal DB, Wu P, Trabulsi J (2016) Differences in the protective effect of exclusive breastfeeding on child overweight and obesity by mother s race. *Matern Child Health J* 20: 1971 1979.
- 10. Whaley SE, Koleilat M, Leonard S, et al. (2017) Breastfeeding is associated with reduced obesity in Hispanic 2- to 5-year-olds served by WIC. *J Nutr Educ Behav* 49 (7, Supplement 2): S144 S150.e1.
- 11. Casazza K, Fernandez JR, Allison DB (2012) Modest protective effects of breast-feeding on obesity: Is the evidence truly supportive? *Nutrition Today* 47: 33–38.
- 12. Lefebvre CM, John RM (2014) The effect of breastfeeding on childhood overweight and obesity: A systematic review of the literature. *J Am Assoc Nurse Pract* 26: 386–401.
- 13. Cope MB, Allison DB (2008) Critical review of the world health organization s (WHO) 2007 report on evidence of the long-term effects of breastfeeding: Systematic reviews and metaanalysis with respect to obesity. *Obes Rev* 9: 594 605.
- 14. Casazza K, Fontaine KR, Astrup A, et al. (2013) Myths, presumptions, and facts about obesity. *N Engl J Med* 368: 446–454.
- 15. American Academy of Pediatrics (2005) Breastfeeding and the use of human milk. *Pediatrics* 115: 496 507.
- 16. Gibson-Davis CM, Brooks-Gunn J (2006) Couples immigration status and ethnicity as determinants of breastfeeding. *Am J Public Health* 96: 641–646.
- 17. Centers for Disease Control and Prevention: Breastfeeding Report Card. (2018) Available from: https://www.cdc.gov/breastfeeding/data/reportcard.htm.
- 18. Rios-Ellis B, Nguyen-Rodriguez ST, Espinoza L, et al. (2015) Engaging community with Promotores de Salud to support infant nutrition and breastfeeding among Latinas residing in Los Angeles County: Salud Con Hyland s. *Health Care Women Int* 36: 711–729.
- 19.

and support breastfeeding.

- 20. Jones KM, Power ML, Queenan JT, et al. (2015) Racial and ethnic disparities in breastfeeding. *Breastfeed Med* 10: 186–196.
- 21. Baker EH, Rendall MS, Weden MM (2015) Epidemiological paradox or immigrant vulnerability? Obesity among young children of immigrants. *Demography* 52: 1295–1320.

- 22. Wouk K, Lara-Cinisomo S, Stuebe AM, et al. (2016). Clinical interventions to promote breastfeeding by Latinas: A meta-analysis. *Pediatrics* 137: e20152423.
- 23. Bigman G, Wilkinson AV, Pérez A, et al. (2018) Acculturation and breastfeeding among Hispanic American women: A systematic review. *Matern Child Health J* 22: 1260–1277.
- 24. Braveman PA, Egerter SA, Cubbin C, et al. (2004) An approach to studying social disparities in health and health care. *Am J Public Health* 94: 2139–2148.
- 25. California Department of Public Health: California Maternal and Infant Health Assessment (MIHA) Technical Documentation. Available from: http://www.cdph.ca.gov/data/surveys/MIHA/Documents/MIHATechnicalDocument.pdf.
- 26. Cubbin C (2015) Survey Methodology of the geographic research on wellbeing (GROW) study. *BMC Res Notes* 8: 402.
- 27. Heck KE, Braveman P, Cubbin C, et al. (2006) Socioeconomic status and breastfeeding initiation among California mothers. *Public Health Reports* 121: 51–59.
- 28. Akinbami LJ, Ogden CL (2009) Childhood overweight prevalence in the United States: The impact of parent-reported height and weight. *Obesity* 17: 1574–1580.
- 29. Gamliel A, Ziv-Baran T, Siegel RM, et al. (2015) Using weight-for-age percentiles to screen for overweight and obese children and adolescents. *Prev Med* 81: 174–179.
- 30. Reynolds D, Hennessy E, Polek E (2014) Is breastfeeding in infancy predictive of child mental well-being and protective against obesity at 9 years of age? *Child Care Health Dev* 40: 882 890.
- 31. Lindsay AC, Wallington SF, Greaney ML, et al. (2017) and practices related to infant feeding: A qualitative study. *J Hum Lact* 33:595 605.
- 32. Hess C, Ofei A, Mincher A (2015) Breastfeeding and childhood obesity among African Americans: A systematic review. *MCN Am J Matern Child Nurs* 40: 313–319.
- 33. Akbulut-Yuksel M, Kugler AD (2016) Intergenerational persistence of health: Do immigrants get healthier as they remain in the US for more generations? *Econ Hum Biol* 23:136–148.
- 34. Baby-Friendly USA: Celebrating 500 baby-friendly designated facilities in the united states. (2018). Available from: https://www.babyfriendlyusa.org/news/baby-friendly-usa-celebrates-major-milestone-of-500-baby-friendly-designated-facilities-in-the-united-states/.
- 35. Whitaker RC, Wright JA, Pepe MS, et al. (1997) Predicting obesity in young adulthood from childhood and parental obesity. *New England J Med* 337: 869 873.
- 36. Ells LJ, Rees K, Brown T, et al. (2018) Interventions for treating children and adolescents with overweight and obesity: An overview of Cochrane Reviews. *Int J Obes* 42: 1823–1833.
- 37. Balcazar AJ, Grineski SE, Collins TW (2015) The Hispanic health paradox across generations: the relationship of child generational status and citizenship with health outcomes. *Public Health* 129: 691–697.
- 38. Padilla YC, Hamilton ER, Hummer RA (2009) Beyond the epidemiological paradox: The health of Mexican-American children at age 5. *Social Sci Q* 90: 1072–1088.
- 39. De Hoog ML, Kleinman KP, Gillman MW, et al. (2014) Racial/ethnic and immigrant differences in early childhood diet quality. *Public Health Nutr* 17: 1308–1317.
- 40. Mennella JA, Ziegler P, Briefel R, et al. (2006) Feeding infants and toddlers study: The types of foods fed to Hispanic infants and toddlers. *J Am Diet Assoc* 106: 96–106.
- 41. Butte NF, Puyau MR, Adolph AL, et al. (2007) Physical activity in nonoverweight and overweight Hispanic children and adolescents. *Med Sci Sports Exerc* 39: 1257–1266.

- 42. Belcher B, Berrigan D, Dodd K, et al. (2010) Physical activity in US youth: Effect of Race/Ethnicity, age, gender, and weight status. *Med Sci Sports Exerc* 42: 2211 2221.
- 43. Xie B, Gilliland F, Li Y (2003) Effects of ethnicity, family income and education on dietary intake among adolescents. *Pre Med* 36: 30 40.
- 44. Wang M, Heck K, Winkleby M, et al. (2016) Social disparities in dietary habits among women: Geographic research on wellbeing (GROW) study. *Public Health Nutr* 19: 1666–1673.
- 45. Goran MI, Reynolds KD, Lindquist CH (1999) Role of physical activity in the prevention of obesity in children. *Int J Obes* 23: S18.
- 46. Gordon-Larsen P, Harris KM, Ward DS, et al. (2003) Acculturation and overweight-related behaviors among Hispanic immigrants to the US: the National Longitudinal Study of Adolescent Health. *Social Sci Med* 57: 2023 2034.
- 47. Fryar CD, Hirsch R, Eberhardt MS, et al. (2010) Hypertension, high serum total cholesterol, and diabetes: Racial and ethnic prevalence differences in US adults, 1999 2006. NCHS Data Brief, 1 8.
- 48. Narayan KV, Boyle JP, Geiss LS, et al. (2006) Impact of recent increase in incidence on future diabetes burden: US, 2005 2050. *Diabetes Care* 29: 2114 2116.
- 49. Ogden CL, Carroll MD, Lawman HG, et al. (2016) Trends in obesity prevalence among children and adolescents in the United States, 1988 1994 through 2013 2014. *JAMA* 315: 2292 2299.
- 50. U.S. Census Bureau; Census 2010, Summary File 1, Table PCT 11; American Fact Finder. (2019) Available from: http://factfinder.census.gov.
- 51. Stettler N, Zomorrodi A, Posner JC (2007) Predictive value of weight-for-age to identify overweight children. *Obesity* 15: 3106–3112.
- 52. Jacknowitz A (2007) Increasing breastfeeding rates: Do changing demographics explain them? *Women's Health Issues* 17: 84–92.



©2019 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0)