

CORRECTION

Open Access



# Correction: Antibacterial efficacy of peracetic acid in comparison with sodium hypochlorite or chlorhexidine against *Enterococcus faecalis* and *Parvimonas micra*

Benjamín Briseño-Marroquín<sup>1,2</sup>, Angelika Callaway<sup>2</sup>, Natascha Gol Shalamzari<sup>2</sup> and Thomas Gerhard Wolf<sup>1,2\*</sup>

**Correction to:** BMC Oral Health (2022) 22:119

<https://doi.org/10.1186/s12903-022-02148-8>

In this article the wrong figures appeared due to a data-base error as figures 1 and 2; the updated Figs. 1 and 2 should have appeared as shown below.

---

The original article can be found online at <https://doi.org/10.1186/s12903-022-02148-8>.

---

\*Correspondence: [thomas.wolf@zmk.unibe.ch](mailto:thomas.wolf@zmk.unibe.ch)


---

<sup>1</sup> Department of Restorative, Preventive and Pediatric Dentistry, School of Dental Medicine, University of Bern, Freiburgstrasse 7, 3010 Bern, Switzerland


Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



**Fig. 1** Inhibition zone diameters boxplot of different concentrations of peracetic acid (PAA), sodium hypochlorite (NaOCl) and chlorhexidine digluconate (CHX) against *E. faecalis* (n = 6). Medians are shown as lines inside the boxes, 25th and 75th percentiles as boxes, maximum and minimum values as whiskers, and outliers as circle on the plot



**Fig. 2** Inhibition zone diameters boxplot of different concentrations of peracetic acid (PAA), sodium hypochlorite (NaOCl) and chlorhexidine digluconate (CHX) against *P. micra* (n=6). Medians are shown as lines inside the boxes, 25th and 75th percentiles as boxes, maximum and minimum values as whiskers, and outliers as circle on the plot

#### Author details

<sup>1</sup>Department of Restorative, Preventive and Pediatric Dentistry, School of Dental Medicine, University of Bern, Freiburgstrasse 7, 3010 Bern, Switzerland.

<sup>2</sup>Department of Periodontology and Operative Dentistry, University Medical Center of the Johannes Gutenberg-University, Mainz, Germany.

Accepted: 28 April 2022

Published online: 10 May 2022

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.