

RESEARCH

Open Access



Comparative assessment of the Willems dental age estimation methods: a Chinese population-based radiographic study

Jian Wang^{1,4†}, Linfeng Fan^{2†}, Shihui Shen^{1,4}, Meizhi Sui^{1,4}, Jiaxin Zhou^{1,4}, Xiaoyan Yuan^{1,4}, Yiwen Wu^{1,4}, Pingping Zhong^{1,4}, Fang Ji^{3,4*} and Jiang Tao^{1,4*}

Abstract

Background: The comparison of the two Willems dental age estimation methods (gender-specific (Willems I) and non-gender-specific (Willems II)) has not been fully investigated. Here we aimed to explore the applicability of the Willems dental age estimation in an Eastern Chinese population, which may cast light on the field of dental age estimation.

Methods: A total of 1211 oral panoramic radiographs (582 boys and 629 girls) of the Chinese Han population aged 11–16 years old were collected. Dental ages (DAs) were calculated using the Willems method. Statistical significance was set at a p-value < 0.05. Age differences between chronological age (CA) and dental age were analyzed by paired t-tests and mean absolute error (MAE).

Results: The differences between CA and DA determined by the Willems I method were + 0.44 and + 0.09 years for boys and girls, respectively. When using the Willems II method, these differences were + 0.57 and – 0.09. The MAEs of the Willems I method between DA and CA were 0.95 and 1.00 years in boys and girls, respectively. For Willems II, MAEs were 1.02 and 1.00 years in boys and girls.

Conclusions: This study showed that the Willems I method was more accurate than the Willems II method in the boys' group for predicting age from a whole scale. In comparison, Willems II is more competitive in the girls' group. Neither method may be satisfactory for 11-to-16-year-old teenagers in Eastern China.

Keywords: Dental age estimation, Willems method, Chinese population, Tooth development

Introduction

The development characteristics of teeth have been applied in legal, forensic, and clinical fields as an effective weapon to decipher age information [1–3]. In the realm of legal, and forensic science, the decoded age information has cast light on many social affairs, such as welfare

distribution, athlete selection, child enrollment, cadaver identification, and refugee checks [4, 5]. More importantly, age estimation helps juveniles delineate to a large extent for doubted age-related problems [6]. In the community of the clinic, applications have been extended to support routine therapeutic diagnosis and strategy decisions in orthodontics and pediatric dentistry [7].

To date, several radiographic dental age assessment methods for children have been proposed and validated, such as Demirjian's [8], Nolla's [9], Willems's [10], Kvaal's [11], Cameriere's [12], Haavikko's [13], and the London atlas [14], which adopt developing

[†]Jian Wang and Linfeng Fan have contributed equally to this work.

*Correspondence: smilefang98@hotmail.com; doctor_taojiang@126.com

⁴National Center for Stomatology, National Clinical Research Center for Oral Diseases, Shanghai Key Laboratory of Stomatology, Shanghai, China
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

features of teeth reflected on radiographs. Taking Willem's method for instance, it was first proposed by Willem's et al. based on a Belgian-Caucasian population in 2001 [10]. The Willem's method, in other words, a modified Demirjian dental age assessment method, simplified the former method by leaving out a step of data processing and testified its accuracy for dental age assessment in Europe. Inspired by Demirjian and his colleagues, the first Willem's method remained a sex-specific trait, similar to the Demirjian method mentioned. Later, Willem's et al. rethought the old method and constructed a reduced non-gender specific method (Willem's II) to better resolve the problem of possible gender uncertainty under some scenarios [15]. Willem's II method provides a common standard for both genders. It will display good performance in bioarchaeological studies under the circumstances that the sex of skeletal remains is unknown. Although studies on the Willem's I method have been well documented on the planet, over- and underestimation of age have been announced in different parts of the world [1, 5, 9, 16–28]. To the best of our knowledge, the practical effect of the Willem's II method has only been reported in Nemsi's, Hedgel's, and Urzi's works and has not been fully investigated thus far [18, 29, 30]. With respect to Chinese populations, dental age estimations concerning the Willem's I method have been reported over recent years [1, 4, 5, 19, 24, 31]. However, no Willem's II method has been tested in China. Hence, the better choice of the two Willem's methods applied in a population of eastern China is meaningful to explore. This work was designed to compare the applicability and accuracy of the two Willem's dental age methods (Willem's I vs Willem's II) among children aged 11–16 years old in an eastern Chinese population.

Methods and materials

Methods

Retrospective cross-sectional research was conducted in a hospital in Shanghai, eastern China. The Independent Ethics Committee of the Shanghai Ninth People's Hospital, affiliated with Shanghai Jiao Tong University School of Medicine (2017–282-T212) approved and authorized the project.

All the included orthopantomograms (OPGs) were randomly selected from the database of the hospital and followed by strict inclusion standards, which have been elucidated in previous work [7].

The patients underwent an OPG check before any oral treatments. All 1211 samples were from a Chinese Han population aged 11–16 years old.

Data analysis

A total of 1211 OPGs were qualified for the current work. The information of each age group at an interval of one year is displayed in Table 1.

CA was calculated by the date of the OPG taken preoperatively minus the date of birth, which was expressed by two decimal points. All X-ray images were evaluated by the Willem's method (Willem's I and Willem's II). To acquire the actual DA, we have to evaluate all the 7 left mandibular teeth (with third molars excluded)'s developmental stages (from A to H) judging from the status of the initial crown's formation to the terminal apex's closure. Then, we marked all the 7 teeth's scores and summed them up to obtain the ideal DA value with the provided dataset tables.

Cohen's kappa tests were employed to make the inner- and inter-agreements' tests [32]. Intra- and inter-agreements were calculated to give better quantitative values of agreements during the studies with the repeated data. The analysis included descriptive and inferential statistics. Descriptive statistics such as the mean differences and standard deviation (SD) were calculated. Differences between DA and CA were generated by subtracting DA from CA, i.e., (DA-CA). Then, differences were stratified based on age and sex and analyzed using the paired *t-test*. The mean absolute error (MAE) was used to assess the precision of the two Willem's methods. All analyses were conducted in SPSS 17.0 for Windows (SPSS, Inc., Chicago, IL). A *p*-value less than 0.05 was considered to be statistically significant.

Results


A total of 1211 OPTs of sub-adults aged from 11 to 16 years were eligible for the present study, including 582 boys and 629 girls. The samples were classified into 6 subgroups at an interval of one year. Details concerning samples were listed in Table 1.

Table 1 Sample distribution by gender and age

Age group	Boys	Girls	Total
11	73	105	178
12	112	97	209
13	109	99	208
14	106	105	211
15	96	110	206
16	86	113	199
Total	582	629	1211

Table 2 Accuracy comparison of Willems I method of dental age estimation

Gender	Age group	CA (years) (mean \pm SD)	DA (years) (mean \pm SD)	CA-DA (years) (mean \pm SD)	p	MAE
Boys	11	11.47 \pm 0.28	11.52 \pm 0.91	(-)0.05 \pm 0.98	0.665	0.78
	12	12.50 \pm 0.28	12.05 \pm 0.83	0.45 \pm 0.84	0	0.75
	13	13.46 \pm 0.28	13.05 \pm 1.18	0.40 \pm 1.20	0.001	1.03
	14	14.42 \pm 0.31	14.19 \pm 1.60	0.23 \pm 1.59	0.147	1.12
	15	15.48 \pm 0.28	14.85 \pm 1.15	0.63 \pm 1.14	0	1.05
	16	16.48 \pm 0.24	15.53 \pm 0.73	0.95 \pm 0.77	0	0.95
	Total	13.98 \pm 1.63	13.54 \pm 1.79	0.44 \pm 1.17	0	0.95
Girls	11	11.45 \pm 1.16	12.11 \pm 1.64	(-)0.66 \pm 1.34	0	1.03
	12	12.46 \pm 0.29	13.03 \pm 1.04	(-)0.58 \pm 1.05	0	0.86
	13	13.49 \pm 0.26	13.62 \pm 1.26	(-)0.13 \pm 1.24	0.289	1.02
	14	14.47 \pm 0.32	14.09 \pm 1.75	0.39 \pm 1.72	0.025	1.25
	15	15.43 \pm 0.27	15.12 \pm 1.04	0.31 \pm 1.04	0	0.76
	16	16.46 \pm 0.29	15.38 \pm 0.82	1.08 \pm 0.80	0	1.08
	Total	14.03 \pm 1.81	13.93 \pm 1.73	0.09 \pm 1.37	0.091	1




Inter and intra-agreements

The Cohen's kappa values reached 0.78 and 0.79 for boys and girls, respectively, which displayed good inter- and intra-agreement in the experiments.

Willems I method

Using the Willems I dataset, the mean dental age for boys and girls was 13.54 and 13.93 years, respectively (Tables 2 and 4). The general mean difference (CA-DA) for boys and girls was 0.44 ± 1.17 ($p < 0.05$) and 0.09 ± 1.37 ($p > 0.05$), respectively. An underestimation (CA-DA) of age was observed in all the subgroups except for boys in the 11–12-year group and girls in the 11–12, 12–13, and 13–14-year groups. The mean absolute error was 0.95 and 1.00 years for boys and girls, respectively (seen in Figs. 1 and 2).



Willems II method

Adopting the Willems II method, the mean dental age for boys and girls was 13.41 and 14.11 years, respectively (Tables 3 and 4). The overall mean difference (CA-DA) for boys and girls was 0.57 ± 1.20 ($p < 0.05$) and $(-)0.09 \pm 1.34$ ($p > 0.05$), respectively. As can be reflected in Figs. 1 and 2, an underestimation (CA-DA) of age was observed in all the subgroups except for the girls in the 11–12, 12–13, and 13–14 years groups. The mean absolute error was 1.02 and 1.00 years for boys and girls, respectively (seen in Figs. 1 and 2).

Discussion

The Willems method has gained popularity to a great extent as an easy-to-check approach for estimating the actual age. The Willems I method's over/underestimation of age has been well reported worldwide. overestimation of chronological age was validated in a variety of counties or regions such as India [33], Turkey [34],

Table 3 Accuracy comparison of Willems II method of dental age estimation

Gender	Age group	CA (years) (mean \pm SD)	DA (years) (mean \pm SD)	CA-DA (years) (mean \pm SD)	p	MAE
Boys	11	11.47 \pm 0.28	11.4 \pm 0.91	0.07 \pm 0.98	0.556	0.8
	12	12.50 \pm 0.28	11.99 \pm 0.84	0.5 \pm 0.86	0	0.78
	13	13.46 \pm 0.28	12.94 \pm 1.20	0.52 \pm 1.21	0	1.07
	14	14.42 \pm 0.31	14.01 \pm 1.66	0.41 \pm 1.64	0.012	1.24
	15	15.48 \pm 0.28	14.64 \pm 1.24	0.84 \pm 1.21	0	1.14
	16	16.48 \pm 0.24	15.41 \pm 0.77	1.06 \pm 0.81	0	1.06
	Total	13.98 \pm 1.63	13.41 \pm 1.78	0.57 \pm 1.20	0	1.02
Girls	11	11.45 \pm 1.16	12.35 \pm 1.58	(-)0.91 \pm 1.30	0	1.01
	12	12.46 \pm 0.29	13.25 \pm 0.99	(-)0.79 \pm 1.00	0	0.94
	13	13.49 \pm 0.26	13.81 \pm 1.21	(-)0.32 \pm 1.19	0.008	0.75
	14	14.47 \pm 0.32	14.26 \pm 1.68	0.21 \pm 1.65	0.205	1.17
	15	15.43 \pm 0.27	15.25 \pm 0.99	0.18 \pm 1.00	0.061	0.75
	16	16.46 \pm 0.29	15.50 \pm 0.79	0.95 \pm 0.77	0	0.95
	Total	14.03 \pm 1.81	14.11 \pm 1.66	(-)0.09 \pm 1.34	0.106	1

Table 4 Comparison of accuracy of Willems I and II methods

Gender	N	Willems I		Willems II		Willems I vs II	
		CA-DA (mean \pm sd)	MAE	CA-DA (mean \pm sd)	MAE	I-II (mean \pm sd)	p-value
Boys	582	0.44 \pm 1.17	0.95	0.57 \pm 1.20	1.02	(-)0.13 \pm 0.23	0
Girls	629	0.09 \pm 1.37	1.00	(-)0.09 \pm 1.34	1	0.18 \pm 0.08	0
Total	1211	0.26 \pm 1.28	0.98	0.23 \pm 1.32	1.01	0.03 \pm 0.23	0

Thailand [35], Spain [36], Kenya [37] and Poland [38]. Elsewhere, there were also reports of underestimation in north China [24], Tunisia [30], and Sri Lanka [39]. In allusion to the differences of the Willems I method among several regions, a systemic review, and meta-analysis we conducted previously concerning the Willems I method revealed the Willems method overestimated CA by 0.18 years and 0.06 years for boys and girls, respectively [4]. It also revealed that ethnicity specificity was necessary when adopting the Willems I method. This conclusion was consistent with other similar meta-analyses [25, 26, 40]. In the current work, the underestimation of age was $+0.44 \pm 1.17$ ($p < 0.05$) and $+0.09 \pm 1.37$ ($p = 0.091$) for the boy and girl groups, respectively. From the whole perspective, the girls' group showed a more accurate effect of age assessment when compared with the boys' group. This phenomenon was also viewed in our previous works and other printed studies [5, 41]. In subgroups' accuracy analysis, the Willems I method was acceptable in 11–12, 12–13, 13–14, and 14–15 years group of boys, while in 13–14, 14–15 and 15–16 years groups of girls, relatively compatible intervals were observed for age assessment.

In the journey of investigating the Willems II method's applicability, few works have been reported over the last decades (Table 4). Urzel et al. observed a mean difference between CA and DA of 0.03 and 0.00 years for both genders with the Willems I and II method, respectively [15]. They concluded that both methods were appropriate when applied to a French population, although the Willems I method was more accurate according to their data. Another similar study from an Indian population with the Willems II method revealed that the mean values were 0.06 ± 0.80 , -0.11 ± 0.79 , and -0.01 ± 0.80 years for boys, girls, and the total sample, respectively [33]. Among Tunisian sub-adults [34], the Willems II method tended to underestimate age by 0.91, and 0.64 years for boys and girls, respectively. Whereas the Willems I method indicated an underestimation of chronological age by 0.40 years for boys, and by 0.69 years for girls. In their investigation, Willems I was more accurate when compared with Willems II in the Tunisian population. In the current study, the mean values of CA-DA were 0.57 ± 1.20 ($p < 0.05$) and -0.09 ± 1.34 ($p = 0.106$) years for boys and girls, respectively. A more accurate estimation was observed in the girls' group in both methods. These results revealed that Willems I was more accurate

and reliable for boys, while Willems II was a better option for girls. Our results are in accordance with the aforementioned studies. To be more specific, Willems I or II was an optional choice for ascertaining age despite Willems I having a slightly higher accuracy for 11–12, 12–13, 14 and 14–15 years groups among boys according to our data. In girls' subgroups, the data appeared to confirm that Willems II lost its advantages in younger age groups of 11–12 and 12–13 years, and regained its leading positions in 13–14 and 14–15 years groups. The results showed that neither method was a reliable choice for evaluating 16–17 years old for both genders. From our perspective, when an individual grows 16 years or older, the apex's closure has been completed in most cases, we could not tell the more differences from the radiograph via Willems method. The inner drawbacks posed restrictions on individuals of age large than 16 years old for assessing dental age via the Willems method.

Despite the accuracy assessment of Willems I and Willems II, the study still has limitations that must be elucidated. The samples we collected ranged only from 11 to 16 years old, we did not retrieve lower age groups younger than 11 years old. In our initial design of the project, we intended to take a clear and accurate evaluation of some important age thresholds like 14 and 16 years old. Because these thresholds provide key information to verdict a teenager's delinquency in China. We have to admit that if we could have recruited individuals less than 11 years old, our research would have a deep and comprehensive understanding of the two Willems methods. Despite we have performed the dental age evaluation among 11–16-year-old sub-adults, more age subgroups will be recruited and show up in our subsequent works.

Conclusions

In conclusion, the present study investigated the applicability of the Willems I and Willems II methods with a sample of Eastern Chinese teenagers aged 11–16 years old. These outcomes suggest that Willems I is more accurate for boys, while the Willems II method prefers girls; Neither method may be a perfect match for the Eastern Chinese Han population despite some subgroups having clear and accurate age assessment. An ethnicity-specific model based on the Willems method or further modifications is encouraged to prosper the science of dental age estimation.

Abbreviations

DA: Dental age; CA: Chronological age; MAE: Mean absolute error; SD: Standard deviation; N: Number; OPG: Orthopantomogram.

Acknowledgements

Not applicable.

Author contributions

JW: Conceptualization, Methodology, Writing—Original Draft, Funding. LFF: Resources. XYY, YWW, PPZ: Software, Investigation, Data Curation. SHS, MZS, JXZ: Formal Analysis, Validation—Investigation. FJ: Supervision, Project Advice. JT: Writing—Review and Editing, Supervision, Funding. All authors read and approved the final manuscript.

Funding

This work was sponsored by Shanghai Sailing Program (21YF1424100), the Interdisciplinary Program of Shanghai Jiao Tong University (YG2019ZDA07), and the Innovation research team of high-level local universities in Shanghai (SHSMU-ZLCX20212401).

Availability of data and materials

The datasets used for the present study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The Independent Ethics Committee of the Shanghai Ninth People's Hospital, affiliated with Shanghai Jiao Tong University School of Medicine (2017-282-T212) approved and authorized the project. All methods were carried out following relevant guidelines and regulations. We declare that Informed consent was obtained from all subjects and/or their legal guardian(s) for study participation.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of General Dentistry, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, College of Stomatology, Shanghai Jiao Tong University, Shanghai, China. ²Department of Radiology, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, College of Stomatology, Shanghai Jiao Tong University, Shanghai, China. ³Department of Orthodontics, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, College of Stomatology, Shanghai Jiao Tong University, Shanghai, China. ⁴National Center for Stomatology, National Clinical Research Center for Oral Diseases, Shanghai Key Laboratory of Stomatology, Shanghai, China.

Received: 20 May 2022 Accepted: 23 August 2022

Published online: 03 September 2022

References

1. Han MQ, Jia SX, Wang CX, Chu G, Chen T, Zhou H, Guo YC. Accuracy of the Demirjian, Willems and Nolla methods for dental age estimation in a northern Chinese population. *Arch Oral Biol*. 2020;118: 104875.
2. Machado MA, Daruge Júnior E, Fernandes MM, Lima IFP, Cericato GO, Franco A, Paranhos LR. Effectiveness of three age estimation methods based on dental and skeletal development in a sample of young Brazilians. *Arch Oral Biol*. 2018;85:166–71.
3. Flood SJ, Franklin D, Turlach BA, McGeachie J. A comparison of Demirjian's four dental development methods for forensic age estimation in South Australian sub-adults. *J Forensic Leg Med*. 2013;20(7):875–83.
4. Wang J, Ji F, Zhai Y, Park H, Tao J. Is Willems method universal for age estimation: a systematic review and meta-analysis. *J Forensic Leg Med*. 2017;52:130–6.
5. Wang J, Bai X, Wang M, Zhou Z, Bian X, Qiu C, Li C, Yang Z, Chen G, Ji F, et al. Applicability and accuracy of Demirjian and Willems methods in a population of Eastern Chinese subadults. *Forensic Sci Int*. 2018;292:90–6.
6. Melo M, Ata-Ali J. Accuracy of the estimation of dental age in comparison with chronological age in a Spanish sample of 2641 living subjects using

- the Demirjian and Nolla methods. *Forensic Sci Int.* 2017;270:276 e271–276 e277.
7. Wang J, Wang M, Shen S, Guo Y, Fan L, Ji F, Tao J. Testing the nonlinear equations for dental age evaluation in a population of eastern China. *Leg Med (Tokyo).* 2021;48: 101793.
 8. Demirjian A, Goldstein H, Tanner JM. A new system of dental age assessment. *Hum Biol.* 1973;45(2):211–27.
 9. Maber M, Liversidge HM, Hector MP. Accuracy of age estimation of radiographic methods using developing teeth. *Forensic Sci Int.* 2006;159(Suppl 1):S68–73.
 10. Willems G, Van Olmen A, Spiessens B, Carels C. Dental age estimation in Belgian children: Demirjian's technique revisited. *J Forensic Sci.* 2001;46(4):893–5.
 11. Kvaal SI, Kolltveit KM, Thomsen IO, Solheim T. Age estimation of adults from dental radiographs. *Forensic Sci Int.* 1995;74(3):175–85.
 12. Cameriere R, Pacifici A, Pacifici L, Polimeni A, Federici F, Cingolani M, Ferrante L. Age estimation in children by measurement of open apices in teeth with Bayesian calibration approach. *Forensic Sci Int.* 2016;258:50–4.
 13. Galić I, Vodačić M, Cameriere R, Nakaš E, Galić E, Selimović E, Brkić H. Accuracy of Cameriere, Haavikko, and Willems radiographic methods on age estimation on Bosnian-Herzegovian children age groups 6–13. *Int J Legal Med.* 2011;125(2):315–21.
 14. AlQahtani SJ, Hector MP, Liversidge HM. Brief communication: the London atlas of human tooth development and eruption. *Am J Phys Anthropol.* 2010;142(3):481–90.
 15. Willems G, Thevissen PW, Belmans A, Liversidge HM, Willems II. Non-gender-specific dental maturity scores. *Forensic Sci Int.* 2010;201(1–3):84–5.
 16. Yang F, Jacobs R, Willems G. Dental age estimation through volume matching of teeth imaged by cone-beam CT. *Forensic Sci Int.* 2006;159(Suppl 1):S78–83.
 17. Djukic K, Zelic K, Milenkovic P, Nedeljkovic N, Djuric M. Dental age assessment validity of radiographic methods on Serbian children population. *Forensic Sci Int.* 2013;231(1–3):398 e391–395.
 18. Urzel V, Bruzek J. Dental age assessment in children: a comparison of four methods in a recent French population. *J Forensic Sci.* 2013;58(5):1341–7.
 19. Ye X, Jiang F, Sheng X, Huang H, Shen X. Dental age assessment in 7–14-year-old Chinese children: comparison of Demirjian and Willems methods. *Forensic Sci Int.* 2014;244:36–41.
 20. Patnana AK, Vabbalareddy RS, NR VV. Evaluating the reliability of three different dental age estimation methods in visakhapatnam children. *Int J Clin Pediatr Dent.* 2014;7(3):186–91.
 21. Gupta S, Mehendiratta M, Rehani S, Kumra M, Nagpal R, Gupta R. Age estimation in Indian children and adolescents in the NCR region of Haryana: a comparative study. *J Forensic Dent Sci.* 2015;7(3):253–8.
 22. Mohammed RB, Sanghvi P, Perumalla KK, Srinivasaraju D, Srinivas J, Kalyan US, Rasool SM. Accuracy of four dental age estimation methods in southern Indian children. *J Clin Diagn Res.* 2015;9(1):HC01–08.
 23. Kumaresan R, Cugati N, Chandrasekaran B, Karthikeyan P. Reliability and validity of five radiographic dental-age estimation methods in a population of Malaysian children. *J Investig Clin Dent.* 2016;7(1):102–9.
 24. Zhai Y, Park H, Han J, Wang H, Ji F, Tao J. Dental age assessment in a northern Chinese population. *J Forensic Leg Med.* 2016;38:43–9.
 25. Mohd Yusof MYP, Wan Mokhtar I, Rajasekharan S, Overholser R, Martens L. Performance of Willem's dental age estimation method in children: a systematic review and meta-analysis. *Forensic Sci Int.* 2017;280:245 e241–245 e210.
 26. Sehwat JS, Singh M. Willems method of dental age estimation in children: a systematic review and meta-analysis. *J Forensic Leg Med.* 2017;52:122–9.
 27. Esan TA, Yengopal V, Schepartz LA. The Demirjian versus the Willems method for dental age estimation in different populations: a meta-analysis of published studies. *PLoS ONE.* 2017;12(11): e0186682.
 28. Van Dyck J, de Llano-Perula MC, Willems G, Verdonck A. Dental development in cleft lip and palate patients: a systematic review. *Forensic Sci Int.* 2019;300:63–74.
 29. Hegde S, Patodia A, Dixit U. Willems I VS Willems II: a comparative study of accuracy in 5–15 year old Indian children. *Forensic Sci Int.* 2016;266:117–22.
 30. Nemsí H, Ben Daya M, Salem NH, Masmoudi F, Bouanène I, Maatouk F, Aissaoui A, Chadly A. Applicability of Willems methods and Demirjian's four teeth method for dental age estimation: cross sectional study on Tunisian sub-adults. *Forensic Sci Int.* 2018;291:281.e281–281.e289.
 31. Yang Z, Geng K, Liu Y, Sun S, Wen D, Xiao J, Zheng Y, Cai J, Zha L, Liu Y. Accuracy of the Demirjian and Willems methods of dental age estimation for children from central southern China. *Int J Legal Med.* 2019;133(2):593–601.
 32. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics.* 1977;33(1):159–74.
 33. Balla SB, Venkat Baghirath P, Hari Vinay B, Vijay Kumar J, Babu DBG. Accuracy of methods of age estimation in predicting dental age of preadolescents in South Indian children. *J Forensic Leg Med.* 2016;43:21–5.
 34. Akkaya N, Yilanci H, Göksülük D. Applicability of Demirjian's four methods and Willems method for age estimation in a sample of Turkish children. *Leg Med (Tokyo).* 2015;17(5):355–9.
 35. Duangto P, Janhom A, Prasitwattanaseree S, Mahakkanukrauh P, Iamroon A. New prediction models for dental age estimation in Thai children and adolescents. *Forensic Sci Int.* 2016;266:583.e581–583.e585.
 36. Paz Cortés MM, Rojo R, Alía García E, Mourelle Martínez MR. Accuracy assessment of dental age estimation with the Willems, Demirjian and Nolla methods in Spanish children: Comparative cross-sectional study. *BMC Pediatr.* 2020;20(1):361.
 37. Kihara EN, Gichangi P, Liversidge HM, Butt F, Gikenye G. Dental age estimation in a group of Kenyan children using Willems' method: a radiographic study. *Ann Hum Biol.* 2017;44(7):614–21.
 38. Sobieska E, Fester A, Nieborak M, Zadurska M. Assessment of the dental age of children in the polish population with comparison of the Demirjian and the Willems methods. *Med Sci Monit.* 2018;24:8315–21.
 39. Ranasinghe S, Perera J, Taylor JA, Tennakoon A, Pallewatte A, Jayasinghe R. Dental age estimation using radiographs: towards the best method for Sri Lankan children. *Forensic Sci Int.* 2019;298:64–70.
 40. Jayaraman J, Wong HM, King NM, Roberts GJ. The French-Canadian data set of Demirjian for dental age estimation: a systematic review and meta-analysis. *J Forensic Leg Med.* 2013;20(5):373–81.
 41. Jayaraman J, Roberts GJ, Wong HM, King NM. Dental age estimation in southern Chinese population using panoramic radiographs: validation of three population specific reference datasets. *BMC Med Imaging.* 2018;18(1):5.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

