

RESEARCH

Open Access



Association of lipid profile and reported edentulism in the elder population: data from the China Health and Retirement Longitudinal Study

Shuping Wang¹, Yutao Wang², Riyue Yu¹, Dingxiang Yuan¹, Yaofeng Ni¹, Lixin Wang¹, Man Sun³ and Xin Wang^{4*}

Abstract

Objectives: Relationship between lipid profile and periodontitis has been reported. However, the association between lipid parameters and edentulism is unclear. This study aimed to investigate the association between lipid profile and reported edentulism in the elder population using a national cohort.

Methods: A total of 3 100 participants aged 65 or above were enrolled in 2011 from China Health and Retirement Longitudinal Study, which was a national population-based survey. We used adjusted logistics models to investigate the relationship between lipid profile and reported edentulism before and after propensity score matching.

Results: The mean (SD) age was 71.96 (5.63) years, and 1 581 (51.0%) were men. There were 254 (8.2%) individuals reporting edentulism, and the low-density lipoprotein cholesterol (LDL-C) was significantly higher in the reported edentulism group, compared with the non-edentulism (122.48 vs. 116.91 mg/dl, $P = 0.015$). In the multivariable model, LDL-C was significantly associated with a higher odds of reported edentulism (adjusted OR [95% CI], 1.004 [1.001–1.008]). In the matched population, LDL-C, non high-density lipoprotein cholesterol, remnant cholesterol, total cholesterol and triglycerides were positively associated with reported edentulism, while HDL-C was negatively associated.

Conclusions: Lipid profiles are probably associated with edentulism, indicating the interaction between oral health and metabolic status in the elder population.

Keywords: Lipid profile, Reported edentulism, LDL cholesterol, Elder population

Introduction

Oral diseases are among the most prevalent diseases worldwide and have serious health, mental and economic burdens [1]. Dental caries and periodontal disease are two major oral diseases impacting the quality of life globally [2]. Edentulism and tooth loss are severe

consequences of these diseases and occur throughout life, especially in the elder population. Although the epidemiology studies show a gradual decline in the prevalence and incidence of edentulism and tooth loss at the global, regional, and country levels [3], the prevalence still increases with aging showing an incidence peak at 65 years [4].

Edentulism and tooth loss are closely related with obesity [5], diabetes [6], dementia [7, 8], chronic obstructive pulmonary disease [9], nonalcoholic fatty liver disease [10] and cancer [11, 12]. Dyslipidaemias are alterations in

*Correspondence: angus.wangxin@outlook.com

⁴ Department of VIP Dental Service, Beijing Stomatological Hospital, Capital Medical University, Beijing, China

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

the lipid profiles which are independently associated with cardiovascular diseases, particularly elevated plasma low-density lipoprotein cholesterol (LDL-C) [13]. The global burden of dyslipidaemias is gradually increasing, and LDL-C level has ranked the 8th leading risk factor of mortality in 2019, which ranked 15th in 1990 [14]. There is emerging evidence regarding the interaction between lipid metabolism and systemic, local inflammation and inflammation related diseases [15–17]. Edentulism and tooth loss, as polymicrobial chronic inflammatory diseases, reflect sustained local inflammatory milieu [18], which play a potential role in the lipid metabolism and lipid level. A recent study reported the association between oral health conditions and changes in lipid profile among people aged 40 years or above [19]. However, the association of edentulism and lipid levels remains unclear in the elder population with the highest risk of tooth loss and edentulism.

Triglycerides and cholesterol perform distinct functions in the body, and are important markers for health. Triglycerides are a most common type of fat or lipid in the blood storing excess energy. Cholesterol is a waxy substance embedded in the lipoprotein, generally classified into high-density lipoprotein cholesterol (HDL-C), LDL-C and remnant cholesterol according to the density. In this study, we aimed to investigate the association of reported edentulism with lipid parameters in the elder population using a national cohort.

Materials and methods

Data source

Participants of the China Health and Retirement Longitudinal Study (CHARLS) at baseline visit between were included in this study. CHARLS is a public open research cohort database collecting a wide range of social-economic data and personal health information (<http://charls.pku.edu.cn/>). CHARLS covers a nationally representative sample of Chinese residents ages 45 and older to promote research on the elderly. The baseline national wave of was initiated in 2011 and included about 10 000 households and 17 500 individuals in 150 counties/districts and 450 villages/resident committees using multi-stage stratified probability proportionate to size (PPS) sampling.

This study was in accordance with the principles of the Declaration of Helsinki and was approved by the Ethics Committee of Capital Medical University (grant number: 2020SY031). All participants provided their written informed consents before participating in the current study. The following criteria were required: (1) taking blood lipid examinations; (2) reporting oral health and tooth status; (3) without history of any malignancy; (4) aged 65 years or above. Figure 1 shows the design and

flowchart of this current study. Finally, there were 3 100 individuals enrolled in the final analysis.


Measurements and definitions

The demographic characteristics, lifestyles, and health status were acquired using the standard questionnaire, including age, sex, education level, marital status, smoking status, drinking status and self-reported health conditions. Educational level was categorized as 'primary school or below', 'middle school', and 'high school or above'. Marital status included 'married', 'unmarried' and 'others'. Smoking status was defined as 'never smoking', 'current smoker', and 'former smoker'. Drinking status was defined as 'current drinking more than once per month', 'current drinking once or less than once per month', and 'no current drinking'. Self-reported health conditions included tooth loss status and the diagnosis of hypertension, diabetes and dyslipidemia. Body mass index (BMI) was calculated as weight (kg)/[height (m)*height (m)]. The definition of obesity was BMI ≥ 28.0 kg/m² for the Asian population [20]. Blood laboratory tests were carried out using fasting venous blood samples, and the lipid profiles included triglycerides, total cholesterol, HDL-C and LDL-C. The nonHDL-C was calculated as total cholesterol minus HDL-C. The remnant cholesterol was calculated using the Martin-Hopkins equation using the Stata.do-file (<https://www.ldlcalculator.com/>). The reported edentulism was assessed by the following question: "Have you lost all of your teeth? (Yes / no)".

Statistical analysis

The characteristics of the population were stratified by reported edentulism or not. The basic characteristics were presented as the mean (standard deviation, SD) for continuous variables and number (proportion) for categorical variables. The differences were compared by student's t-test test for continuous variables and chi-square test for categorical variables. The distributions of lipid profiles were presented using box-plots.

The unadjusted and adjusted logistics regression models were used to analyze the association of the lipid parameters with reported edentulism. To control the potential confounding factors, age, sex, education level, marriage status, smoking status, drinking, BMI level, hypertension or not, dyslipidemia or not, diabetes or not were adjusted in multivariable model. In addition, we used propensity score method to match controls for individuals of reported edentulism. The matching ratio was 1:2, and all the potential confounding factors mentioned above were considered in the matching process. The association between lipid parameters and reported edentulism was further explored in the matched population.



All the analyses above were analyzed using R software (version 4.1.0). The differences were considered statistically significant at two-side $P < 0.05$.

Results

Population characteristics

The mean (SD) age of the whole population was 71.96 (5.63) years, and 1 581 (51.0%) were men. There were 254 (8.2%) individuals with reported edentulism. The individuals of reported edentulism were older than people of non-edentulism (72.63 vs. 71.91, $P = 0.049$). There were no significant differences for sex, education, marriage, BMI, smoking, drinking and self-reported diagnosis of hypertension, diabetes and dyslipidemia as shown in Table 1. LDL-C was significantly higher in people with reported edentulism (122.48 vs. 116.91 mg/dl, $P = 0.015$), and no significant differences were observed for HDL-C, nonHDL-C, remnant cholesterol, total cholesterol and triglycerides ($P > 0.05$). In the matched population, 208 individuals with reported edentulism and 416 controls were selected, and the covariates were balanced as shown in Table 2. Of note, LDL-C (122.53 vs. 115.00 mg/dl, $P = 0.009$), nonHDL-C (145.14 vs. 134.02 mg/dl, $P < 0.001$), remnant cholesterol (22.40 vs. 20.16 mg/dl, $P = 0.004$), total cholesterol (197.76 vs. 190.47 mg/dl, $P = 0.022$) and triglycerides (116.30 vs. 101.64 mg/dl,

$P = 0.017$) were significantly higher in the reported edentulism group, while HDL-C (52.62 vs. 56.45 mg/dl, $P = 0.005$) was significantly lower. The distributions of lipid profiles between the reported edentulism and non-edentulism groups in the whole population and matched population were shown in Fig. 2.

Association of lipid profiles and reported edentulism

In the whole population, LDL-C was significantly associated with a higher odds of reported edentulism, and the adjusted OR (95% CI) was 1.004 (1.001–1.008). We did not observe independent association between reported edentulism and other lipid parameters as shown in Table 3. In the matched population, LDL-C, HDL-C, nonHDL-C, remnant cholesterol, total cholesterol and triglycerides were all significantly associated with reported edentulism as shown in Fig. 3, and the adjusted OR (95% CI) values were 1.007 (1.002–1.012), 0.985 (0.974–0.995), 1.008 (1.004–1.013), 1.027 (1.008–1.046), 1.005 (1.001–1.010) and 1.003 (1.001–1.005) respectively.

Discussion

In this study, we found that reported edentulism was independently associated with lipid metabolism in the elder population, particularly LDL-C. In the matched population, six lipid parameters were significantly

Table 1 Characteristics of this study

Characteristics	Non reported edentulism	Reported edentulism	P value
Participants, No	2846	254	
Age, years	71.91(5.58)	72.63(6.12)	0.049
Men, n (%)	1463(51.4)	118(46.5)	0.148
Education, n (%)			
Primary or below	2381(83.8)	223(88.1)	0.195
Middle school	282(9.9)	19(7.5)	
High school or above	177(6.2)	11(4.3)	
Marriage status, n (%)			
Married	2121(74.5)	181(71.3)	0.316
Unmarried	25(0.9)	1(0.4)	
Others	700(24.6)	72(28.3)	
Smoking status, n (%)			
Never	1618(56.9)	149(58.7)	0.831
Current	872(30.7)	76(29.9)	
Former	354(12.4)	29(11.4)	
Drinking status, n (%)			
> once a month	664(23.3)	53(20.9)	0.503
≤ once a month	173(6.1)	13(5.1)	
No drinking	2007(70.6)	188(74.0)	
Hypertension, n (%)	959(33.8)	77(30.3)	0.286
Dyslipidemia, n (%)	271(9.7)	18(7.2)	0.245
Diabetes, n (%)	182(6.4)	13(5.1)	0.491
BMI, kg/m ²	23.42(3.99)	22.97(3.34)	0.106
Obesity, n (%)	270(11.1)	17(7.9)	0.181
Total cholesterol, mg/dL	193.30(38.34)	197.44(40.65)	0.101
LDL-C, mg/dL	116.91(34.52)	122.48(37.45)	0.015
HDL-C, mg/dL	52.02(15.91)	52.20(15.78)	0.865
nonHDL-C, mg/dL	141.27(38.15)	145.24(40.77)	0.115
Remnant cholesterol, mg/dL	23.34(11.62)	22.62(9.01)	0.338
Triglycerides, mg/dL	125.96(90.20)	118.57(70.62)	0.204

Data are presented as mean (SD), or number (proportion)

BMI, body mass index; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol

associated with reported edentulism including LDL-C, HDL-C, nonHDL-C, remnant cholesterol, total cholesterol and triglycerides.

The relationships between periodontitis and blood lipid parameters have been reported in different populations and regions. Previous studies showed that periodontitis was associated with low HDL-C, high triglycerides and total cholesterol [21–24]. On the contrary, some studies found no significant associations between periodontitis and lipid profiles [25, 26]. On the other hand, Song et al. found that periodontitis was both associated with baseline lipid profile and the longitudinal changes of blood lipid parameters [19]. The age levels of different populations could partially contribute to the inconsistent results. Kim et al. reported that the number of tooth loss

was positively correlated with triglycerides level [27]. Our study provided further information on the association between lipid parameters and reported edentulism in the elder population.

The local and systematic inflammation may partially explain the underlying mechanism between edentulism and blood lipid parameters. Tooth loss causes the surrounding tissues damages and leads to chronic local inflammation [28–30]. Through the local tissue injury and chronic inflammation, oral bacteria could harm blood vessels or cause systemic inflammatory response [31–33]. In addition, control of the oral local infection is associated with a reduction in serum inflammatory markers including C-reactive protein (CRP) and interleukin-6 (IL-6) [34, 35]. These inflammatory reactions

Table 2 Characteristics of the matched population

Characteristics	Non reported edentulism	Reported edentulism	P value
Participants, No	416	208	
Age, years	72.08(5.59)	72.66(6.09)	0.235
Men, n (%)	165(39.7)	95(45.7)	0.177
Education, n (%)			
Primary or below	380(91.3)	183(88.0)	0.408
Middle school	24(5.8)	17(8.2)	
High school or above	12(2.9)	8(3.8)	
Marriage status, n (%)			
Married	290(69.7)	149(71.6)	0.751
Unmarried	4(1.0)	1(0.5)	
Others	122(29.3)	58(27.9)	
Smoking status, n (%)			
Never	256(61.5)	118(56.7)	0.51
Current	111(26.7)	63(30.3)	
Former	49(11.8)	27(13.0)	
Drinking status, n (%)			
> once a month	85(20.4)	45(21.6)	0.854
≤ once a month	19(4.6)	11(5.3)	
No drinking	312(75.0)	152(73.1)	
Hypertension, n (%)	120(28.8)	59(28.4)	0.975
Dyslipidemia, n (%)	19(4.6)	13(6.2)	0.48
Diabetes, n (%)	19(4.6)	12(5.8)	0.648
BMI, kg/m ²	23.29(3.59)	23.04(3.34)	0.394
Obesity, n (%)	37(8.9)	17(8.2)	0.88
Total cholesterol, mg/dL	190.47(35.66)	197.76(40.93)	0.022
LDL cholesterol, mg/dL	115.00(31.45)	122.53(37.69)	0.009
HDL cholesterol, mg/dL	56.45(15.90)	52.62(16.06)	0.005
nonHDL cholesterol, mg/dL	134.02(34.90)	145.14(40.83)	< 0.001
Remnant cholesterol, mg/dL	20.16(8.86)	22.40(9.36)	0.004
Triglycerides, mg/dL	101.64(71.66)	116.30(72.38)	0.017

Data are presented as mean (SD), or number (proportion)

BMI, body mass index; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol

play an important role in the lipid metabolism [16], especially the atherogenic lipid components, such as LDL-C and remnant cholesterol [36, 37]. Similarly, the oral-derived systematic inflammatory is associated with glycemic control, complications, and incidence of diabetes, which have been reported in previous studies [38–40].

In our study, reported edentulism was positively associated with LDL-C, nonHDL-C, remnant cholesterol, total cholesterol and triglycerides, and negatively associated with HDL-C. Similarly, Meisel et al. [41] showed that statins use and reduction in LDL-C were associated with diminished tooth loss as a long-term response, which also supported the interaction effect between tooth loss or edentulism and lipid metabolism.

There were some limitations in this study. First, our study failed to collect the dietary information, as the dietary fat and cholesterol intake have an effect on plasma lipid profiles. Second, oral hygiene indicators, such as tooth brushing frequency were not considered in this current study, which could affect the local inflammatory response and lipid metabolism. Third, our study was a cross-sectional observational study, and we could not claim the causal association between edentulism and lipid parameters. Whether oral hygiene improvement could improve lipid level or lipid control could prevent edentulism remain to be investigated in further studies. Fourth, although there was a statistically significant association, the effect size was weak from the clinical perspective, indicating oral health as a systemic indicator





Fig. 2 Distribution of lipid parameters in the whole and matched population

Table 3 Association between lipid profiles and tooth loss in the whole population

	Unadjusted model		Adjusted model	
	OR (95% CI)	P value	OR (95% CI)	P value
LDL cholesterol	1.005(1.001–1.008)	0.014	1.004(1.001–1.008)	0.046
HDL cholesterol	1.001(0.992–1.008)	0.911	1.001(0.992–1.010)	0.838
Remnant cholesterol	0.994(0.981–1.006)	0.393	0.992(0.976–1.006)	0.302
nonHDL cholesterol	1.003(0.999–1.006)	0.099	1.002(0.999–1.006)	0.211
Total cholesterol	1.003(1.000–1.006)	0.092	1.003(0.999–1.006)	0.187
Triglycerides	0.999(0.997–1.001)	0.241	0.999(0.996–1.002)	0.172

OR, odds ratio; CI, confidence interval

Age, sex, education level, marriage status, smoking status, drinking, BMI level, hypertension or not, dyslipidemia or not, diabetes or not were adjusted in model



of many health indicators. In the matched population, there were significant associations of LDL-C, HDL-C, nonHDL-C, remnant cholesterol, total cholesterol and triglycerides with reported edentulism. In the whole population, only LDL-C was significantly associated with reported edentulism probably due to the confounding bias. The effect size of lipid profile on edentulism needs further investigation in other studies.

Conclusions

In conclusion, lipid profiles are probably associated with edentulism in the elder population. The interaction between oral condition and metabolic health needs further attention in clinical practice.

Abbreviations

CHARLS: China Health and Retirement Longitudinal Study; LDL-C: Low-density lipoprotein cholesterol; HDL-C: High-density lipoprotein cholesterol; BMI: Body mass index; SD: Standard deviation; CRP: C-reactive protein; IL-6: Interleukin-6.

Acknowledgements

Not applicable.

Author contributions

YTW, XW and MS conceptualized and developed the study, reviewing and interpreting the results, and editing of the manuscript. SPW and RYY were responsible for data analysis, development of tables, and writing first draft of the manuscript. DXY, YFN and LXW helped with statistical analyses, reviewed and interpreted the results. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The CHARLS study was approved by the institutional review board of Peking University (IRB00001052-11015). All procedures were performed in accordance with the 1964 Declaration of Helsinki and its later amendments or with comparable ethical standards. All participants have provided the informed consents.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Stomatology, Beijing Shijitan Hospital, Capital Medical University, Beijing, China. ²Shanghai Fufan Information Technology Co., Ltd, No.323 Guoding Road, Yangpu District, Shanghai 200433, China. ³Community Health Service Center, Huangcun Town, Daxing District, Beijing, China. ⁴Department of VIP Dental Service, Beijing Stomatological Hospital, Capital Medical University, Beijing, China.

Received: 5 May 2022 Accepted: 4 October 2022

Published online: 15 October 2022

References

1. Peres MA, Macpherson LMD, Weyant RJ, Daly B, Venturelli R, Mathur MR, Listl S, Celeste RK, Guarnizo-Herreño CC, Kearns C, et al. Oral diseases: a global public health challenge. *Lancet*. 2019;394(10194):249–60.
2. Peres MA, Macpherson LMD, Weyant RJ, et al. Oral diseases: a global public health challenge. *Lancet*. 2019;394(10194):249–60.
3. Kassebaum NJ, Bernabé E, Dahiya M, Bhandari B, Murray CJ, Marcenes W. Global burden of severe tooth loss: a systematic review and meta-analysis. *J Dent Res*. 2014;93(7 Suppl):20s–8s.
4. Müller F, Naharro M, Carlsson GE. What are the prevalence and incidence of tooth loss in the adult and elderly population in Europe? *Clin Oral Implants Res*. 2007;18(Suppl 3):2–14.
5. Vallim AC, Gaio EJ, Oppermann RV, Rösing CK, Albandar JM, Susin C, Haas AN. Obesity as a risk factor for tooth loss over 5 years: a population-based cohort study. *J Clin Periodontol*. 2021;48(1):14–23.

6. Genco RJ, Borgnakke WS. Diabetes as a potential risk for periodontitis: association studies. *Periodontol* 2000. 2020;83(1):40–5.
7. Fang WL, Jiang MJ, Gu BB, Wei YM, Fan SN, Liao W, Zheng YQ, Liao SW, Xiong Y, Li Y, et al. Tooth loss as a risk factor for dementia: systematic review and meta-analysis of 21 observational studies. *BMC Psychiatry*. 2018;18(1):345.
8. Thomson WM, Barak Y. Tooth loss and dementia: a critical examination. *J Dent Res*. 2021;100(3):226–31.
9. Dwivedi N, Wiener RC, Findley PA, Shen C, Sambamoorthi U. Asthma, chronic obstructive pulmonary disease, tooth loss, and edentulism among adults in the United States: 2016 Behavioral Risk Factor Surveillance System survey. *J Am Dent Assoc*. 2020;151(10):735–744.e731.
10. Chen Y, Yang YC, Zhu BL, Wu CC, Lin RF, Zhang X. Association between periodontal disease, tooth loss and liver diseases risk. *J Clin Periodontol*. 2020;47(9):1053–63.
11. Chen Y, Zhu BL, Wu CC, Lin RF, Zhang X. Periodontal disease and tooth loss are associated with lung cancer risk. *Biomed Res Int*. 2020;2020:5107696.
12. Michaud DS, Fu Z, Shi J, Chung M. Periodontal disease, tooth loss, and cancer risk. *Epidemiol Rev*. 2017;39(1):49–58.
13. Chen P, Zhang M, Zhang Y, Su X, Chen J, Xu B, Tao J, Wang Z, Ma A, Li H. Economic burden of myocardial infarction combined with Dyslipidemia. *Front Public Health*. 2021;9: 648172.
14. Pirillo A, Casula M, Olmastroni E, Norata GD, Catapano AL. Global epidemiology of dyslipidaemias. *Nat Rev Cardiol*. 2021;18(10):689–700.
15. Khovidhunkit W, Kim MS, Memon RA, Shigenaga JK, Moser AH, Feingold KR, Grunfeld C. Effects of infection and inflammation on lipid and lipoprotein metabolism: mechanisms and consequences to the host. *J Lipid Res*. 2004;45(7):1169–96.
16. He J, Zhang P, Shen L, Niu L, Tan Y, Chen L, Zhao Y, Bai L, Hao X, Li X, et al. Short-chain fatty acids and their association with signalling pathways in inflammation, glucose and lipid metabolism. *Int J Mol Sci*. 2020. <https://doi.org/10.3390/ijms21176356>.
17. Zhang C, Wang K, Yang L, Liu R, Chu Y, Qin X, Yang P, Yu H. Lipid metabolism in inflammation-related diseases. *Analyst*. 2018;143(19):4526–36.
18. Singhrao SK, Harding A, Simmons T, Robinson S, Kesavalu L, Crean S. Oral inflammation, tooth loss, risk factors, and association with progression of Alzheimer's disease. *J Alzheimers Dis*. 2014;42(3):723–37.
19. Song TJ, Kim JW, Kim J. Oral health and changes in lipid profile: a nationwide cohort study. *J Clin Periodontol*. 2020;47(12):1437–45.
20. Zhou BF. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults—study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci*. 2002;15(1):83–96.
21. Jaramillo A, Lafaurie GI, Millán LV, Ardila CM, Duque A, Novoa C, López D, Contreras A. Association between periodontal disease and plasma levels of cholesterol and triglycerides. *Colomb Med (Cali)*. 2013;44(2):80–6.
22. Lösche W, Karapetow F, Pohl A, Pohl C, Kocher T. Plasma lipid and blood glucose levels in patients with destructive periodontal disease. *J Clin Periodontol*. 2000;27(8):537–41.
23. Thapa S, Wei F. Association between high serum total cholesterol and periodontitis: National Health and Nutrition Examination Survey 2011 to 2012 Study of American Adults. *J Periodontol*. 2016;87(11):1286–94.
24. Penumarthi S, Penmetsa GS, Mannem S. Assessment of serum levels of triglycerides, total cholesterol, high-density lipoprotein cholesterol, and low-density lipoprotein cholesterol in periodontitis patients. *J Indian Soc Periodontol*. 2013;17(1):30–5.
25. Taleghani F, Shamaei M, Shamaei M. Association between chronic periodontitis and serum lipid levels. *Acta Med Iran*. 2010;48(1):47–50.
26. Machado AC, Quirino MR, Nascimento LF. Relation between chronic periodontal disease and plasmatic levels of triglycerides, total cholesterol and fractions. *Braz Oral Res*. 2005;19(4):284–9.
27. Kim SW, Cho KH, Han KD, Roh YK, Song IS, Kim YH. Tooth loss and metabolic syndrome in South Korea: The 2012 Korean National Health and Nutrition Examination Survey. *Medicine (Baltimore)*. 2016;95(16): e3331.
28. Kinane DF, Marshall GJ. Periodontal manifestations of systemic disease. *Aust Dent J*. 2001;46(1):2–12.
29. Petersen PE, Ogawa H. The global burden of periodontal disease: towards integration with chronic disease prevention and control. *Periodontol* 2000. 2012;60(1):15–39.
30. Nazir MA. Prevalence of periodontal disease, its association with systemic diseases and prevention. *Int J Health Sci (Qassim)*. 2017;11(2):72–80.
31. Loos BG. Systemic markers of inflammation in periodontitis. *J Periodontol*. 2005;76(11 Suppl):2106–15.
32. Pink C, Kocher T, Meisel P, Dörr M, Markus MR, Jablonowski L, Grotevendt A, Nauck M, Holtfreter B. Longitudinal effects of systemic inflammation markers on periodontitis. *J Clin Periodontol*. 2015;42(11):988–97.
33. Shi D, Meng H, Xu L, Zhang L, Chen Z, Feng X, Lu R, Sun X, Ren X. Systemic inflammation markers in patients with aggressive periodontitis: a pilot study. *J Periodontol*. 2008;79(12):2340–6.
34. Cairo F, Nieri M, Gori AM, Tonelli P, Branchi R, Castellani S, Abbate R, Pini-Prato GP. Markers of systemic inflammation in periodontal patients: chronic versus aggressive periodontitis. An explorative cross-sectional study. *Eur J Oral Implantol*. 2010;3(2):147–53.
35. D'Aiuto F, Parkar M, Andreou G, Suvan J, Brett PM, Ready D, Tonetti MS. Periodontitis and systemic inflammation: control of the local infection is associated with a reduction in serum inflammatory markers. *J Dent Res*. 2004;83(2):156–60.
36. D'Aiuto F, Ready D, Tonetti MS. Periodontal disease and C-reactive protein-associated cardiovascular risk. *J Periodontol Res*. 2004;39(4):236–41.
37. Mustapha IZ, Debrey S, Oladubu M, Ugarte R. Markers of systemic bacterial exposure in periodontal disease and cardiovascular disease risk: a systematic review and meta-analysis. *J Periodontol*. 2007;78(12):2289–302.
38. Genco RJ, Graziani F, Hasturk H. Effects of periodontal disease on glycemic control, complications, and incidence of diabetes mellitus. *Periodontol* 2000. 2020;83(1):59–65.
39. López-Gómez SA, González-López BS, Scougall-Vilchis RJ, Pontigo-Loyola AP, Márquez-Corona ML, Villalobos-Rodelo JJ, Rueda-Ibarra V, Medina-Solís CE. Tooth loss in patients with and without diabetes: a large-scale, cross-sectional study of Mexican adults. *J Am Dent Assoc*. 2020;151(4):276–86.
40. Helal O, Göstemeyer G, Krois J, Fawzy El Sayed K, Graetz C, Schwendicke F. Predictors for tooth loss in periodontitis patients: Systematic review and meta-analysis. *J Clin Periodontol*. 2019;46(7):699–712.
41. Meisel P, Kroemer HK, Nauck M, Holtfreter B, Kocher T. Tooth loss, periodontitis, and statins in a population-based follow-up study. *J Periodontol*. 2014;85(6):e160–168.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

