

DETECTING AND QUANTIFYING LAND COVER AND LAND USE CHANGE IN EASTERN MAU BY REMOTE SENSING

Kundu P.M¹, China S.S, Chemelil M.C and Onyando J.O

Department of Agricultural Engineering, Egerton University, Box 536 Njoro, Kenya

Abstract

Eastern Mau is an important water catchment area for lake Nakuru drainage basin. It has great potential for agricultural development as manifested by the high yields of wheat, barley and dairy products from the few remaining large scale farms. This area was once covered by rich vegetation of highland evergreen forests which extended from the Mau hills and turned into woodland dominated by acacia trees in the Rongai-Njoro plains. Remote sensing techniques and ground survey methods were used to evaluate the land cover /land use change that occurred in the area over a period of about 40 years since 1964. A significant change of land use from forest to agriculture was evident from the interpreted results of the photos taken in 1964 and 1969, the SPOT image of 1987, the Landsat image of 1989 and the ground survey of 2003. Deforestation, land fragmentation, cultivation of wetlands and rapid increase in human settlements have had negative impacts on water resources resulting in reduced stream flows and ground water. Land use has changed from planted forests and large scale farms of wheat, barley, and dairy to homesteads, kitchen gardens and small farms of maize, beans, potatoes and vegetables with some people keeping one or two dairy animals. Currently, studies show increasing land degradation in the area. Sheet and rill erosion is observed when it rains and gullies are beginning to develop along some of the farm tracks. To contain the situation, satellite remote sensing was used to monitor and provide data that can be used to evaluate and manage land use and land use change for sustainable development in Mau East.

Keywords: Remote Sensing, Air photo-interpretation, Landsat, SPOT, Catchment

Introduction

Eastern Mau is in the Rift Valley Province of Kenya and lies between Rongai-Njoro plains and the upper slopes of the Mau hills. It is mainly covered by the Quaternary and the Tertiary volcanic deposits. The area is drained mainly by rivers Njoro and Lamriak which originate from Eastern Mau escarpment and empty into Lake Nakuru and rivers kaumara, Nessuit and Rongai which drain northwards to lake Baringo and Bogoria.

Remotely sensed data and ground survey methods were used to evaluate the land cover /land use change in the area for a period of about 40 years. A combination of fine, medium and coarse resolution images from different platforms were used to detect changes. The study showed a significant change of land use from forest and woodlands to agriculture, and rural built-up lands. This was evident from the analysed results for the air photos of 1969, the SPOT image of 1987, the Landsat images of 1984 and 1989 and the ground survey of 2003 which gave a chronological landscape record. Deforestation, land fragmentation, cultivation of wetlands and rapid increase in human settlements have had negative impacts on water sources resulting in reduced stream flows. Land use has changed from planted forests, large farms of wheat, barley, and dairy to homesteads, kitchen gardens and small farms of maize, beans, potatoes and vegetables with one or two dairy animals.

Methodology

A systematic sequence of API was done on air photographs of 1969 to ensure that definite, reliable and significant information was extracted. The central photograph was analysed using mirror stereoscopes. A full scene SPOT image (fig.5) covering Lake Nakuru and Lake Naivasha drainage basins was manually interpreted and analysed. The general character as well as details of the geology, drainage, plant cover and land use were drawn from surrogate indicators and interpretation elements in the images. Vegetation cover was determined by the colour tone, the distribution, form and texture of plant-covered areas. Land cover change was determined by comparing multi-temporal images. A subscene Landsat image of 1989 (fig.8) covering the recharge areas and the mature sections of the rivers originating from Mau hills were interpreted. The large area covered in one image was important for various kinds of regional studies with details of size 30-80 m being observed. TM bands 4, 5, 3 were selected as they showed vegetation and land use features more clearly.

Ground survey was carried out in the selected sample sites in 2003. In order to achieve unbiased samples, surveys were based on an unaligned systematic random sample of fixed-size segments of 1 km x 1 km. Individual land parcels and ground cover classes were identified in each sample segment. By the method of direct expansion, the area for each cover class was determined for the entire study area. A high densification of sample sites was preferred in order to validate the consistency of land use cover database.

In order to classify the heterogeneous land use in eastern Mau, a simplified method known as EU-LUCAES based on Anderson 1977 was applied. Field survey provided the locations of known land use cover types such as wheat farms and improved pasture fields. It provided the locations of unknown features such as rural built-up lands, heterogeneous crop and subsistence farms which cannot be identified on the image using visual interpretation techniques. It helped

identify the main physical environments of the study area such as hills, valleys and rolling land. Data verification process involved rechecking photographs and images to determine if the interpretation was correct, and where questions existed, the site was field verified. At least a third of the segments were field checked, results tabulated and compared to the original interpretation.

Area by direct expansion

Mean proportion:

$$\bar{y}_c = 1/n \sum_{i=1}^n y_i \quad (1)$$

Where,

\bar{y}_c = Mean proportion of each coverage area

y_i = Coverage proportion for each segment

n = number of segments in the sample

Total area under coverages

$$Z_c = D\bar{y}_c \quad (2)$$

Where,

Z_c = Estimate of the coverages area

D = Total study area

Variances of area estimator

$$Var(\bar{y}_c) = (1 - n/N) / (n(n-1)) \sum_{i=1}^n (y_i - \bar{y}_c)^2 \quad (3)$$

$$Var(Z_c) = D^2 Var(\bar{y}_c) \quad (4)$$

Where,

N = Total number of segments in the study area

Results and Discussion

The land cover classes identified in the area were as shown in table 1.

Table 1: Land cover classes

Category	Air Photograph (1969)	Lands (1969)
		(i)

parts of the image. The bright yellowish areas have high reflectance in all wave-lengths. These are areas of open pasture and grassland, bare soil, overgrazed, just harvested, roads or built-up areas. Patches of dark red represent trees or scrub vegetation. In agricultural lands, the bright tones are an indication of cleared or fallow fields at this time of the year in January. The greenish tones are ploughed fields. Water bodies like lake Nakuru, Lake Elmentaita, Lake Bogoria and Sasumua dam are indicated in black tones. Lake Baringo appears as bright green because of high sedimentation. At the middle of Lake Nakuru, the bright red indicates aquatic vegetation. Woodlands in the image appear as red mottled tones.

Landsat TM (July 1989)

Land use cover change was detected on the image after analyzing it and comparing the results with the photograph of 1969. The colour key shows a significant change of land use from the dominant forest and large scale farms seen on the photograph. It can be seen from figure 8 that the forest which was at the confluence of the little shuru and enjoro has been cleared. The blue patches within the forest indicate new-clear lands now under subsistence cultivation.

Configuration and Composition of Land cover/Land Use

It was ascertained through remote sensing and time series analysis that there has been a significant land use cover change, particularly the conversion of forest and woodlands to arable land as shown in table 2 and figures 1 and 2. The air photographs of 1969 showed large scale farming, forest conservation and little built-up lands. The pre-change geo-topes of planted forests and farms showed clear corridors, sizes, shapes and connectivity. The dominant land use cover types were forests (75%), woodlands (12%) and farms (13%). By

1989, the landscape had changed tremendously as showed by the landsat sub-scene image which showed 60 % forest and woodland and 40 % agriculture and built-up area. The dominance of mixed agriculture and subsistence farming in much of the former Nessuiet forestlands and Ngondu large scale farm is shown by the different shades of blue and whitish patches. Due to decrease of spatial resolution on the images, the dominant land use cover types of forest and mixed farming exhibit a rapid increase and overestimation at the coarse levels of spatial aggregation at the expense of non-dominant land use cover types like rural built-up lands. The dominant land use cover types are overestimated due to spatial mixing of covers hence the high variances. Settlements like Ngondu and Njokerio now stand on former large scale farms and grasslands. Ngondu formerly known as Wright farm produced wheat and dairy products while Njokerio formerly Sebiens farm for horses and flower production. Forest lands in Beeston area have been converted into arable and built-up lands. Expansion into forest and riverine areas has been taking place for quite a while now. From ground survey of 2003, the dominant land cover types were 5% forest, 7% woodland, 82% agriculture and 6% built-up area. It was evident that loss of forest cover in the reserves was by both clear cut and progressive thinning due to poaching by local residents. There are many small land use patches of maize, beans, potatoes, wheat, pasture and homesteads with spectrally similar land cover which tend to clump to the bigger entities contaminated by mixed pixels in remotely sensed data. The EU-LUCAES method used for this study provided a simple calibration which has been validated to assist with interpreting images to determine land use cover types of heterogeneous type, especially subsistence agriculture and rural built-up land on former forests and large scale farms.




Figure 1: Change in woodland




Figure 2: change in arable land

Table 2: Land use cover statistics

Impacts of Land use cover change

As land use changes from forest to rural built-up lands, urban lands and subsistence agriculture, surface runoff increases while surface and groundwater quantity and quality deteriorates. The effect is manifested in reduced natural recharge, reduced stream flow and elimination of wet lands. Currently, there is a steady recognition in the area

the an-
hl

Acknowledgement

I would like to thank the Trust Fund Amsterdam 2000 for giving me the financial assistance to attend the conference and Egerton university for supporting me in the preparation of this paper and giving me time off to attend the conference.

References

Anderson J.R, 1977: Land use and land cover changes-A framework for monitoring: US Geol. Survey, Jour. Research, V.5, (2) pp. 223-224.

Chemelil C.M, 1995: The effects of Human-induced Watershed changes on stream flows. PhD thesis, Loughborough university of Technology

China S.S and Kundu P.M, 1985: Evaluation and mapping of erosion processes on Tatton farm and its surroundings (unpublished)

China S.S, 1993: Land Use Planning using GIS. PhD thesis, University of Southampton

Hoekman D.H and Quinones M.J, 1998: Land cover type and biomass classification using AirSAR data for evaluation of monitoring scenarios in the Colombian Amazon, proc. 2nd Intl workshop on Retrieval of Bio and Geophysical parameters from SAR data for land application, 21-23 Oct. 1998, ESA-ESTEC, Noordwijk, the Netherlands.

Hugh W, Murray J and Peter S, 1995: Land use mapping using satellite data of Manawatu, New Zealand, ITC Juornal, 1996-2

Frank T.D, 1988: Mapping dominant vegetation communities in the Colorado Rocky Mountain Front Range with Landsat Thematic Mapper and digital terrain data. Photogrammetric Engineering and Remote Sensing, 54: 1727-1734.

Gichugi N, 1998: Wetlands coordinator-National Museums of Kenya

Jones R.G.B, 1967: Specific Applications of Air photo Interpretation in agricultural development planning, The journal and proceedings of the Institution of Agricultural engineers Vol. 24 No 2 1969

Karanja A, China S.S, Kundu P.M, 1986: The influence of land use on the Njoro River Catchment between 1975 and 1985. In: Soil and Water Conservation in Kenya. D.B Thomas, E.K Biamah, A.M Kilewe, L. Lundgren, B.O Mochoge (eds) .

Kalyango S.N, 1990: Remote Sensing in assisting weather forecasting and monitoring environmental degradation, Newsletter of the Regional Centre for services in surveying, mapping and Remote Sensing, Nairobi-Kenya, No.1/1991

Kinyanjui H.C.K, 1979: Detailed Soil Survey of Tatton farm, Egerton College

Leprieur C.E, Durand M.J and Peyron J.L, 1988: Influence of topography on forest reflectance using Landsat Thematic Mapper and digital terrain data. Photogrammetric Engineering and Remote Sensing 54: 491-496

Matiru V, 2000: Forest Cover and Forest Reserves in Kenya: Policy and Practice, IUCN Eastern Africa Program: Forest and Social Perspectives in Conservation, Working Paper No.5, January 2000

Megier J and Winkler R, 1996: The use of satellite remote sensing for land cover mapping in Europe. Proceedings, XVIII ISPRS Congress, 21-24 April 1996, Baltimore, MD

Mulders M.A & G. l'Homme, 1997: Multispectral aerial photography as land quality indicator for maize growth in the Limagne area. Proceedings, ITC-ISSS Conference on Geo-information for planning of sustainable Land management, ITC-Land Resources and urban Development Department, Enschede, 17-21 August 1997.

Muthaura F, 2002: Workshop on Population in Nakuru, Ministry of Environment

Njuguna P, Mbegera M, Mbithi D, 1999: Reconnaissance Survey of Forest Blocks in the West and East of Rift Valley, Permanent Presidential Commission on Soil Conservation and Afforestation.

Ottichilo W, 1998: Kenya Wildlife Services (KWS)

Ramesh T, 1998: World Wildlife Fund (WWF)-Lake Nakuru Ramsar Project

Rasch H, 1994: Mapping of vegetation, land cover and land use by satellite: experience and conclusions for future project applications. PE&RS 60, 3 pp 265-271.

Rosenholm, D.G. 1993: Land use and vegetation mapping by satellite: some experiences gained by SSC satellite in 1987-1993, ITC, pp 19-23.

Shivoga W.A 2001: The influence of hydrology on the structure of invertebrate communities in two streams flowing into Lake Nakuru, Kenya, Hydrobiologia 458: 121-130, 2001

Taylor J.C and Eva H.D, 1992: Regional Inventories on Beds, Cambs and Northants (UK), 1992-final report to CEC-JRC, Silsoe College ISBN 1 87156452 2, 70p.