

these, 41.2% were characterized as GII.3 (mostly with a GII.Pb pol), 26.5% as GII. Pg_GII.1, and 17.6% as GII.4 variants. From spring to fall 2012, the variant New Orleans 2009 became the predominant GII.4 strain, and the variant Sydney 2012 strain apparently disappeared.

During November–December 2012 and January 2013, ISGEV detected NoV infection in 90 (28.9%) of 311 children hospitalized for gastroenteritis. This finding is comparable to a prevalence of 25.2% in the same period (November–January) of the 2011–12 winter season. A representative subset of 45 samples was randomly selected for sequencing, and 26 (74.3%) of 35 fully typed strains were characterized as GII.4 Sydney 2012, which suggested that the new NoV variant had become the predominant strain in Italy.

Our surveillance seem to mirror observations of a report from Denmark that documented the onset and circulation at low prevalence of the variant GII.4 Sydney 2012 at the beginning of 2012 with a marked increase in the prevalence only by the end of 2012 (10). Our surveillance detected the emergence of this variant in Italy at the end of 2011 and provided us with one of the earliest strains of the variant GII.4 Sydney 2012. This novel variant has a common ancestor with the NoV GII.4 variants Apeldoorn 2008 and New Orleans 2009 and has several amino acid changes on the main epitope in the capsid P2 domain (10).

Sequence analysis of these early strains of the GII.4 variant Sydney 2012 could help clarify the mechanisms driving its global emergence and spread. Continued surveillance for NoV infections through ISGEV and additional data on clinical and epidemiologic features will enable further assessment of the public health implications of the new variant GII.4 Sydney 2012 in Italy.

This study was supported by the grants “Studio dei meccanismi evolutivi

dei calicivirus umani” (Italian Scientific Research Fund PRIN 2008), “Caratterizzazione molecolare di norovirus circolanti nella popolazione pediatrica” (University of Palermo, Italy, Fondi di Ateneo 2007), and “Epidemiologia molecolare e studio dei meccanismi evolutivi di norovirus” (University of Parma, Italy, Fondi di Ateneo 2008).

**Giovanni M. Giammanco,
Simona De Grazia,
Fabio Tummolo,
Floriana Bonura,
Adriana Calderaro,
Alessio Buonavoglia,
Vito Martella, and
Maria C. Medici**

Author affiliations: Università degli Studi di Palermo, Palermo, Italy (G.M. Giammanco, S. De Grazia, F. Bonura); Università degli Studi di Parma, Parma, Italy (F. Tummolo, A. Calderaro, M.C. Medici); Medical Practitioner, Bari, Italy (A. Buonavoglia); and Università Aldo Moro di Bari, Valenzano, Italy (V. Martella)

DOI: <http://dx.doi.org/10.3201.130119>

References

- Green KY. Caliciviridae. In: Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, et al., editors. *Fields virology*, 5th ed. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 949–79.
- Kroneman A, Vega E, Vennema H, Vinjé J, White PA, Hansman G, et al. Proposal for a unified norovirus nomenclature and genotyping. *Arch Virol*. 2013; April 25. Epub ahead of print.
- Bok K, Abente EJ, Realpe-Quintero M, Mitra T, Sosnovtsev SV, Kapikian AZ, et al. Evolutionary dynamics of GII.4 noroviruses over a 34-year period. *J Virol*. 2009;83:11890–901. <http://dx.doi.org/10.1128/JVI.00864-09>
- Siebenga JJ, Vennema H, Renckens B, de Bruin E, van der Veer B, Siezen RJ, et al. Epochal evolution of GGII.4 norovirus capsid proteins from 1996 to 2006. *J Virol*. 2007; 81:9932–41. <http://dx.doi.org/10.1128/JVI.00674-07>
- van Beek J, Ambert-Balay K, Botteldoorn N, Eden J, Fonager J, Hewitt J, et al. Indications for worldwide increased norovirus activity associated with emergence of a new variant of genotype II.4, late 2012. *Euro Surveill*. 2013;18:8–9.
- Medici MC, Tummolo F, Albonetti V, Abelli LA, Chezzi C, Calderaro A. Molecular detection and epidemiology of astrovirus, bocavirus, and sapovirus in Italian children admitted to hospital with acute gastroenteritis, 2008–2009. *J Med Virol*. 2012;84:643–50. <http://dx.doi.org/10.1002/jmv.23231>
- Giammanco GM, Rotolo V, Medici MC, Tummolo F, Bonura F, Chezzi C, et al. Recombinant norovirus GII.g/GII.12 gastroenteritis in children. *Infect Genet Evol*. 2012;12:169–74. <http://dx.doi.org/10.1016/j.meegid.2011.10.021>
- De Grazia S, Martella V, Chironna M, Bonura F, Tummolo F, Calderaro A, et al. Nationwide surveillance study of human astrovirus infections in an Italian paediatric population. *Epidemiol Infect*. 2013;141:524–8. <http://dx.doi.org/10.1017/S0950268812000945>
- Kroneman A, Vennema H, Deforche K, Avoort HVD, Peñaranda S, Oberste MS, et al. An automated genotyping tool for enteroviruses and noroviruses. *J Clin Virol*. 2011;51:121–5. <http://dx.doi.org/10.1016/j.jcv.2011.03.006>
- Fonager J, Hindbaek L, Fischer T. Rapid emergence and antigenic diversification of the norovirus 2012 Sydney variant in Denmark, October to December, 2012. *Euro Surveill*. 2013;18:pii: 20413.

Address for correspondence: Giovanni M. Giammanco, Dipartimento di Scienze per la Promozione della Salute e Materno Infantile “G. D’Alessandro,” Via del Vespro 133, 90127, Palermo, Italy; email: giovanni.giammanco@unipa.it

Group C Betacoronavirus in Bat Guano Fertilizer, Thailand

To the Editor: Bats play a critical role in the transmission and origin of zoonotic diseases, primarily viral zoonoses associated with high case-fatality rates, including those caused by Nipah virus (NiV) and severe acute respiratory syndrome (SARS)-like coronavirus (CoV) infections (1). Recently, the World Health Organization (WHO) reported 44 confirmed cases of human infection with Middle East

respiratory syndrome CoV, resulting in 22 deaths. Full-genome and phylogenetic analyses of these Middle East respiratory syndrome CoVs have been published elsewhere (2). The identified viruses from 2 patients (previously referred to as England/Qatar/2012 and EMC/2012) are genetically related and belong to group C betacoronavirus, which is most related to CoVs from *Nycteris* bats in Ghana and *Pipistrellus* bats in Europe (2,3). In addition, bat CoVs HKU4 and HKU5 originated from *Tylonycteris pachypus* and *Pipistrellus abramus* bats, respectively, in the People's Republic of China (4). Bats are also known to harbor and transmit nonviral zoonotic pathogens, including the fungal pathogen *Histoplasma capsulatum*, which causes histoplasmosis in humans (5).

Bat guano is sold for use as a fertilizer in several countries, including Thailand, Indonesia, Mexico, Cuba, and Jamaica. The practice of collecting and harvesting bat guano may pose a considerable health risk because guano miners have a high level of contact and potential exposure to bat-borne pathogens. To assess pathogens in bat guano, we examined bat guano from a cave in the Khao Chong Phran Non-hunting Area (KCP-NHA) in Ratchaburi Province, Thailand, where bat guano was sold as agricultural fertilizer, for the presence of NiV, CoV, and *H. capsulatum* fungi. Bats from 14 species in 7 families have been found roosting within this area. *Tadarida plicata* bats are the most abundant species (2,500,000 bats), and 3 other species of bats found at the site each had thousands of members: *Taphozous melanopogon*, *Taphozous theobaldi*, and *Hipposideros larvatus*.

A random sample of dry bat guano, ≈ 100 g, was collected in a sterile plastic bag weekly from the main cave at KCP-NHA from September 2006 through August 2007. The specimens were sent for analysis by express mail

(at room temperature within 2–3 days) to the WHO Collaborating Centre for Research and Training in the Viral Zoonoses Laboratory at Chulalongkorn University. Samples were frozen immediately at -80°C until nucleic acids were extracted and PCR assays were run. A total of 52 collected bat guano specimens were examined in this study.

Two aliquots of feces from each weekly specimen (104 samples total) were screened for CoV, NiV, and *H. capsulatum* by PCR. RNA was extracted from 10 mg of fecal pellet by using the QIAamp Viral RNA Mini Kit (QIAGEN, Hilden, Germany). CoV RNA was detected by using nested reverse transcription PCR with the degenerated primers to amplify the RNA-dependent RNA polymerase (*RdRp*) gene (6). NiV RNA was detected by duplex nested reverse transcription PCR (7). To detect *H. capsulatum* and other fungi, we extracted genomic DNA directly from bat guano by using the silica-guanidine thiocyanate protocol, NucliSense Isolation Reagent (bioMérieux, Boxtel, the Netherlands), according to the manufacturer's protocol. We tested for fungal ribosomal DNA (rDNA) in extracted total nucleic acid specimens by using the PCR protocol designed to amplify all rDNA from 4 major fungus phyla at the internal transcribed spacer 1 and 2 regions (8).

Four (3.8%) of 104 samples were positive for CoV. They were collected on September 2, 2006 (KCP9), October 26, 2006 (KCP12), November 14, 2006 (KCP15), and March 4, 2007 (KCP31). Three of the 4 positive CoV sequences (KCP9, KCP12, and KCP15) were identical at 152 nt of the *RdRp* region (ATCGTGCTATGCCTAATATGTGTAGGATTTTGCATCTCTCATATTAGCTCGTAAACACAATACTTGTTGTAGTGTTCAGACCGCTtTtATAGACTTGCaAACGAGTGTGCGCAAGTCTTGAGTGAGTATGTGCTATGTGGTGGTGGCTAT) and phylogenetically clustered with

the group C betacoronavirus (Figure), with 76%, 80%, and 77% nt identity to bat CoV HKU4, bat CoV HKU5, and human CoV EMC and England1_CoV, respectively. The other CoV sequence (KCP31: ATCGTGCACTTCCCAATATGATACGCATGATTTCCGCCATGATTTTGGGATCAAAGCATGTACTTGCTGTGACACATCTGATAAGTATTACCGTCTTTGTAATGAGCTtGCACAAGTTTTCAGAGGTTGTTTATTCTAATGGTGGTTTC) showed 82% nt identity with bat CoV HKU8, an alphacoronavirus. Although we recognize that longer sequences or full genomes may alter the topology of the phylogeny slightly and give stronger branch support, we expect that the overall topology and placement of these CoVs would remain consistent. Samples from particular bat species could not be identified because bats of different species roost in this cave, and samples were pooled during collection for bat guano fertilizer. The detection of CoVs in bat guano from the KCP-NHA cave in Ratchaburi was consistent with the previous finding of alphacoronavirus from *Hipposideros armiger* bats from the same province in 2007, but those researchers tested fresh bat feces (9).

All bat guano samples screened by PCR were negative for NiV and *Histoplasma* spp. but were positive for group C betacoronavirus. The natural reservoir and complete geographic distribution of this CoV are currently unknown. Although we did not isolate live virus from these samples, the detection of nucleic acid and previous isolation of viruses from bat feces and urine (10) warrants some concern that guano miners might be exposed to bat pathogens in fresh excreta as well as in soil substances. We suggest that guano miners use preventive measures of personal hygiene and improved barrier protection to reduce the possibility of exposure to zoonotic pathogens.

This research was supported by the Thailand Research Fund (DBG5180026, RDG5420089); the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission (HR1160A-55); the Thai Red Cross Society, Thailand; the Naval Health Research Center BAA-10-93 under Cooperative Agreement no. W911NF-11-2-004; the Centre National de la Recherche Scientifique; a nonbiodefense Emerging Infectious Diseases Research Opportunities Award from the National Institute of Allergy and Infectious Diseases of the National Institutes of Health (1 R01 AI079231-01); and the United States Agency for International Development Emerging Pandemic Threats Program, PREDICT project, under the terms of Cooperative Agreement no. GHN-A-OO-09-00010-00.

**Supaporn Wacharapluesadee,
Chirapol Sintunawa,
Thongchai Kaewpom,
Kritsada Khongnomnan,
Kevin J. Olival,
Jonathan H. Epstein,
Apaporn Rodpan,
Paiboon Sangsri, Nirun Intarut,
Ariya Chindamporn,
Kanyarat Suksawa, and
Thiravat Hemachudha**

Author affiliations: Chulalongkorn University and King Chulalongkorn Memorial Hospital, Bangkok, Thailand (S. Wacharapluesadee, T. Kaewpom, K. Khongnomnan, A. Rodpan, N. Intarut, A. Chindamporn, K. Suksawa, T. Hemachudha); Mahidol University, Nakornpathom, Thailand (C. Sintunawa); EcoHealth Alliance, New York, New York, USA (K.J. Olival, J.H. Epstein); and Ministry of Natural Resources and Environment, Bangkok (P. Sangsri)

DOI: <http://dx.doi.org/10.3201/eid1908.130119>

References

- Olival KJ, Epstein JH, Wang LF, Field HE, Daszak P. Are bats unique viral reservoirs? In: Aguirre AA, Ostfeld RS, Daszak P, editors. *New directions in conservation medicine: applied cases of ecological health*. 2nd ed. Oxford: Oxford University Press; 2012. p. 195–212.
- Cotten M, Lam TT, Watson SJ, Palser AL, Petrova V, Grant P, et al. Full-genome deep sequencing and phylogenetic analysis of novel human betacoronavirus. *Emerg Infect Dis*. 2013;19:736–42. <http://dx.doi.org/10.3201/eid1905.130057>
- Annan A, Baldwin HJ, Corman VM, Klose SM, Owusu M, Nkrumah EE, et al. Human betacoronavirus 2c EMC/2012-related viruses in bats, Ghana and Europe. *Emerg Infect Dis*. 2013;19:456–9. <http://dx.doi.org/10.3201/eid1903.121503>
- Woo PC, Lau SK, Li KS, Poon RW, Wong BH, Tsoi HW, et al. Molecular diversity of coronaviruses in bats. *Virology*. 2006;351:180–7. <http://dx.doi.org/10.1016/j.virol.2006.02.041>
- Lyon GM, Bravo AV, Espino A, Lindsley MD, Gutierrez RE, Rodriguez I, et al. Histoplasmosis associated with exploring a bat-inhabited cave in Costa Rica, 1998–1999. *Am J Trop Med Hyg*. 2004;70:438–42.
- Tong S, Conrardy C, Ruone S, Kuzmin IV, Guo X, Tao Y, et al. Detection of novel SARS-like and other coronaviruses in bats from Kenya. *Emerg Infect Dis*. 2009;15:482–5. <http://dx.doi.org/10.3201/eid1503.081013>
- Wacharapluesadee S, Hemachudha T. Duplex nested RT-PCR for detection of Nipah virus RNA from urine specimens of bats. *J Virol Methods*. 2007;141:97–101. <http://dx.doi.org/10.1016/j.jviromet.2006.11.023>
- White TJ, Bruns T, Lee S, Taylor JW. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. *PCR protocols: a guide to methods and applications*. New York: Academic Press, Inc.; 1990. p. 315–22.
- Gouilh MA, Puechmaille SJ, Gonzalez JP, Teeling E, Kittayapong P, Manuguerra JC. SARS-coronavirus ancestor's foot-prints in South-East Asian bat colonies and the refuge theory. *Infect Genet Evol*. 2011;11:1690–702. <http://dx.doi.org/10.1016/j.meegid.2011.06.021>
- Chua KB, Koh CL, Hooi PS, Wee KF, Khong JH, Chua BH, et al. Isolation of Nipah virus from Malaysian Island flying-foxes. *Microbes Infect*. 2002;4:145–51. [http://dx.doi.org/10.1016/S1286-4579\(01\)01522-2](http://dx.doi.org/10.1016/S1286-4579(01)01522-2)

Address for correspondence: Supaporn Wacharapluesadee, WHO Collaborating Centre for Research and Training on Viral Zoonoses, Faculty of Medicine, Chulalongkorn University, Bangkok, 10330, Thailand; email: spwa@hotmail.com




Figure. Phylogenetic tree of 3 coronaviruses (CoVs) isolated from bat guano collected in this study (KCP9, KCP12, and KCP15); 19 additional human and animal CoVs from the National Center for Biotechnology Information database are included. Construction of the tree was based on 152 nt of the *RNA-dependent RNA polymerase* gene region by maximum-likelihood method and GTR+I model with the 1,000 bootstrap resampling method implemented in MEGA5 (<http://megasoftware.net/>). Numbers on branches indicate percentages of bootstrap support from 1,000 replicates. The scale bar indicates the estimated 0.1 nt substitutions per site. HCoV-HKU1C, human CoV HKU1C (DQ415913); HCoV-HKU1A, human CoV HKU1A (DQ415903); HCoV-HKU1B, human CoV HKU1B (AY884001); MHV, murine hepatitis virus (NC001846); HCoV-OC43, human CoV OC43 (AY585229); PHEV, porcine hemagglutinating encephalomyelitis virus (DQ011855); BCoV, bovine CoV (AF391541); BatCoV-HKU9, *Rousettus* bat CoV HKU9 (NC009021); BatCoV/KW2E-F93/Nyc, *Nycteris* bat CoV (JX899383); BatCoV/KW2E-F53/Nyc, *Nycteris* bat CoV (JX899384); BatCoV/KW2E-F82/Nyc, *Nycteris* bat CoV (JX899382); BatCoV HKU4, *Tylonycteris* bat CoV HKU4 (NC009019); BatCoV HKU5, *Pipistrellus* bat CoV HKU5 (NC_009020); HCoV-EMC, human betacoronavirus 2c EMC/2012 (JX869059); England1_CoV, human betacoronavirus England 1 (NC_019843); SARS_BatCoV HKU3, severe acute respiratory syndrome (SARS)-related *Rhinolophus* bat CoV HKU3 (DQ022305); SARSr-CoV-CFB, SARS-related Chinese ferret badger CoV (AY545919); SARSr-ciCoV, SARS-related palm civet CoV (AY304488); SARS_huSARS-CoV, SARS human CoV (NC_004718). An expanded version is online at wwwnc.cdc.gov/EID/article/19/8/13-0119-F1.htm