

Special Considerations for Prophylaxis for and Treatment of Anthrax in Pregnant and Postpartum Women

Technical Appendix

Technical Appendix Table 1. Oral Antimicrobial Post-Exposure Prophylaxis for infection with *Bacillus anthracis**

a. For all strains, regardless of penicillin susceptibility or if susceptibility is unknown	
Non-pregnant Adults:	Modifications for Pregnant Women:
ciprofloxacin 500 mg every 12H OR doxycycline 100 mg every 12H OR levofloxacin 750 mg every 24H OR moxifloxacin 400 mg every 24H OR clindamycin† 600 mg every 8H	ciprofloxacin is preferred no change in dosing
OR	
b. Alternatives for penicillin-susceptible strains	
amoxicillin 1 g every 8H OR penicillin VK 500 mg every 6H	
Duration of Post-Exposure Prophylaxis for <i>Bacillus anthracis</i> : 60 d	no change in duration

***Boldface** indicates preferred agent. Alternative selections are listed in order of preference for treatment for patients who cannot take first-line treatment, or if first-line treatment is unavailable.

†Based on in vitro susceptibility data, rather than studies of clinical efficacy.

Technical Appendix Table 2. Intravenous Antimicrobial Treatment for Systemic Anthrax with Possible/Confirmed Meningitis*

Nonpregnant Adults:	Modifications for pregnant Women:
1. A Bactericidal Agent (Fluoroquinolone) ciprofloxacin 400 mg every 8H OR levofloxacin 750 mg every 24H OR moxifloxacin 400 mg every 24H PLUS	ciprofloxacin is preferred
2. A Bactericidal Agent (β-lactam) a. For all strains, regardless of penicillin susceptibility or if susceptibility is unknown meropenem 2 g every 8H OR imipenem† 1 g every 6H OR doripenem 500 mg every 8H OR	at least one antibiotic with transplacental passage is recommended; ciprofloxacin, levofloxacin, meropenem, ampicillin, penicillin, clindamycin, rifampin
b. Alternatives for penicillin-susceptible strains penicillin G 4 million units every 4H OR ampicillin 3 g every 6H PLUS	
3. A Protein Synthesis Inhibitor linezolid‡ 600 mg every 12H OR clindamycin 900 mg every 8H OR rifampin§ 600 mg every 12H OR chloramphenicol¶ 1 g every 6–8 H	
Duration of treatment: for ≥2–3 weeks until clinical criteria for stability are met. Patients exposed to aerosolized spores will require prophylaxis to complete an antimicrobial drug course of 60 d from onset of illness (see Technical Appendix Table 1).	No change in duration
<p>*Systemic anthrax includes anthrax meningitis; inhalation, injection, gastrointestinal anthrax; and cutaneous anthrax with systemic involvement, extensive edema, or lesions of the head or neck. Boldface indicates preferred agent. Alternative selections are listed in order of preference for treatment for patients who cannot take first-line treatment, or if first-line treatment is unavailable.</p> <p>† Increased risk of seizures associated with imipenem/cilastatin treatment</p> <p>‡Linezolid should be used with caution in patients with thrombocytopenia, as it may exacerbate it. Linezolid use for >14 d carries additional risk for hematopoietic toxicity.</p> <p>§Rifampin is not a protein synthesis inhibitor; however, it may be used in combination with other antimicrobials based on its in vitro synergy.</p> <p>¶Should only be used if other options are not available, due to toxicity concerns.</p>	

Technical Appendix Table 3. Intravenous Antimicrobial Treatment for Systemic Anthrax When Meningitis Has Been Excluded*

Non-pregnant adults	Modifications for pregnant Women:
1. A Bactericidal Antimicrobial	
a. For all strains, regardless of penicillin susceptibility or if susceptibility is unknown	
ciprofloxacin 400 mg every 8H	ciprofloxacin is preferred
OR	
levofloxacin 750 mg every 24H	
OR	
moxifloxacin 400 mg every 24H	
OR	
meropenem 2 g every 8H	
OR	
imipenem† 1 g every 6H	at least one antibiotic with transplacental passage is recommended; ciprofloxacin, levofloxacin, meropenem, ampicillin, penicillin, clindamycin, rifampin
OR	
doripenem 500 mg every 8H	
OR	
vancomycin 60 mg/kg/day IV divided every 8 h (maintain serum trough concentrations of 15 – 20 µg/mL)	
OR	
b. Alternatives for penicillin-susceptible strains	
penicillin G 4 million units every 4H	
OR	
ampicillin 3 g every 6H	
PLUS	
2. A Protein Synthesis Inhibitor	
clindamycin 900 mg every 8H	
OR	
linezolid‡ 600 mg every 12H	
OR	
doxycycline§ 200 mg initially, then 100 mg every 12H	
OR	
rifampin¶ 600 mg every 12H	
Duration of treatment: for ≥2 weeks until clinical criteria for stability are met. Patients exposed to aerosolized spores will require prophylaxis to complete an antimicrobial drug course of 60 d from onset of illness (see Technical Appendix Table 1).	No change in duration
<p>*Systemic anthrax includes anthrax meningitis; inhalation, injection, gastrointestinal anthrax; and cutaneous anthrax with systemic involvement, extensive edema, or lesions of the head or neck. Boldface indicates preferred agent. Alternative selections are listed in order of preference for treatment for patients who cannot take first-line treatment, or if first-line treatment is unavailable. †Increased risk of seizures associated with imipenem/cilastatin treatment</p> <p>‡Linezolid should be used with caution in patients with thrombocytopenia, as it may exacerbate it. Linezolid use for >14 d carries additional risk for hematopoietic toxicity.</p> <p>§A single 10–14 course of doxycycline is not routinely associated with tooth-staining.</p> <p>¶Rifampin is not a protein synthesis inhibitor; however, it may be used in combination with other antimicrobials based on its in vitro synergy.</p>	

Technical Appendix Table 4. Oral Antimicrobial Treatment for Cutaneous Anthrax without Systemic Involvement*

Non-pregnant adults	Modifications for pregnant women
a. For all strains, regardless of penicillin susceptibility or if susceptibility is unknown ciprofloxacin 500 mg every 12H OR doxycycline 100 mg every 12H OR levofloxacin 750 mg every 24H OR moxifloxacin 400 mg every 24H OR clindamycin† 600 mg every 8H	ciprofloxacin is preferred
OR	
b. Alternatives for penicillin-susceptible strains amoxicillin 1 g every 8H OR penicillin VK 500 mg every 6H	
Duration of Treatment: 60 d	No change on duration
*Recommendations are specific to cutaneous anthrax in the setting of bioterrorism. Boldface indicates preferred agent. Alternative selections are listed in order of preference for treatment for patients who cannot take first-line treatment, or if first-line treatment is unavailable.. †Based on in vitro susceptibility data, rather than studies of clinical efficacy.	

Members of the Workgroup on Anthrax in Pregnant and Postpartum Women:

Mahmoud Ahmed University of Texas Medical Branch

Martina Louise Badell Emory University

Kristin Banks Centers for Disease Control and Prevention (CDC)

Wanda Barfield CDC

Michael Bartenfeld CDC

Rich Beigi* Magee-Women's Hospital

Tina Bhavsar* CDC

William Bower* CDC

John Bradley* American Academy of Pediatrics

Deborah Brower Health Resources and Services Administration

Renee Brown-Bryant* CDC

William Callaghan* CDC

Tegan Callahan Association of Maternal & Child Health Programs

Amanda Cohn* CDC

Ava Marie S. Conlin* Naval Health Research Center

Andreea Creanga* CDC

Deborah Dee* CDC

Gary Disbrow Biomedical Advanced Research and Development Authority

Jeffrey Ecker* Massachusetts General Hospital

Jesse Geibe CDC

Alma Gomez Association of Maternal & Child Health Programs

Susan Gorman CDC

Jacqueline Grant* State of Georgia

Violanda Grigorescu CDC

Gary Hankins* University of Texas Medical Branch

Publisher: CDC; Journal: Emerging Infectious Diseases
Article Type: Online Report; Volume: 20; Issue: 2; Year: 2014; Article ID: 13-0611
DOI: 10.3201/eid2002.130611; TOC Head: Online Report
Piia Hanson Association of Maternal & Child Health Programs

Karen E, Hays* American College of Nurse-Midwives

Mary F. Hebert University of Washington

Robert Heine* Duke University

Katherine Hendricks* CDC

Jack Herrmann National Association of County and City Health Officials

Kate Howe* Association of Maternal & Child Health Programs

Mark L. Hudak* University of Florida College of Medicine -Jacksonville

Denise Jamieson* CDC

Ayten Kadanali, Umraniyae EAH, Turkey

Sedat Kadanali, Medical Park Hospital Turkey

Juliette Kendrick* CDC

Hye-Joo Kim* CDC

George Korch Office of the Assistant Secretary for Preparedness and Response

Marian McDonald CDC

Michael M. McNeil* CDC

Dana Meaney-Delman* CDC

Nancy Messonnier* CDC

Lara Misegades* CDC

Mirjana Nesin* National Institutes of Health

Stacey Parker CDC

Alison Patti, CDC

Andrew Pavia* University of Utah

Georgina Peacock* CDC

Roberta Prepas* American College of Nurse Midwives

Conrad P. Quinn* CDC

Lauren Raskin Ramos Association of Maternal & Child Health Programs

Sonja Rasmussen* CDC

Publisher: CDC; Journal: Emerging Infectious Diseases
Article Type: Online Report; Volume: 20; Issue: 2; Year: 2014; Article ID: 13-0611
DOI: 10.3201/eid2002.130611; TOC Head: Online Report
Mirelys Rodriguez CDC

Paula Rosenberg CDC

Catherine Ruhl* Association of Woman's Health, Obstetric and Neonatal Nurses

Mark Sewell U.S. Army

Sean Shadomy* CDC

Jeanne S. Sheffield* University of Texas Southwestern Medical Center

Tom Shimabukuro* CDC

David Siegel National Institutes of Health

Laura Smith CDC

David Soper* Medical University of South Carolina

Helen Stallings Office of the Assistant Secretary for Preparedness and Response

Tracee Treadwell CDC

Methodius Tuuli* Washington University in St. Louis

Etobssie Wako* CDC

Lee Warner CDC

D. Heather Watts* National Institutes of Health

George Wendel Jr.* American Board of Obstetrics and Gynecology

Linda West* CDC

Amy William CDC

Melissa Willis Biomedical Advanced Research and Development Authority

Mary Wright* National Institutes of Health

Yon Yu* CDC

Laurie Zephyrin Department of Veterans Affairs

Marianne Zotti* CDC

*Designates pre-meeting work group participant.